

The Himalayan Gazetteer or

**The Himalayan Districts of the
North Western Province of India**

In 3 Vols. (6 Parts)

Edwin T Atkinson

Vol. 1 in 2 parts

Part - 1

COSMO PUBLICATIONS

New Delhi

PREFACE.

THE system of transliteration used in the present volume is that which has been adopted in the Gazetteer of the North-Western Provinces. For the vowels it is briefly as follows :—

a	as in	'woman.'	a	as in	'pall.'
d	"	'father.'	d	"	'rule.'
e	"	'grey.'	ai	"	'aiaia.'
i	"	'bit.'	o	"	'hole.'
ɪ	"	'machine.'	su	as	'ou' in 'house' (nearly).

The original plan of this Memoir provided for a thorough examination of the country between the Tons and the Kāli somewhat on the plan of Tennent's Ceylon, and in addition a less detailed summary of our knowledge of the Himālaya-Tibetan region. It was believed that only thus could the physical unity and with it the historical connections of the entire region be correctly understood. From or through the Himālaya came the pre-Aryan and Aryan races now inhabiting India, as well as the successive waves of Baktrian, Skythian, and Musalmān invaders who have conquered India in historical times. As we shall see hereafter, forms of belief having their origin in or beyond the Himālaya have influenced the religious systems of India from the earliest ages to the present day, so that for the political and religious history of the plains an adequate conception of the physiography, ethnography, and history of the Himālaya-Tibetan tract is a necessary preparation. It is to be regretted that this extended programme cannot be carried out, but the pressure of official duties prevents its completion, and all that can now be attempted is to work up the materials that have been collected for the Himālaya districts of the North-Western Provinces. At the end of Chapter I. will be found a series of 'References' to materials illustrating the history and resources of the Himālaya from Asām to Afghānistān. These were collected in continuation of a plan suggested by Mr. R. N. Cust in 1866 for the preparation of a 'Catalogue raisonné of every kind of printed information connected with the North-Western Provinces;' and I certainly know of no better aid to good administration than an index to the special and local reports of those engaged in it. The necessity for such a catalogue has ceased in great part with

PREFACE.

the publication of the District Memoirs, but there are subjects of general importance which the publication of the local accounts does not subserve. One of these is the history, using this word in its widest sense, of the Himálaya-Tibetan region, and I offer the references as the nucleus of a complete index to our knowledge of its physiography, products, peoples, and institutions.

As observed in the foot-note to Chapter I., much remains to be done, but at the same time very much more has been accomplished than is generally known. Scientific and economic botany have been carefully explored; the chapter on economic mineralogy leaves little to be desired: those on meteorology and geology have been brought up to the level of our present scientific knowledge; and the introduction to physical geography gives a popular and suggestive summary of the information that we possess, whilst the references afford a guide to materials for the study of details. All this is new, or embodies information buried in correspondence and reports, and practically as inaccessible to the public as if it had never been committed to writing. To General Richard Strachey I am indebted for the use of an unpublished work of his own on the physical geography of the Himálaya, which has been specially made use of in the chapters on 'Geology' and 'Meteorology.' Mr. H. B. Medlicott, Superintendent of the Geological Survey of India, prepared the chapter on 'Geology,' and Mr. S. A. Hill, Meteorological Reporter to the Government of the North-Western Provinces, contributed the valuable chapter on 'Meteorology.' Dr. King, Superintendent of the Royal Botanical Gardens, Calcutta, furnished the list of the *flora* of Western Garhwál, Dehra Dún, and Jaunsár-Báwar, and Dr. Watson, the list for Eastern Garhwál, Kumaon, and the Bhábar. To General Strachey I am further indebted for the list of plants collected by himself and Mr. Winterbottom in Kumaon, Garhwál, and the neighbouring parts of Tibet, and which has never before been published. This list has been admirably edited by Mr. F. Duthie, Superintendent of the Botanical Gardens at Saháranpur, to whom also I am indebted for the sketch of the history of the Tea industry in the Himálayan districts. The sheets of the portions relating to economic botany have had the criticism of Dr. Watson and Mr. Duthie; and for the forest history Mr. Greig and Major Garstin have advised me in many matters of detail.

PREFACE.

Of the maps, the map of Kumaon was prepared under the instructions of Colonel Walker, R.E., Superintendent of the Great Trigonometrical Survey, and is the result of great labour and care for detail : indeed, a less accurate map might possibly be more useful. The great 'one-inch to one-mile' sheets were photographed down to a smaller scale and sent to me. I then obtained the correct names in Hindi of all the villages, rivers, &c., in Kumaon and Garhwál (about 8,000) and entered against each the correct transliteration for use in a new edition of the large maps, and then drew up a draft list of names for the district map. I desired to enter the names of all halting-places on every ordinary route ; villages in which the patwári usually resides ; places of note ; trade-centres ; and all villages containing over one hundred inhabitants. After much trouble, correspondence, and inquiry, this was fairly accomplished. The villages retained were marked off on the photographic copies of the maps, and these were returned to the Surveyor-General with lists showing the actual and revised spelling for the further compilation. In addition, the principal lines of road have been laid down, and for the first time the pattis or subdivisions adopted at the recent settlement by Mr. J. Beckett have been shown. In the trans-Himálayan portion to which the survey had not extended I have added places from the village maps prepared by the patwáris, and in their selection was aided by Mr. Beckett. The map of the Himálaya-Tibetan region is taken from one prepared by Mr. Trelawny Saunders, omitting the eastern portion. The geological map is a revised copy of that which accompanied General Strachey's paper 'On part of the Himálaya Mountains and Tibet' read before the Geological Society in 1851. It may be necessary to explain that these volumes are not intended to be solely a popular account of the districts which they describe, but to contain, first of all, a record of all facts of permanent scientific or economic value that have been gathered by me during my visits to Kumaon and the Dehra Dún or which have been contributed by others. This work has occupied my leisure time for several years, and I can only hope that the labour and care bestowed on it will be of some use to my successors, and enable them to produce a more worthy record for those who are truly interested in the moral and material progress of the North-Western Provinces.

BOMBAY,
17th December, 1881. }

E. T. ATKINSON.

CONTENTS

VOL. I PART I

CHAPTER I.

INTRODUCTION.

	Page.
LIMITATION of this work—Systematic geography—Herbert ; Vigna, Jacquemont, &c. ; Hodgson ; R. Strachey ; Thompson ; Cunningham ; H. Strachey ; Rawlinson ; Saunders ; Markham ; Calcutta Notices ; Blanford—Nomenclature—Ethnical sketch : Plains of Assam and Bengal ; Plains of Upper India ; Clothing ; Food ; Customs ; Eastern Himálaya—Eastern tribes ; Tribes of the Southern watershed ; Bhután, Sikkim, Nepál ; Tribes of the Lower Himálaya—Karnál to Tume ; Bieahr ; the Satlaj to the Indus—Afghánistán ; Galkha States ; Kára-Tangutans ; Tibet—General distribution	1

Refs

CHAPTER II.

[PHYSICAL GEOGRAPHY OF THE HIMÁLAYA.]

The Himálayan river basins—The Indus basin—Subordinate systems of the Indus basin—The Ganges basin—Subordinate systems of the Ganges basin ; The Brahmaputra basin—Subordinate systems of the Brahmaputra basin ; Systems of Tibet—Lob-Nor basin ; Oxus basin. Plains of Hindustán—Indus plain ; Indian desert ; Gangetic plain ; Assam valley ; Bhábar ; Tarál ; Siwálík ; Dána. Extent of Bhábar—Cause of the deposit—Hodgson's oceanic theory ; Fluvialite theory ; Tarál ; Siwálík ; Dána ; Ganges to Brahmaputra ; Lower Himálaya. Materials of the mountains—Eocene period ; Mode of mountain formation ; Mountain-sculpture ; Ice-wedges ; Avalanches ; Glaciers ; Rain ; Rivers	61
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----

CHAPTER III.

GEOLOGY.

Position of the ground. Early observers : Herbert, Cantley, Falconer. The mountain system. The sub-Himálayan some formed of tertiary rocks. Siwálík. Post-tertiary deposits. Supposed glacial deposits. Lower Himálayan region. Simla region. Sirmúr formation. Relation of Sirmúr and Siwálík series. The Satlaj valley. East of Simla. The Kumaon section. The Nepál section. Sikkim section. Slight correspondence between the rocks of the Himálaya and of the Peninsula. Frequent occurrence of carbonaceous deposits throughout the Lower Himálaya. Conjectural affiliation of the Lower Himálayan sections. The Central Himálaya. Strachey. Stolickaa. Western Tibet. The principal crystalline areas. Nummulitic deposits. Post-eocene eruptive rocks. Central gneiss. Kachhar-Kishtwár region. Central Tibet. Its granites, schists, slates, conglomerates and fossiliferous rocks. Palaeozoic strata,

	<i>Page.</i>
Silurian, Trias, Jurassic. Fossils Tertiary deposits. So-called Niti fossils. General Strachey's conclusions. Stolickza's views	... 111

CHAPTER IV.

THE HIMÁLAYA.

The outer Himálaya. Mountain systems; Jumna; Ganges; Káli. Subordinate boundaries The Nayár. Western Rám-ganga; Káli. The snowy range. Axis of highest elevation. The Tibetan plateau. Glaciers. Glacier of the Pindar; feeder of the glacier; moraines; crevasses and ice tables. Glacier of the Kuphinf. Motion of the Pindar glacier. Snow-line. Observations in the eastern Himálaya. Observations in Kumaon: Bisahr Across the snowy range. Kumaon; Garhwál; Kumaon; Western Tibet. Lakes	... 160
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------

CHAPTER V.

METEOROLOGY.

Preliminary sketch of climate. Contrast between the eastern and western parts of the plain and the Himálaya. Radiation, solar and nocturnal. Temperature:—Diurnal and annual ranges,—Vertical decrement,—Height of snow-line. Pressure and winds:—Barometric tides,—Mountain winds,—Annual variation of pressure,—Monsoons. Humidity:—Vertical distribution of vapour,—Relative humidity,—Cloud. Rain-fall:—Distribution on plains and on the Himálayan slope,—Annual variation,—The winter rains,—The monsoon rains	... 201
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------

CHAPTER VI.

ECONOMIC MINERALOGY.

Mining history: experiments in Garhwál. The Kumaon Iron Works Company. Causes of present poor returns—Mode of mining: excavation: removing the ore: crushing the ore: cleaning the ore: drainage. Smelting operations: refining the ore. Division of profits. Improvement needed in appliances: Chilli furnace: English system. Outturn. Financial results of the settlement of the revenue derived from minerals. Gold: Tibetan gold-fields: Thok Jalung mines. Silver. Copper mines: Rái mine and its ores: Sira mines: Gaul and Sor Gurang. Bournonite. Copper mines of Garhwál: Dhanpur: Dhoori: Pokhri: Chaumattiya mines: Rájá's mine: Nola mine: Thála mine: Danda mine: Tálapungia mine: Kharna mine: Agar Sera mine. Iron mines: Kumaon Iron mines: Giwar mines: Garhwál mines: Nágpur: Dasauli: Irlyakot. Lead Arsenic. Lignite. Graphite. Sulphur. Borax. Gypsum. Soap. Stone. Asbestos. Silajit. Limestone. Building stone. Roofing slate. Alum	250
<i>References</i>	297

CHAPTER VII.

SCIENTIFIC BOTANY.

A list of the *flora* of Garhwál, Jaunár-Báwar and the Dehra Dén, by Dr. King. Tract to which the list refers: The Siwálík. Sub Siwálík Tract.

	<i>Page.</i>
Dehra Dún. Himálayan tract. Characteristic vegetation from the plains to Mussooree. From Mussooree northwards. A list of the <i>flora</i> of Kumaon and the Kumaon Bhábar by Dr. Watson	... 299

MAP AND ILLUSTRATIONS.

- 1 Geological map of the country between the Satlaj and the Sarda (page 115).
2. Front view of the Pindar glacier (page 182).
3. The Pindar and Kuphiní glaciers (page 184).

THE HIMÁLAYAN DISTRICTS OF THE NORTH-WESTERN PROVINCES.

PART I.

CHAPTER I.

INTRODUCTORY.

Limitation of this work—Systematic geography—Herbert; Vigne, Jacquemont, &c.; Hodgson; R. Strachey; Thompson; Cunningham; H. Strachey; Rawlinson; Saunders; Markham; *Calcutta Review*; Blanford—Nomenclature—Ethnical sketch: Plains of Assam and Bengal; Plains of Upper India; Clothing; Food; Customs; Eastern Himalaya—Eastern tribes; Tribes of the Southern Watershed; Bhután; Sikkim; Nepál; Tribes of the Lower Himalaya—Karnáli to Tons; Bishahr; the Satlaj to the Indus—Afghánistán, Galcha States; Kára-Tangutans; Tibet—General distribution—Appendix: list of authorities for this Chapter.

UNDER the name Himálayan districts of the North-Western Provinces of India we include the British districts of Kumaon, Garhwál, Taráí, Dehra Dún and Jaunsár-Báwar, and the independent State of Tihri or foreign Garhwál, comprising the tract within the Himalaya bounded by the

¹ The collection of materials for the 'Memoir on the Kumaon Himálaya' has been in progress since 1872, but other and more pressing duties have hitherto prevented their arrangement for publication. It was then intended to give as complete a description of the entire Himálaya as the means at our disposal would permit. There is little hope, however, that the leisure necessary for such an undertaking will occur within any reasonable time, and it therefore seems better to work up the materials already existing in their present form than to wait until opportunity is found for completing the original design. There are many and great gaps in our information regarding Kumaon, but the first step towards remedying this defect will be this attempt to take stock of our present knowledge, which is far more complete than is supposed. Without the aid of the materials entrusted to me by General R. Strachey, Sir John Strachey, Mr. J. H. Batten, Rudradatta Pant, and others, whose assistance will be found acknowledged in the preface, as well as the co-operation of Mr. H. B. Medlicott, Mr. S. A. Hill, Mr. Duthie, Dr. King, Dr. Watson, Captain G. Marshall, and others, my own work would be very meagre and unsatisfactory, as well from the great range of subjects discussed as from their frequent special and technical character.

Tons on the west, and the Káli or Sárda on the east. The adjoining portion of Tibet, to which British subjects resort for the purposes of trade, or in order to visit the sacred lakes, also comes within the scope of our investigations. A glance at the accompanying map will show that both the Himálayan and Tibetan portions of the tract with which we are concerned form but a small slice or segment of a greater system, which must be studied as a whole before an adequate conception of the structure and relation of its parts can be arrived at. The Himálaya itself is but the southern belt of that great girdle of mountains which encloses within them the country of which the southern half is commonly called Eastern Turkistán. From or through the southern slope of the Himálaya flow the great rivers known as the Indus, Ganges, and Brahmaputra. To the east, the continuation of the Himálaya is traced in the mountain ranges through which flow the Yang-tse-kiang and the Hoang-ho, and which are prolonged to the north in the Ala-shán, Inshán, and Khing-han mountains. The last of these ranges joins the Yablonnoi Khrebet branch of the Altai system at right angles in about 55° north latitude. The Yablonnoi mountains are the north-eastern continuation of the Altai range and form the water-parting between the Lena and the Amour. The Altai rises, on the right bank of the Irtysh river, at the north-western angle of the central plateau, and separates the Upper Gobi from the Siberian steppes. It consists of a belt of mountains varying in breadth from 400 to 1,000 miles, though in one place contracting to 150 miles, of no great elevation, and descending in a succession of broad terraces to the Siberian plains. It is pierced by the rivers draining into Lake Baikal, and, east of the 88th meridian, consists of three almost parallel ranges—the Saianak, Tangnou, and Ulangomula. From the sources of the Kára-Irtysh the Barluk Orochuk and Zungarian Ala-tau connect with the Tian-shán, the Celestial mountains of the maps, in which are the sources of the Syr-Darya or Jaxartes. The Tian-shán to the north and north-east of Káshgar consists of a series of parallel ranges having a direction from the east to west and an increased elevation as they proceed northwards. They abut on the elevated mountainous region known as the Alai and Pámír plateau, the latter of which contains the source of the Oxus in the little *lul* or lake of the lesser Pámír. The eastern margin of the Pámír is

described by M. Severtsof, who visited it in 1878, as neither a mountain chain nor the precipitous extremity of a table-land, but an extensive mountainous region filled with numerous ranges. The peak measured by Hayward (21,000 feet), and the Tágharma or Muztágh-Ata peak measured by Trotter (25,350 feet), and supposed to form a part of the Kizil-Art range connecting the Tian-shán with the Hímalayan system, are now reported to be only the culminating points of groups of snowy peaks separated from each other by a distance of over thirty miles. The intervening country contains the basin of the little Kára-kul lake surrounded by a complicated system of short and comparatively low mountain ranges having an altitude of from 14,000 to 15,000 feet. These ranges, however, connect with the Hindu Kush, Kárákorum, and Western Hímalaya, and thus complete the girdle of mountains from one or another side of which flow all the great rivers of Asia.

The country thus defined forms a part of the great Empire of China. To the south lies the rugged, elevated plateau of Tibet, regarding which we shall have more to say hereafter. To the north of this comes the depression known as the Lob-nor basin, which receives the drainage of the northern slope of the Tibetan plateau called, in the 39th degree of north latitude, the Altyn-Tágh range, but more generally known as the Kuen-lun, a name given probably from some well-known pass. The drainage from the eastern slope of the Tian-shán flows in the same direction, as well as that from the south-eastern slopes of the Altai range; but all is swallowed up in the great sandy desert of Gobi, which at one time apparently formed the bed of an ancient sea some 300,000 square miles in extent. According to Prejevalsky this plateau varies in height from 6,000 feet on the margin to about 2,000 feet in the middle. It is intersected from west to east by a depressed valley called Shamo, or 'sea of sand,' containing salt. West of this lies the Han-hai, or 'dry sea.' Prejevalsky has crossed the desert between Kuldja and what may turn out to be Lob-nor, lying in east latitude 90° and north latitude 39° 30,' and found himself at only 2,500 feet above the level of the sea on the banks of the Tarim. To the west, the desert presented a thin loam impregnated with salt, and to the east a plain of drift-sand. Towards the Kurugh-tágh hills, from which he descended on the

Eastern Turkistán.

desert, lay a belt of pebble and gravel some 15 to 18 miles wide. He also crossed the desert to the north between the Ala-shán range and Urga, where he found it to vary in height between 3,000 and 5,500 feet, whilst it still preserved its sandy character. On the route between Urga and Kalgan explored by the same traveller, there is a great depression towards the middle, where the elevation is as low as 2,400 feet. Here the soil of the Gobi proper is composed of coarse reddish gravel and small pebbles interspersed with drifts of yellow shining sand. Leaving these regions, we shall in future restrict ourselves to the southern plateau, of which the Kuen-lun mountains are the northern boundary, and which is so intimately connected in its physical relations with the Himálaya.

Before proceeding with our examination of the structure of the Himálaya-Tibetan region, it will be convenient to pass in review the different theories¹ that have been advanced in regard to its systematic geography, since a complete understanding as to what has been done in this direction will enable us to arrive at some conclusion as to what remains to be done. Captain Herbert, who conducted the mineralogical survey of the Himálayan country between the Káli and the Satlaj in 1818, was the first who attempted to give a general account of its physical characteristics. His description² was intended to serve as an introduction to his geological account of the Himálaya, as well as to be a distinct contribution to general geography; the existing works on the subject "being singularly deficient in details, as well as erroneous in the few that are given." His idea of the country north of India was apparently derived only from maps. He describes it as a large central space strongly marked by the feature that it was little intersected by rivers, whilst from its sides flowed the streams which united to form the greatest rivers in the world. As the source of every river must be higher than any other part of its course, he inferred that the zone in which those rivers originated must be higher than the plains through which they flowed to seek the ocean, and that the entire central tract itself was completely surrounded by lofty mountains. He considered the upper beds of the Brahmaputra

¹ For a brief summary of these theories see Mr. C. R. Markham's 'Memoir on the Indian Surveys,' p. 341.

² *J. As. Soc. Ben.*, XI., Pt. I., p. x.

and the Satlaj as forming part of the barrier zone which surrounds the central tract, and not as a part of the plateau itself. He further showed that the true "line of boundary is undoubtedly the chain of water-heads, and that this is by no means synonymous with the line of greatest elevation." At first view the arrangement of the mountain masses in the tract between the Káli and the Satlaj appeared to be irregular and confused, but by tracing the courses of the rivers and their tributary streams, a clue was found to lead the observer out of this labyrinth. "By connecting their sources and by following out the devious windings of the several feeders, an idea is obtained of the extent, the direction, and the connection of the several ranges. . . . Instead of a succession of parallel and continuous ranges running south-east to north-west, and rising one behind the other in regular array and increasing elevation till the series is closed in the farthest distance by the line of snow-clad peaks, we see only one continuous range of any extent forming an irregularly curved line which bends round the tract, commencing on the north-east angle, and with a north-westerly direction, which it gradually alters to a south-easterly one on the south-west angle, and latterly due south just before it is lost in the plain country. This range forms one of the boundaries of the basin of the Satlaj which bends around the convex side, while within its concavity are contained the numerous sources of the Ganges." This he called the Indo-Gangetic chain, "a ramification of that more extensive line of water-heads which would exclude from the central plateau all the mountain tract watered by the Sanpu and the Indus as well as by the Ganges. Next in extent are the two principal ramifications separating the basin of the Jumna from that of the Ganges, and the basin of the latter from that of the Káli. From these two principal ramifications proceed a number of minor ones which, but for the assistance derived from a study of the course of the rivers, would almost bid defiance to any analysis. Transverse ridges, several thousand feet higher in elevation, ramify from the Indo-Gangetic chain towards the Ganges basin, and a line or plane connecting their summits would be that of the greatest elevation, which, however, has no connection with the disposition of the water-heads. It is a fact that in a line of 500 miles two summits are found exceeding five miles in perpendicular height, not isolated, but connected to appearance by a

regular series of peaks of very little inferior elevation. If we confined ourselves to heights of 21,000 feet, we should find a connected line of peaks extending over 1,000 miles; that is, one apparently without breaks, but in reality connected only through the line of water-heads from which they ramify. Whether the word 'line' or 'plane' is used, the idea of considerable breadth must be conceded, and in that case its surface would be very irregularly studded with peaks, and in this way it may be said to be parallel to the common boundary of mountain and plain land, and to intersect instead of bounding the river districts." The above summary gives a *resumé* of Herbert's speculations on the physical structure of the Himálaya. His errors were those of his time, when the knowledge even of descriptive geography was in its infancy. He was unable to recognise the unity of the great central mass and its bulwarks, and was wrong in saying that the groups of snowy peaks intersected the river basins, when, in fact, they bound the drainage area, and are the determining causes of its existence. Still Herbert is to be remembered as the first who attempted to give a systematic account of the Himálaya as a whole,¹ and is therefore worthy of a prominent place in this brief notice of its geography.

Next to Herbert comes Hodgson, who in an admirable article in

the Asiatic Society's Journal² also alludes to
Hodgson.

the difficulty experienced by a traveller in

the Himálaya in getting "rid of that tyranny of the senses which so strongly impresses almost all beholders of this stupendous scenery with the conviction that it is a mighty maze without a plan." His first step towards freedom was his grasping the fact "that the vast volume of the Himálayan waters flows more or less at right angles to the general direction of the Himálaya, but so that the numberless streams of the mountains are directed into a few grand rivers of the plains either at or near the confines of the two regions." Secondly, a study of the river systems like the "Sapt Gandaki" and the "Sapt Kausiki" urged him "to discover, if possible, what cause operated this marked convergence of innumerable transverse parallel streams, so as to bring them into a series of distinct main rivers." Thirdly, he found that "the transcendent

¹ Neither Moorcroft, Vigne, Jacquemont, Hooker, Shaw, Henderson, nor Bellew attempt any description of the Himálaya as a whole. ² *J. As. Soc. Ben.*, XVIII., Pt. II., p. 761.

elevation and forward position, at right angles to the line of gháts of the great snowy peaks, presented that casual agency : the remotest radiating points of the feeders of each great river being coincident with the successive loftiest masses belonging to the entire extent of the Himálaya." The great peaks bound and do not intersect the principal Alpine river basins, as Herbert had thought, and, by so bounding, create the basins, whereas their intersection would destroy them. Hodgson's Himálaya proper is the ghát line or watershed between Tibet and India, and the watershed between the valleys of the Indus and Sanpu and the great plateau is called by him the Nyenchhen Thangla chain. The cause of the convergence of the various streams which form the great rivers upon or near the verge of the plains is shown by him to be "the superior elevation of the lateral barriers of these river basins, between which there are synclinal slopes of such decided preponderance that they overrule the effect of all other inequalities of surface, how vast soever the latter may sometimes be." These lateral barriers are crowned by the great peaks which stand forth from the watershed and send forth southward ridges proportionally immense. Equally effective with the divergent power of these peaked ridges is the convergent power of two ridges upon the single contained river basin. "The synclinal lines from the inner faces of the two adjacent ridges draw the waters together, and because these ridged peaks are the loftiest masses of the entire mountains, the effect of all other masses, even that of the spine of Himáchal or the ghát line of the snows, is overruled or modified, so that in the most rugged region on earth a very limited series of distinct main rivers appear in the plains from innumerable independent Alpine feeders." We may assume that where the loftiest peaks occur, there is a proportionate intumescence of the general mass, and therefore that these grand peak-crowned ridges determine the essential character of the aqueous distribution along the entire line. A further proof is adduced from the fact that the lower rivers, which take their rise in the middle region, do not show this unitizing principle, such as the Bágmati and Rámghanga. With regard to the mountain systems, Hodgson divides them into the lower, central, and upper ; sub-dividing the first into the sandstone range with its contained Dúns or Máris, the Bhábar or *adl* forest, and the Taráí. The lower region extends from the level of the

plains to 4,000 feet above the level of the sea; the central region from 4,000 to 10,000 feet; and the upper region to the watershed or ghát line: divisions which fairly correspond with the distribution of both organic life and inorganic matter. Though unable to follow Mr. Hodgson in all his theories and the deductions that he draws from them, credit must be given for his recognition of the position of the great mountain masses in regard to the alpine river basins and for his appreciation of the influence of climatic conditions on the animal and vegetable world.

Captain (now General) R. Strachey, in his paper on the Physical Geography of the Provinces of Kumaon and Garhwál, read before the Royal Geographical Society¹ in 1851, pointed out distinctly for the first time that the Himálaya was in truth the broad mountainous slope of the great Tibetan table-land descending to the plains of Northern India, while a slope of corresponding character descending to the north is known as the Kuen-lun. He remarks that the great peaks in Kumaon and Garhwál "are not found on a continuous ridge, but are grouped together in masses that are separated one from the other by deep depressions, through which flow the streams that drain those parts of the mountains that are immediately contiguous to the north." To the east the same sort of arrangement obtains, but to the west it is much less distinct. The river-beds to within a distance of ten miles in a direct line from the snowy peaks seldom exhibit a rise of more than four or five thousand feet; but when we cross "the line on which the great peaks are situated, the ascent very rapidly increases, and a very few miles carries the river-bed up to an altitude of nine or ten thousand feet; thus showing that the sudden increase of height of the mountains along this line is not confined to the peaks alone, but is a general elevation of the whole surface." Dr. Thomson²

Dr. T. Thomson. substitutes the name *cis-Satlaj Himálaya* for Herbert's Indo-Gangetic chain, and gives the name *trans-Satlaj Himálaya* to the chain which, commencing in Kailás, separates the waters of the Satlaj from those of the Indus. He refers to these two great chains the whole of the mountains between the Indus and the plains, and says: "The northern boundary

¹ *J. R. G. S.*, XXI, p. 57: adopted in Somerville's *Physical Geography*, 7th edition, London, 1877, p. 56.

² *Travels*, p. 456.

of Tibet is formed by the great chain north of the Indus, to which Humboldt gave the name Kouen-lun": and again, that every part of Tibet is traversed by mountains having their origin either in the trans-Satlaj Himálaya or the Kouen-lun. So far the unity of the

Himálaya-Tibetan region is acknowledged
Major A. Cunningham.

by this distinguished traveller. Major A. Cunningham¹ makes the Bara-lacha range, which forms the watershed between the Indus and its five affluents, the continuation of the main Himálaya or watershed between the Sanpu and the Ganges. To the south of this lies two distinct and independent ranges stretching in the same general direction from south-east to north-west, which he calls the mid-Himálaya, or Pir Panjál, and the outer or sub-Himálaya, leaving the name Siwálik unchanged for the lowermost sandstone ranges. Beyond the Himálaya the same system of parallel chains is observed, comprising at least three distinct ranges of mountains, which Cunningham proposes to call the trans-Himálayan, or that which divides the head waters of the Satlaj from those of the Indus and extends to the western limits of Rongdo and Astor; second, the Kailás or Gangri range which runs through the midst of Western Tibet along the right bank of the Indus to its confluence with the Shayok; and third, the trans-Tibetan range, also called

Captain H. Strachey.

Bolor and Kárákoram. These distinctions are however, purely local and geographical and are so far convenient and to be accepted. Captain H. Strachey, in his paper² on the Physical Geography of Western Tibet, shows us that the Indian watershed is not the Great Himálaya as seen by the Indian observer, but is found in a succession of valley heads much depressed and penetrating that mass to such a depth that the passes from India to Tibet are never visible from any station fairly south of the perpetual snow. The Turkish watershed divides the waters of Tibet from those of Turkistán, including Khoten and Káshgar. "The general plan of the mountain system of Western Tibet appears to consist of a series of parallel ranges running right across the breadth of the table-land in a direction so extremely oblique to the general extension of the whole as often to confound the one with the other, or to convert the transverse direction to a longitudinal one. Short transverse necks connecting the main ranges in some parts, and cross fissures cutting

¹ Ladák. p. 41.

² London. 1854.

through them in others, together with projecting spurs of a secondary order, will suffice to convert the supposed primary arrangement into all the existing variety of valley and drainage.**The great snowy peaks lying mostly on the terminal butt-ends of the primary ranges, sometimes widened by lateral spurs ; and the Tibetan passes crossing the low connecting links, whose alignment forms the main watershed, but not the main mountain-crest." Sir H. Rawlinson¹ recognises

Sir H. Rawlinson.

the unity of the entire mass, and writes that the "whole country between India and

Tartary may be considered as a broad mountain range, the Himálaya forming the southern crest, and the Kuen-lun the northern. The direction of this range is from east to west, trending to the northward, while the parallel chain which bounds Siberia to the south, and the outer crest of which is the Tian-shán, trends somewhat to the south ; so that at a short distance to the west of Yárkand and Káshgar the great interior depression of Chinese Tartary terminates, and the bounding ranges coalesce in the elevated table-land of Pámír."

We have now come to the theory set forth by Mr. Trelawny

Mr. Saunders' theory.

Saunders,² Geographer to the India Office, according to whom the summit of the Himá-

laya consists of a double range of peaks enclosing a series of valleys running parallel to the axis of the mass, and which he would call the northern and southern Himálaya respectively. The first of the two forms the water-parting between the Ganges basin and that of the Sanpu. To the latter must be assigned nearly all the great snowy peaks which are seen from the plains of India, and which are separated from the former by the valleys already mentioned. These valleys are comparatively elevated, and at length burst through the southern range by intersecting gorges. Both Herbert and Hodgson are set aside, and the great peaks are described as forming a chain, broken at intervals by intersecting gorges. "The upper valleys of the Sanpu, the Satlaj, and the Indus appear to form a huge elevated trough separating the Himálaya from the northern part of the table-land of Tibet and from the snowy range into which the table-land contracts at its western end." This range is crossed by the Muztágh, Kárákoram, and

¹ England and Russia in the East : London, 1875, p. 236.
Mountains and River Basins of India : London, 1870.

² Sketch of the

and Changchoumo passes, and is remarkable for the great length of its glaciers and the great height of its peaks. The Indus forms its southern base as well as the northern base of the Himálaya. The Indus, Satlaj, and Sanpu, "are the only rivers which, washing the northern base of the Himálaya in channels parallel to the range, break through the entire breadth of the range and water the plains at its southern base." The eastern base of the mountainous highland of Tibet is marked by the Min river, and on the north-east the slope is defined by the basin of the Hoang-ho. From the latter river westward to the Muztágh the Kuen-lun mountains descend to the plains of Gobi from the northern edge. These unite with the Himálaya, Pámír, and Hindu Kush in the lofty peak or knot called Pasht-khar or Tághdambash. The accompanying map, prepared by Mr. Saunders for Mr. C. R. Markham's Memoir on the Indian Surveys, will illustrate better than any further quotations his views on the subject of the relations of the great mountain systems, as well as serve our own purpose. Mr. Markham¹ divides the Himálayan system into

three great culminating chains, which he calls
 Mr. Markham. the inner, central, and outer, running more

or less parallel to each other from the gorge of the Indus to that of the Dihong. "The lofty region of Great Tibet lies mainly between the inner and outer range, with the central chain, whence most of the rivers of Northern India take their rise, running through its length." The western extremity of his inner and most northern range is the Kárákoram, which separates the Indus valley from the affluents of the Lob-nor system, and the eastern section is the Gangri mountains of the map, the Nyenchhen Thangla of Hodgson and Ninjinthangla or Nyenchhen-tang-la of Markham, which commences in peak or knot called Kailás. Parallel to the northern range runs the central range, the eastern section of which commences at the Mariam-la pass near the Kailás peak. "Here a comparatively low saddle connects the northern and central ranges and separates the valley of the Satlaj from that of the Brahmaputra. To the eastward the northern side of the central chain forms the southern watershed of the Brahmaputra, whilst on its southern slopes are the sources of many important rivers, which, forcing their way through the southern chain of the Himálaya, eventually join the Ganges or

¹ Tibet, p. xxiii., 1876.

the Brahmaputra." The southern chain is made up of the series of snowy peaks which, to the east, overhang Nepál, Sikkim, and Bhután. Thus, Mr. Markham is at one with Mr. Saunders in his theory as to the Himálayan system, only substituting the terms "inner or northern, central and outer or southern," for the terms "Gangri, northern and southern Himálaya," used by Mr. Saunders.

A writer in the *Calcutta Review*¹ has taken objection to the creation of the southern chain, which, "being occasionally intersected by rivers of more remote origin, is not a chain at all, but a series of spurs running southwards from an extended line of elevation more to the north, in the neighbourhood of which the said rivers rise." He also suggests for the whole system the name Indo-Tibetan, correctly urging that it is undesirable to give to the whole a name which belongs only to a part. He prefers simply to lay down two lines of watersheds, the northern corresponding for the most part with Mr. Markham's inner range, and the southern extending from Chilás by the Zoji-la, Baralacha, Níti, and No passes to the Laghalangla above Shikatsé. He then examines the river basins and shows that Hodgson's theory regarding them is in accordance with facts; that these basins derive much of their water from certain prominent peaks which, standing in advance—that is, southwards of the watershed—are connected with it, and from which ridges with dependent spurs project, that serve as lateral barriers to the basins. "The preponderating synclinal slopes of the ridges and spurs which overrule the effect of all other intervening inequalities of surface, however vast, cause the several groups of mountain streams between them to converge till they unite and constitute a main river near the edge of the plains." This is practically Hodgson's law re-affirmed in the full light of all that modern research has shown us regarding the geography of Tibet, the Karakorum and Káshgar, a *terra incognita* to our early writers.

Both Mr. Markham and Mr. Saunders have issued rejoinders to the criticisms in the *Calcutta Review* in two articles in the *Geographical Magazine*.²

¹ Rejoinders of Mr. Markham and Mr. Saunders.

² January, 1877, p. 145. ³ By Mr. C. R. Markham in May, 1877, and Mr. Saunders in July, 1877, *Geo. Mag.* IV, 113, 173, London. The other matters in controversy between the reviewer and Mr. Markham are omitted as foreign to the subject of this notice.

Mr. Markham chiefly confines himself to a defence of his use of the word 'chain' as applied to a series of culminating ridges, whether rivers force their way through its gorges or not; but Mr. Saunders goes more fully into the entire question at issue between him and the reviewer, and supports his arguments by a re-statement of his views on the physical geography of the entire Himālaya-Tibetan system. For this purpose he draws largely on his "Sketch of the Mountains and River Basins of India," already noticed, in which the theory of the southern chain of snowy peaks was first developed; and whether we agree with his deductions or not, we must consider his summary as a valuable contribution to our knowledge of the subject. He recapitulates the arguments in favour of considering the line of snowy peaks a southern chain, and concludes that they are entitled to that name, "(1) as the culminating summit of the southern or Indian slope; (2) as the common origin of a succession of rivers; (3) as cut off from the northern range by a succession of remarkable valleys, sometimes very long, sometimes very deep, and sometimes very broad and flat, and all containing considerable rivers running parallel to the chains which they divide." He objects to the inclusion of the mountain ranges on both sides of the troughs of the Indus and Sanpu under the term Himālaya, the northern watershed of those rivers composing the contreforts, buttresses and slopes or escarpments of the great central plateau which they uphold and from which they cannot be separated. The table-land is Tibetan; therefore its southern slope cannot be called Himālayan. The remaining portion of Mr. Saunders' article will be noticed as we proceed.

The latest contribution to the physical geography of the Himālaya is to be found in Mr. H. Blanford's Manual¹ and Mr. W. Blanford's introduction² to the "Manual of the Geology of India." In the latter work, which may presumably be taken as giving Mr. W. Blanford's conclusions on the subject, he considers the Himālaya to form a curved belt of mountains with their convexity to the southward which mark the southern scarp of the Tibetan plateau as the

¹ Physical Geography for the use of Indian Schools, Calcutta. 1879, L, ix. It should be remembered that the term 'range', is used here for geological purposes which are not always the same as those intended by geographers. Its precise meaning depends on the context.

² Calcutta,

Kuen-lun define the northern. The western terminal portion of the Himálayan chain comprises a number of great ranges variously named. It is doubtful whether any of these "should be considered the prolongation of the main Himálayan axis, although, if any be really a continuation of the Himálaya proper, it is either the Pir Panjal or the Zaskar range." Geological considerations would lead him to suppose that "the main range commences on the westward in the Dhauradhar near Dalhousie, and extends to the east-south-east till it rises into the main snowy range of the north-west Himálaya. Many geographers distinguish two parallel ranges from the neighbourhood of Simla to the eastward; the snowy range proper, formed of the highest peaks (Saunders' theory), and a more northern ridge, forming the watershed between the Tibetan plain and the rivers running to the plains of India. Others consider the latter to be the true Himálayan range, and look on the higher peaks as belonging to the spurs between the rivers flowing from that range. It is certain that the great peaks, such as Nandadevi, &c., are separated from each other by deep valleys, through which flow streams coming from the northern range, and that, although the peaks of the latter are inferior in elevation, the passes by which it is traversed are much higher; but it has not yet been ascertained whether the great peaks are on the strike of any continuous band of rock, or whether they merely consist of hard nuclei left undenuded." There is little doubt that, until the geologist is able to assist us, the question whether the line of snowy peaks should be considered a true chain or merely spurs from the main water-parting must be left undecided. Though year by year fresh materials are added to our stock of knowledge regarding the Himálaya, they are yet too imperfect for us to offer little more than a suggestion as to the views that should be adopted regarding its structure. A glance at Mr. Saunders' map will show us the vastness of the subject, and that the Himálaya of Kumaon and Garhwál, with which we are more immediately concerned, is but a very small portion of the great girdle of snowy peaks that uphold between them the elevated plateau of Tibet. Herbert showed us that this girdle, as seen from the plains of India, is not a continuous line of parallel ranges rising one behind the other, and increasing in elevation until the series is closed in the farthest distance by the line of

snow-clad peaks ; but that these peaks or groups of peaks are ramifications from the line of water-parting which itself is lower than the line of greatest elevation. Hodgson subsequently explained the influence of these groups of peaks on the river-systems, and Captain Strachey showed us that the Himálaya was the southern slope of the Tibetan plateau as the Kuen-lun formed its northern slope. These are, broadly, the more important additions to our knowledge of the physical geography of the Himálaya that have been made of late years. To our mind the recognition of the unity of the entire Himálaya-Tibetan system is the most important of them all, and that alone which will lead to practical results. The division of the Himálaya into ranges may be allowed as a matter of convenience, but should not be permitted to cloud the great fact that all are but variations in the southern slope of the great table-land due to the influence of the elements on the materials of which they are composed, and to the disturbing action of subterranean forces. We can lay down the line of water-parting and the line of greatest elevation with some precision, but must call in the aid of the geologist and mineralogist to distinguish which amongst the ranges is entitled to be called the real main axis of the Himálaya ; and, until their labours are communicated to the world, must rest content with the somewhat arbitrary distinctions afforded by the prominence or otherwise of existing physical features.

Seeing the misunderstandings that have arisen from a too loose use of words and phrases, it will be as well
 Nomenclature. to state here that we adopt the word 'water-parting' to represent the ridge which separates the flow of water on either side of a range of hills.¹ The word 'range' will include a series of mountains or hills continuing in one direction along a common axis, whether broken by chasms or not ; and the word 'spur' will be used of a ramification from a range, whether connecting it with another range or sinking gradually into a plain.

The great mountain chain lying between Tibet and the plains of India is generally known to the natives of India by the term *pahār* (mountain), to which they prefix the local name where such exists.

¹ The use of this word in this sense is one of the subjects of controversy between the Calcutta reviewer and Mr. Markham. The former (p. 147, note) objects that the old word 'water-shed' is sufficient.

The more educated give the name Himáchal¹ (snowy-range) or Himálaya (abode of snow) to the snow-covered ranges; whilst Europeans popularly include under the name Himálaya the entire mountainous region lying between the gorge of the Brahmaputra on the east and that of the Indus on the west, and between the upper valleys of the same rivers on the north and the plains of India on the south. A first glance at any good map will convince us of the general unity of the physical relations of the range within the limits commonly assigned to the Himálaya, whilst a closer examination will induce us to include much more. For our part we accept the popular definition of the Himálaya as extending from the gorge of the Indus on the west to that of the Brahmaputra on the east, and from the upper courses of the main branches of those rivers on the north to the plains of India on the south, speaking of its connections beyond those limits as the western and eastern extensions respectively.

It will materially aid the reader if we further preface our remarks with a short description of the ethnical and political divisions of its surface, and of the regions in immediate contact with it. We have arrived at some idea of the physical relations of the tract itself, and shall now, at the risk of being thought diffuse, endeavour to trace the ethnical affinities of its inhabitants. Commencing, then, with the plain on the south, we find the provinces of British India flanking the foot of the Himálaya along its entire length from the 96th to the 72nd meridian of east longitude. Following the direction of the Himálaya from east to west, we find in Upper Asám a number of tribes speaking different languages and dialects, and so intermixed and blending the one into

¹ The word Himáchal (हिमाचल) is derived from two Sanskrit words, 'Hima' (snow) and 'achala' (mountain), meaning 'snowy-mountain' or 'snowy-range'. Similarly the word Himálaya (हिमालय) is derived from 'Hima' and 'alaya' (abode), meaning the 'home' or 'abode of snow.' The proper pronunciation is therefore Him-a-lay-a, not Him-a-láy-a as commonly obtains. The plains-men speak of the Simla-paádr, the Mansuri-paádr, and sometimes of the snowy-range as the barf (ice)-paádr. "The people south of the Himálaya in Nepál call all snowy mountains *langer*, by which they mean the highest points. They call the peaks that have no snow *barjang*, and the low ground under the said *barjang* they call *phedi*. The term Himálaya is not used by uneducated people, who only talk of the snowy mountains as '*barfáni langer*.'—*G. T. S. Rep.*, 1872, p. 46.

the other that, beyond a mere cursory description, their classification cannot be attempted here. In the extreme north-east they are

Plains of Asám and allied with or are members of the tribes inhabiting the neighbouring hills, of whom more

will be said hereafter. They speak a language having an affinity with the great Barina-Tibetan group, and are more pagan savages. Along either side of the Brahmaputra in its course through the valley we find the settled tribes of Asám speaking a language akin to Bengáli. Though differences exist they so closely resemble in habits and character the people of the conterminous parts of Bengal that it is difficult to draw a strict line of severance between them without entering into long historical and ethnical discussions quite out of place here: many that are now Hindu or Musalmán Bengális in all outward appearance can be shown to be converts in recent times from the pagan tribes in their neighbourhood of unmistakably aboriginal origin. The people of Bengal, the flat alluvial plains of which lie along the lower courses of the Ganges and Brahmaputra, exhibit all the features characteristic of a race inhabiting a region of tropical heat and moisture. They are small in stature, of dark complexion, and effeminate in character, living chiefly on rice grown in the lowlands subject to annual inundation. Their dress is of the scantiest proportions, consisting chiefly of one or two pieces of cotton cloth simply wrapped around their waist and shoulders, and not wrought into any form of garment. Their heads and feet are usually left bare. Their houses, constructed of mats, lie scattered among the thick groves of bambus and palms that spring up in wild luxuriance on the uncultivated ground.

As we ascend the Ganges,¹ we find a drier climate with greater contrasts between the summer and winter temperature, and a taller, more manly, and more robust race, of whose food the millets and unleavened bread of wheat, barley, and other grains form the principal element. Their clothing is more elaborate and warmer than that of the Bengáli. All wear turbans, and those who can afford it have short jackets fastening on the right breast in the case of Hindus, and on the left breast in the case of Musalmáns. Their houses are built of mud and are either

¹ This account of the Hindus in the plains is partly based on Elphinstone and Notes by General R. Strachey.

tiled or thatched in the villages, but in many of the towns very small bricks are used in the construction of the better class of dwellings. The country is open and unenclosed, and almost the only trees are groves planted near towns and villages, with occasional patches of *dhák* (*Butea frondosa*) and *babul* (*Acacia arabica*) jungle. The people are mostly Hindu in religion and speak dialects of Hindi. On approaching the Satlaj the language passes into Panjābi, also Sanskrit in its character, and we find the religion of the Sikhs the seal of the double dispensation of Brahma and Muhammad.¹ To the west of the Panjāb, or country of the five rivers, the religion of Islām predominates amongst a motley group of tribes of very varied origin. The name Hindustān, which is more correctly applied to the northern Gangetic districts alone, may without impropriety be used so as to include the entire tract below the Himālaya. Intersected by the innumerable streams that flow from the mountains above it, watered by the copious falls of periodical rains, and enjoying a semi-tropical climate, the great unbroken plain is thus supplied with the two great requisites, heat and moisture, that are the necessary and certain agents for the development of vegetable life. We naturally, therefore, find an agricultural population often in the older settled parts extremely dense, and attaining to no small degree of civilisation. Cities and large towns are common, many with a population of over 50,000 souls; and the inhabitants, without coming up to a European standard, enjoy considerable wealth. Literature, both indigenous and of European origin, is cultivated; schools are numerous, and the useful arts are highly advanced and eagerly followed. The hot climate which induces a love of repose and fertility of soil which renders severe labour unnecessary has, in some measure, modified the habits of patient industry which are usually characteristic of an agricultural population; but the dislike to change which marks those communities in all parts of the civilised world is nowhere more strikingly exhibited than in Hindustān. The great wealth of the country and its open and easily accessible character, together with the insuperable obstacles to union presented by differences in race and caste, have, for many centuries, subjected it to the reiterated attacks of foreigners. With very few exceptions all truly national government has ceased to exist; and from what little

¹ Cunningham, History of the Sikhs, p. 18.

we know regarding it, the people have small cause to regret the successive changes of masters. India has never had in the whole course of its history so strong, universal, and just a government as it has enjoyed under the British since the memorable mutiny of 1857. Englishmen may well point with just pride to the lengthy catalogue of measures attesting true moral and material progress that have been introduced during the last quarter of a century, and have been assimilated by the people to such a degree that the advance—political, moral, and social—made has eclipsed all that had been previously effected under British rule.

Taking the people of the plains as a whole, their clothing is cotton and their food is vegetable, though
 Clothing, food, customs. Musalmáns and some Hindús eat meat and a few eat fish. The dress of the men, as a rule, is white, though they often wear coloured jackets; the women, who have no other covering for their heads than a corner of the cloth that they wrap round their bodies, frequently wear bright colours, usually indigo blue, Turkey red, or safflower yellow. The Hindús preserve their moustache, but shave their beards, and frequently their heads, except a small scalp-lock, whilst the Musalmáns allow their beards to grow. The seclusion of women seems to be a custom introduced by the Muhammadans, but amongst all religions and races in Hindustán the position of females is essentially inferior to that of the other sex, with whom they do not mix in society. Marriages are almost always contracted in childhood, and the betrothed bride is always under the age of puberty. Amongst Hindús, a dowry is given with the daughter, though the practice of accepting a sum of money for a daughter is in many parts of the country gaining ground. Female infanticide has been rife in the Gangotic districts of Upper India, due to the disgrace supposed to be attached to the expression 'father-in-law,' as well as to the great expenses ordinarily consequent on the marriage of a daughter. The education of women is absolutely neglected, and the efforts of Government in this direction have proved a total failure. The men, however, for the most part, amongst the classes above those actually engaged in the cultivation of the soil, can read and write, and even the men who have not acquired those attainments possess the power of mental calculation to a remarkable degree. The Brahmanical faith is with few exceptions

dominant throughout Hindustán. The Musalmáns are, however, numerous everywhere, and in some districts are in the majority. They belong chiefly to the Suni sect, but Shiahhs are also found in certain tracts where the influence of powerful families of their persuasion has been felt. The peculiar modification of Hinduism adopted by the Sikhs is chiefly confined to that portion of the Panjáb which lies east of the Chináb. To the west of that river, the great mass of the population is Muhammadan. Taking the people of the Panjáb proper, excluding Peshawar and the trans-Indus districts attached to the Leia and Multán divisions, we have a population of ten millions, of whom seven-twelfths are Musalmáns, four-twelfths are Hindús, and one-twelfth are Sikhs. Distributing them according to race, General Cunningham¹ makes 3 per cent. of so-called early Turanian origin, 27 per cent. Aryans, and 70 per cent. later Turanians.

We shall now consider the ethnical and political divisions of the Himálaya itself, proceeding in the same direction from east to west. At the extreme Eastern Himálaya. east we have the same races speaking a Barma-Tibetan language that we found in the plains, but a line drawn north and south across the Brahmaputra, in the general direction of the Dhansiri river, and continued southwards so as to leave Kachár to the west of it, would, according to Hodgson, divide them from the Alpine races of more pronounced Tibetan stock, as well as from the so-called aboriginal tribes of the central Himálaya. These Barma-Tibetan tribes are known as Abors, Bor-Abors, Daphlas, Akas, Mishmis, Miris, &c., and their communities are reported to have a sort of rough republican constitution. This conjecture of Hodgson appears to be supported by the result of the most recent investigations.

The country lying on the Táwáng route by the Dhansiri river from Asám to Chetang in the valley of the Eastern tribes. Sanpu, in the 92nd meridian of east longitude, has been traversed by one of the Pandits of the Great Trigonometrical Survey, from whom we learn that to the east of that line the Himálaya is inhabited by Lhoba Daphla tribes.² These men are remarkable for the abnormal development of the muscles of the arms and calves of the legs. They wear cylindrical-shaped hats made of

¹ Cunningham's Arch. Rep., II., 2, 4.
attached to this chapter.

² See section A. of references

bambus, and their only garment is a long blanket folded somewhat after the fashion of a plaid and fastened round the waist by a cloth girdle, which is used as a quiver for their arrows, which all carry, as well as a bow slung over their left shoulder. The greater part of their legs and arms is bare. They wear no boots, but ornamental rings made of rope, fastened very tightly both on the wrists and legs below the knee. They have a decided Tibetan caste of feature, high cheek-bones, and Chinese-looking eyes. They wear no hair on the face, but the hair of the head is allowed to grow to a great length, and is drawn together behind the head and then allowed to hang down.¹ They appear to be distributable into two groups—those living in the great rice country to the north on the banks of the Sanpu, called Lho-khals, and who are independent of the Lhasa authorities; and the Shiyar Lhobas, a wild race who inhabit the country through which the great river flows to Gaya, Asám, and who may be identified with the wilder tribes of Mishmis. The Mishmis are distributed into three great divisions comprising numerous clans—the Chulikata or crop haired, the Midhu, and the Digaru, each of whom have a separate dialect, and the last reside within British territory along the hills as far west as the Digaru river. The Abors or Padams inhabit the country to the west of the Dibong river. They are described by their neighbours as exceedingly fierce and blood-thirsty: “like tigers, two cannot dwell in the same den. Their houses are scattered singly or in groups of two and three over the immense extent of mountains inhabited by them.” They manufacture the weapon called *dao*, and weave coarse cloth, which with *manjit*, beads, bell-metal cooking vessels, female slaves or rather wives, and the breed of cattle called *mithans*, are exchanged for salt and coarse cloth imported by the Miris from the plains. The Miris are more civilised than the Abors, and dwell in villages both in the hills and plains. The Akas or Hrussos live between the Miris and the Daphlas on the upper waters of the Sundari, and call themselves Tenaes. The women of the Akas wear blue or black petticoats and jackets of white cotton of their own manufacture. Their faces are tattooed, whence the name “Aka” given them by the people of Asám. The males wear a girdle of canework painted red, which hangs down behind in a long bushy tail. Their staple food is rice, but everything edible is made use of. The

¹ G. T. S., 1872, p. 70.

Daphlas belong to the same stock, and all are mere pagan savages, debased, cruel, and treacherous, though in the last respect the Akas have a somewhat favourable report.

To complete our review, we shall diverge to the south of the
 Tribes of the southern Brahmaputra and briefly notice the tribes
 water-shed. inhabiting its southern water-shed.¹ In the
 extreme east we find the Khamtis or Shans, a tribe linguistically
 allied to the Siamese and Buddhists in religion ; next we have the
 Singphos, or Kakhyens, and the Jilis, on the northern slopes of the
 Patkoi range, both of whom are pagans and speak a language
 intermediate between Barmese and Tibetan. Further west come
 the Nágas, who are distributed into three great classes—the Namsang,
 Khari, and Angámi. They are the most numerous of all the pagan
 tribes to the south of the Brahmaputra, extending from the Kopili
 river in the meridian of Nowgong on the west to the meridian of
 Sadiya on the east. They bury their dead and appear to manage
 their affairs in a sort of republican assembly. The Kopili river
 separates the Nágas from the Khasiyas of the Jaintiya and Khasiya
 hills around Shillong. The Khasiyas appear to be an isolated group,
 speaking a monosyllabic language which cannot be classed with any
 other of the same family. The form of government is republican
 and the religion is mere paganism. To the west are the Gáros, who
 also are pagans, though their language has affinities with the Aryan
 dialects spoken on the north, south, and west. The language,
 however, has a Tibetan basis, and Hodgson would include it in the
 Bodo group, of which more hereafter. South-west of the Nágas come
 the tribes of the Manipur, Lushái, Tipura (Tipperah), and Chitta-
 gong hills. From McCulloch and Damant we learn that there are
 numerous dialects in Manipur, and that the principal has a character
 of its own derived from the Nágari. The inhabitants have adopted
 the Brahmanical faith. The people further west are known as Kúkis,
 and appear to speak four dialects of a common stem-language :—
 the Lushái spoken by the Dzoes of the Lushái highland, the Thadu
 in northern Kachár, the Kúki in the same district, and the Hallámi
 in the Tipura hills. The Kúkis are pagans, but are gradually yield-
 ing to the influence of their Brahmanical neighbours, as indeed are
 all the pagan tribes similarly situated in the Asám valley. We shall

¹ See section B. of references attached to this chapter.

now return again to the tribes inhabiting the Himálaya to the west of the Dhansiri river.

The whole country along the Táwáng route from Asám to the head of the valley leading down to Chetang on the Sanpu is under the rule of the Lhásan Jongpen of the Chona Jang. The Lámas of the great lamasery of Táwáng, however, own the country to the south of the range of hills which form the water-parting between the Táwáng and Dhirang valleys, and are entirely independent of Lhása. They manage all public business in an assembly of the principal Lámas, called the *Kato*, which is also the supreme court of justice. To the north, near the Chetang valley, the elevated highlands are occupied by nomad tribes; but to the south, in the Mon-yul or Sub-Himálaya, the country within which Táwáng is situated, the people are called Monpas or Hill Indians, and differ materially in language, dress, and manners from the Bodpas, or people of Bod-yul to the north of Chona. The Monpas resemble the inhabitants of Bhután on the west. They wear their hair closely cut round the head, not in plaited tails as in Tibet, and as a covering have a small skull-cap of woollen cloth or felt. Instead of the long gown of Tibet a short coat is worn which reaches to the knee, and is fastened by a woollen girdle that invariably holds a long, straight knife. The people keep cattle, sheep, and pigs. Thus, the line drawn north and south by the Dhansiri river passes through this wedge of Tibetan territory, separating the Barma-Tibetan tribes on the east from those who have a more pronounced Tibetan origin in the central Himálaya and the so-called aboriginal tribes of the lower Himálaya on the west. Hodgson assigns to the latter the name Tamulian, but it cannot stand, involving as it does linguistic and ethnical associations which modern research has failed to establish.

To the west of the Dhansiri river we have the countries of Bhután, Sikkim, and Nepál, all of which possess a more or less established form of government.¹ Much has been written regarding the people inhabiting those countries which we can but very lightly touch upon here. The entire tract may be divided into three great belts,—the elevated region beyond the snowy range that is visible from the plains, varying

¹ See section C. of references attached to this chapter.

from 10,000 to 16,000 feet above the level of the sea ; the central region, varying from 4,000 to 10,000 feet ; and the lower region, extending from the plains to 4,000 feet.¹ To the central region are confined the Lhopás, Lepchás, Limbus, Kirántis, Murmis, Newars, Sunwárs, Chepángs, Gurungs, Magars, and Khasiyas. To the lower regions belong the Koch, Bodo, Dhimál, Kichak, Tháru, Denwar, and Pallah tribes. The inhabitants of the central region belong to a comparatively recent Tibetan immigration, whilst those of the lower region, the so-called Tamulian aborigines, are apparently to be ascribed partly to an early Tibetan immigration and partly to an Aryan source. To the north, along the entire line of gháts from the 92nd meridian to the Jumna, we find the Bhotiyas or Bod-pas of pure Tibetan origin and Buddhists in religion. Bhután, the Lho-pato, Lho-duk, or Lho-mon of the Tibetans, is also a Buddhist country, as well as Sikkim, the Demojong of the Tibetans. Nepál, called Palbo by the Tibetans, is partly Buddhist and partly Brahmanical in religion. In the central Himálaya of Bhután and Sikkim we find the Lhopás, Lepchás, and Limbus. The name 'Lhopa' seems to be a generic term signifying the people of Lho or Bhután, as 'Bod-pa' means a person of Bod or central Tibet, and 'Kham-pa,' a person of Kham or eastern Tibet. So also the term 'Dok-pa' is the religious equivalent of the territorial term 'Lho-pa.' The Lepchás extend from Panákha in mid-Bhután on the east into eastern Nepál on the west. They are divided into the Rong and Khamba tribes. The women of both divisions wear a loose coat of the fibre of the silk-worm that feeds on the castor plant, or of unbleached cotton with a wrapper of the same material around the waist to form a petticoat. The men wear a robe of striped red and white cotton cloth crossed over the breast and shoulders and descending to the calf of the leg, leaving the arms bare ; a loose jacket of red cotton cloth is worn over the robe by those who can afford it, and both are bound round the waist by a red girdle. Some strings of coloured beads round the neck, silver and coral earrings, a bambu bow with a quiver of iron-pointed arrows, and a long knife complete their costume. This knife, called '*bán*' by the Lepchás and '*chipsa*' by the Bhotiyas, is worn on the right side, suspended from the left shoulder, and serves as

¹ Better known as Hodgson's belts.

an axe, hoe, spade, sword, and knife. The Lepchás eat any flesh of bird or beast, all cultivated and many wild grains, and drink beer and tea. They are Buddhists in religion. Their language, though allied to Tibetan, is not Tibetan, and has a character of its own. They bury their dead; though the Murmis, a tribe of the same country, first burn their dead and then bury the ashes. The Lepchás are short in stature, averaging about five feet, bulky for their height, and rather fleshy than sinewy. They have a fair complexion, pleasantly marked Tibetan features, and part their hair along the crown of the head. Both sexes allow it to grow long; the younger males allow it to hang loose over the shoulders, whilst the elder males and women plait it into tails, and the latter tie the ends with braid and silken cords and tassels. Like all Buddhist hill-tribes they are very filthy in their habits, ablution being unknown.

The Limbus are an important tribe of the central Himálaya, found between the Tista on the east and the

Limbus.

Dúdh-Kosi on the west. The word 'Limbu,' according to Dr. Campbell, is a corruption of the term 'Ekthumba,' the correct name of this people, and used generally to designate the whole population of this portion of the Himálaya not included amongst the well-known divisions, such as Lepchás, Murmis, Bhotiyas, and Parbatiyas. Under the name Limbu are included the Kirántis or Kirátis, Ekas, and Ráis, and their country is divided into Kiráti-des from Dúdh-Kosi to the Arun and Limbua from the Arun to the Konki river, which leaves the Nepál hills about twenty miles to the west of the Mechi river. Further east and west they occur only in small colonies. Hodgson records the vocabulary of seventeen dialects of the Kiráti language, none of which are referable to the written Tibetan or Hindi. They are pagans in religion, though willing to pass themselves off as followers of Hinduism or Buddhists where those religions prevail. Their features, the absence of a beard, and the colour of their skin, all show them to be of Tibetan origin. The Limbu wears his hair long, but does not plait it; he carries a *kukhri* or curved knife instead of the *ban*, and has a wide trousers and jacket instead of the robe and long jacket of the Lepcha. Both tribes are found at elevations of from 2,000 to 4,000 feet above the level of the sea.

The Hayus or Vahus prefer the lowest elevations in the valleys, and occupy the central and lower ranges of the mountains of eastern Nepál between the Arun river and the Konki. They are found mixed with the Eka division of the Limbus, but possess clearances and villages of their own. They differ from all around in language, religion, and habits, and are esteemed an outcast race by the Gurkhális. They do not intermarry or hold intercourse with other tribes. The Bramhus, similarly placed in the Noakot valley to the west, speak a Barina-Tibetan dialect and are also pagans. The great bulk of the Murmis are found between the Nepál valley and the Dúdh-Kosi, whence in smaller numbers they extend to the Tista on the east and as far as twenty miles west of Kathmandu on the west. They are divided into two classes: one from the Ni district in Tibet, and the other from the Tsang district; hence the generic name Nitsang or Nishang applied to the whole tribe. They prefer elevations of from 4,000 to 6,000 feet and engage in pastoral and agricultural operations, living in cottages built of stone and thatched with grass. They are Buddhists in religion and their language is akin to Tibetan. In their physical traits they resemble the Lepchás, though somewhat taller. The Newars compose the majority of the inhabitants of the Nepál valley, but are not numerous beyond its limits. They are a shorter race than the Gurkhális, and their appearance betrays their transmontane origin. The greater number are Buddhists, and the remainder profess the Brahmanical faith. They possess a written character of their own and speak a distinct dialect, though the Buddhist portion also use Tibetan, in which their religious books are chiefly written. In the forests to the west of Nepál, close to the plains, we have the wild tribes known as Chépáangs and Kusundas, the former clearly akin to the Rájis of Kumaon. They speak a language allied to that of the Lhopás of Bhután. In the same direction are the Hinduised tribes of Khasiya, Magars and Gurungs, generically known as Parbatiyas. They speak a language having a Tibetan basis, and into which many Hindi and Urdu vocables have been introduced. The Gurungs, like the Murmis, prefer elevations of 6,000 feet, and are partly pagan and partly Hindu in religion. The Magars are entirely Hindu in religion, and to them

belong the Thápa clan so famous in later Nepálese history. Both these tribes supply numerous recruits to the regiments in the British service, and to this is probably due the Indianised form of their speech. North of them we find the Sunwárs, and on the west the Thaksyas (Thakuris?), and on the east the Pahría. Vocabularies of the languages of these tribes have been preserved by Hodgson.

The Gurkhális speak the Hindi dialect called Nepálese Khas or Parbatiya. In summer they wear a sort of pantaloons called *pájámas* and a jacket or coat of white or blue cotton, and in the winter the same padded with cotton or lined with fur and fastened by a cotton girdle, which invariably holds the heavy, crooked knife, called *kukhri*. Turbans of dark cloth or loosely-folded cotton are used as a head-dress, or small tinsel, embroidered skull-caps. The Newárs wear a waist-cloth of cotton and a jacket of the same, or some woollen materials. Some adopt the Tibetan costume of full short trousers, a long tunic, and a fur-edged cap. The head-dress is a small skull-cap of black or white cloth thinly wadded with cotton and generally turned up an inch or so at the border. The dress of the other inhabitants of Nepál differ little from that already described. The women of the Newárs wear their hair gathered into a short thick club at the crown of the head, whilst others have it plaited into a long tail. Flesh is much more commonly used by all classes than in the plains. The lower classes drink a coarse fermented stimulant called *rakshi*, and the higher classes, when they can afford it, consume large quantities of tea.

Taking now the tribes inhabiting the lower Himálaya to the Tribes of the lower north of the Brahmaputra, we find on the Himálaya, extreme east the Deoriya Chutiya, the remnants of a powerful tribe, who though Hinduised in religion preserve their old language, which is affined to the Barma-Tibetan group. Next come the Dhimál, Kachári or Bodo, and the Koch. Hodgson tells us that in travelling between Gwálpára in Asám and Aliganj in the Morang Taráí of Nepál one has to pass through the country of the following tribes:—the Koch, Bodo, Dhimál, Rábha, Hájong, Kúdi, Batar or Bor, Kebrat, Pallah, Gangai, Maráhi, and Dhanuk. The Rábha, Kúdi, Hájong, Mech, Gáro, and Páni-koch, are all affined to the Kachári or Bodo type. The last six of Hodgson's list are doubtful and undefined

and require further investigation. The Dhimáls are found in the *sál* forest between the Konki and the Tarsa, mixed with the Bodos, but without intermarriage and living in separate villages. The Bodos extend from the Surma to the Dhansiri, and thence by Bijni and the Bhután and Sikkim Taráí to the Konki; besides occupying outside the forest limits a large proportion of central and lower Asám.¹ The Rábhús and Hájongas are found in the Gwálpára district and are Hindús, whilst the Páni-koch occupy the tract along the foot of the Gáro hills and are still pagans. The Dhimáls and some of the Bodos, Kochs, and Mechis are still pagans. The last-mentioned tribe is found all along the Taráí with the Kochs and Dhimáls. Their dialect and religion differ from those of the neighbouring tribes of the hills and plains. They are fairer than the Kochs and have strongly marked Mongolian features, but softer than those of the Lepcha or Limbu, resembling more the Newárs than the other hill-tribes of Tibetan origin. They live at elevations between 800 and 1,000 feet, and almost always keep to the forest, where they make temporary clearances. Their religion is connected with the Bhairava form of Sivaism. The Koch tribe is now nearly completely converted to Islám or Brahmanism, and with their conversion have dropped their old name and language, speaking a corrupted form of Bengáli, in which, however, many of the ancient vocables are retained. The Páni-koch, according to Hodgson, represents the unimproved primitive Koch stock; but Dalton considers them a plains tribe driven upwards by the Aryan invader. Hodgson estimates the number of all religions at over a million souls. The settled Koch assimilate in their food and clothing with the Bengális, and show no marked differences. The Bodo women wear garments of coarse silk, the produce of the worm that feeds on the castor-plant. The Bodo men and Dhimáls of both sexes wear cotton clothes. The men wear one cloth thrown over the shoulders and another wrapped round the waist and drawn up between the legs. The female garment consists of a cloth wrapped around the body and enveloping it from the arm-pits to the centre of the calves. Wooden sandals are worn, but ornaments are rare, though the women sometimes wear small silver rings in their noses and ears and heavy bracelets of mixed metal. Meat, fish, and

¹ See Hodgson's *Aborigines of India*, p. 151.

vegetables are eaten by all, and beer made of rice or millet is a favourite beverage. Thárus and Denwars are found westwards in Nepál mixed with the Mechis, and especially the former in the malarious tract in the Gorakhpur and Tirhút Taráis along the foot of the mountains where no other human being can live. They, however, seem to be healthy, robust races.

The mass of the people of the Doti district of Nepál and the British district of Kumaon belong to the race
 Karnáli to the Tons. generically known as Khasas or Khasiyas.

In Garhwál they are more mixed, though the difference is scarcely discoverable. The northern inter-Alpine valleys are, however, inhabited by Bhotiyas, who are decidedly of Tibetan origin. Amongst the Khasiyas there is a great admixture of immigrants from the plains, and most of the better classes look down with contempt on the purer members of the Khasiya class, who appear to represent the oldest inhabitants of these hills, though now much modified by centuries of close connection and intermarriage with the more civilized tribes of the plains. Throughout Kumaon, the Kyunam of the Tibetans, the inhabitants dress and eat like those of the plains, the only difference being that to the north woollen materials find more favour, and there is greater license in matters of food and drink. In Garhwál, which is known as Galdiya to the Tibetans and to the north as Chongaa, garments made of hempen fibre are common amongst the poorer classes. The language spoken throughout is pure Hindi, though for purposes of trade the Bhotiyas also use Tibetan, and amongst themselves speak a dialect of Tibetan origin. In the land of marsh and forests which borders the plains we find the Thárus in the eastern Tarái and the Bhuksas, a tribe of similar character, occupying the tract between Purnapur-Sabna in the Bareilly district and Chandpur in the Bijnor district. The Rájis in eastern Kumaon are akin to the Chepángs of Nepál, and the Lúls and Rawats of the same tract are now absorbed in the Hinduised population.

To the west of the Tons we have a number of petty independent states known as the "protected hill-states,"
 Bisahr. followed by British territory. Amongst the former the most important is Bisahr,¹ of which the northern part,

¹ See section D. of references attached to this chapter.

called Kunawar, or more correctly Knaor, the Kunu of the Tibetans, marches with Tibet. The people of upper Knaor are of Tibetan origin and Buddhists in religion, and correspond to the Bhotiyas further east. Buddhism extends down the valley of the Satlaj as far as Sarahan, between which and Pangi is a sort of debateable ground common to Hindús and Buddhists; but north of Pangi Buddhism prevails, and south of Sarahan, Hinduism. With our approach to these Buddhist countries the curious custom of polyandry appears. Commencing in north-western Tihri, we trace it through Bisahr and Láhul, but find it confined to the inhabitants of the valleys of the central and higher ranges professing both the Buddhist and Hindu religions. The central tract is inhabited by a fair, slight, and muscular race of mixed origin known as Kunets, and akin to the Khasiyas on the east. To the south, in the portion adjoining the plains, the people resemble the inhabitants of the lower country, and appear of every shade of colour from dark-brown to a tawny yellow or yellowish-white. The hair is black and worn long at the sides and back of the head down about the ears, where it is cut short. The crown of the head is shaved bare, but moustache and beards are worn. The dress is a short coat of coarse cotton reaching to the knee, pleated in folds to give it fulness, and fastened round the waist by a girdle of the same material. A pair of cotton *páejámas* and a sheet of the same material complete the hot-weather costume. In winter these are exchanged for a pair of woollen drawers and a blanket, but the poorer classes remain content with a coarse waist-cloth and a blanket all the year round. The food of the people from the Káli to the Indus differs very little in each tract, or, indeed, from that of the people in the adjoining plains. Wheat, barley, rice, and various millets and pulses are grown in the lower hills, and to the north hardier varieties suited to a sub-arctic climate are cultivated. When the produce is insufficient for the wants of the inhabitants, a supply is imported from the lower districts. To the north, woollen home-spun replaces the cotton worn in the lower hills, and the girdle supports an axe. The women wear a similar dress, the coat reaching down to the ankles, and the hair, done up in long plaits, is twisted into rolls and covered with a piece of cloth wound like a turban round the head.

To the west of the Satlaj we find the British territory of Kulu and Mandi, the independent state of Chamba, and the territories of Kashmir and Janu.¹

Satlaj to the Indus.

Kulu, called in Tibetan Nyungti and Mandi, lies along the upper course of the Byás; Chamba, known to Tibetans as Punga, along the Rávi; and Láhul, the Tibetan Garzha, on the Upper Chináb. To the north in Láhul and Ladák the people are Bhotiyas or Bodpas of Tibetan origin, professing the Buddhist religion, and amongst them is a servile race known as Bem or 'low.' Further east in *Baltistan* are Tibetan Musalmáns who have adopted with their religion the Arabic alphabet. All these have decided Mongolian features, and are noted for their strength of body and power of enduring fatigue. In this respect the Baltis are somewhat inferior to their Buddhist brethren in race. The men wear a coat of woollen material reaching to the knees, fastened by a girdle, in which a knife is usually carried. Round their legs, from knee to ankle, they have coarse woollen leggings secured by a tape of the same material wound spirally round the leg from the ankle upwards. The head-dress is either a quilted skull-cap or a fur cap with the hair or wool inside, and with a large flap behind which covers the neck and ears. They wear boots of felt with soles of sheep or goat-skin. The women wear a black woollen jacket with a striped parti-coloured petticoat and baggy trousers, and over all a skin coat with the fur turned inside. The hair is arranged in a number of small plaits, and is ornamented by a band of cloth, on which is sown a number of turquoises and beads. The food of the common people consists of thick barley cakes, though those who can afford it eat wheaten bread and drink tea and a fermented liquor called *chung*. The name Kunet seems properly to designate only the mixed race in southern Knaor, but it is used for the population of the central tract in Bisahr, Kulu, Chamba, and Kashtwár, which borders on Ladák. In Chamba we find the Gaddis, who cross over into the neighbouring territory of Kashmir and meet the Thakars or Takkas, the chief cultivating class in those hills, and apparently in the same position with reference to other Hindús as the Játs of the plains. In the valley of Kashmir we have the Kashmiris, and amongst them the servile class called Bátal. To the south-west, along the left bank of the

¹ See section E. of references attached to this chapter.

Jhelum, we have the Musalmán Dogras, called Chibhális, and the Musalmán Sadams of Pūnch. On the right bank of the same river we have Musalmán Gakkars, Satis, and Dunds. To the east of the Chibhális come the Hindu Dogras, and amongst them the servile tribes of the Meghs and Dúms, who are scattered about everywhere and form a considerable part of the population. The Dogras have a light-brown complexion, clearly-cut features, and black hair, which is cut to form a fringe below the turban. The hair is worn on the face. The Thakars are a well-made race, somewhat more powerful in body than the Dogra Rajpúts, whilst the Meghs and Dúms are darker in colour, smaller in limb, shorter in stature, and less bearded. The food and clothing differ in no marked respect from that of the hill-tribes at a similar elevation to the east. Passing to the north-west of the Kashmir valley we come upon the Dárls, an Aryan tribe called Brokpa by the Tibetans, and most of whom are Musalmáns, though the Dal section have adopted the Buddhist religion, language, and customs. They occupy Astor and the trans-Indus Kashmiri district of Gilgit, as well as the neighbouring Kanjúd states of Nagar and Hanza, and the Kashkára states of Chitrál, Yassan, and Mastáj. They are a strongly-built race, with decidedly Aryan features, wearing woollen garments, except among the higher classes, who wear cotton in summer. The ordinary costume consists of trousers, a coat reaching to the knees and confined by a girdle, and a cap of woollen cloth about half a yard long and turned up at the edges until it fits the head, the outward roll thus forming a protection against heat and cold. On their feet they have scraps of leather put under, over, and around the foot, and kept in their place by straps of the same material wound around them. A servile race is also found amongst them, known as Dúms, and performing the same duties as the Dúms of Kumaon. To the south-west of Kashmir, in the salt range, we find the Awáns and Janjúhas, tribes of Aryan origin and of considerable antiquity. From the Káli to the Indus, dialects of Tibetan are spoken to the north. Mr. Drew tells¹ us that from near the Nunkun group of peaks which form the water-parting between the Maru-wadwan and Suru rivers, "and from no other spot in Asia, one may go westward through countries entirely

¹ Northern Barrier of India, p. 20.

Muhammadian as far as Constantinople; eastward among none but Buddhists, to China; and southward over lands where the Hindu religion prevails, to the extremity of the Indian peninsula."

Amongst the great mountain groups comprising the western or trans-Indus extension of the Himálaya, and including the ranges known as the Hindu Kush, Kára-koram, and those connecting them with the Tian-shán, are several petty states, regarding which much has been learned of late.¹ To the south lies the country called Afghánistán, the north-eastern portion of which is included in the western prolongation of the Himálaya, while the south-western part is a mountainous country confluent with the Himálaya on the one hand and extending far in the opposite direction to the table-land of Persia. To the north the boundaries are ill-defined and vary almost from decade to decade. The authority of the ruler of Kabul, in many places, depends on the forces at his disposal to coerce his unwilling subjects; but in 1879, the Afghán province of Turkistán included the whole of the countries between the Hindu Kush and the Oxus, comprising Balkh, Kunduz, and Badakshán, with their dependent states. The seat of the government is Balkh, with cantonments at Faizabad. Karátigin, on the upper valley of the Surkháb, pays tribute to Bukhára; also Darwáz, on the Panja branch of the same river, and Shighnán-e-m-Roshán, on the Ghund and Murgháb rivers. Wakhán is tributary to Badakshán, and south of it lies Kashkára, also called Chitrál. Northern Kashkára, including Yassan and Mastúj, is subordinate to the ruler of southern Kashkára, who resides at Chitrál. North of Gilgit we have the robber states of Hanza and Nagar or Kanjúd. South of Gilgit, in the valley of the Indus below Banji or Bawanji, are a number of small republics, who manage their affairs in assemblies called *sigás*, with which we may compare the similar institutions at the opposite extremity of the Himálaya. Some of these, such as Dúrel and Hodar, owe a nominal subjection to Kashmir. Further south we come to the independent tribes of Buner and Swát. West of the Indus, in the Kunar valley, are the Bajaur and Dir states, and between them and the Hindu Kush range the country of Káfiristan.² It may well

¹ See section F. of references attached to this chapter.
explored by Major Tanner, R.E.

² Lately partially

be supposed that this rugged tract, the meeting-place of the Turanian, Iranian, and Aryan races, presents many points of interest to the ethnographer as well as to the geographer; but here we can but very briefly refer to them, however important they may be.

We have already noticed an Aryan race, the Dárds, occupying Gilgit and forming a part of the population of the neighbouring states. Here they meet the great Iranian race, which, under the name Tájik or Galeha, form the bulk of the cultivating population of the Musalmán states between the Indus and Jaxartes. History tells us that from the third century before Christ to the sixth century after Christ this tract was subject to the continued incursions of a Skythian race, traces of whom may be seen in the Brahúis of Biliuchistán, the Hazárah of the Hindu Kush, the Gujars there and in India, the Gakkars and Kohistánis of the Indus, and the Játs of India, and who have continued to influence the entire history of this tract to the present day. In the eleventh century the Afgháns were a small tribe in the Sulaimán mountains, of no importance and but little known. Since then they have increased so much as to have been able to annex a considerable extent of country, and to impose their language, Pukhtu or Pushtu, on the populations which they have absorbed. Along the Indus, Afgháns occupy the villages as far as Batera in 34° 53' north latitude, where the Kohistán commences. The Afgháns themselves are tolerably fair, robust, and of moderate stature. They have long faces, high cheek-bones, and dark hair, which they wear unshaved. Their underclothing is of cotton, over which they throw a loose coat of woollen cloth, felt, or, more commonly, of sheepskin. They wear low caps on their heads, around which a cloth is twisted to form a turban. Boots are generally worn, and they carry a matchlock, scymitar, and shield. Leavened bread and meat are eaten by all classes. The women are rigorously secluded in the towns, but in the country, beyond the influence of the local mauvi or mullah, much more liberty is allowed. The use of wine is forbidden, but in the hills it is taken in secret, and that made in the Dárel valley has more than a local reputation. Pushtu approaches the Pehlavi or Zendic form of old Persian on the one hand, and the Sindhi form of Prákrit on the other. It is spoken throughout Afghánistán, and, with dialectal variations, in

Bajaur, Panjkora, Dir, and Swát. Afgháns are Sunni Muhammadans.

The Kohistánis of the Indus claim an Arab descent and speak a language different from Pushtu and akin to Kshikári and Dárdui. They are fair and have sandy hair, and are of a robust make. They wear a tight-fitting coat and trousers resembling somewhat those of the European, with a cap of brown woollen material in the form of a bag rolled up at the edges. Around the legs and the feet a goatskin is wound, kept in its place by a strap of leather, the great-toe and the heel being left bare. A matchlock and sword are always carried, and they are expert shots. The women wear a loose jacket and trousers, and a cap of cotton or wool, over which they throw a woollen or cotton sheet when proceeding far from their villages. Unleavened bread of wheat, barley or millet (*Holcus sorghum*) is eaten with vegetables, milk, butter, or stewed meat. Further north the people are less prosperous and more regardless of cleanliness. In some of the villages there are colonies of pastoral Gújars, and in the Yassan villages settlements of Dúms, who supply the musicians and dancers of the country. Following next the Kunar valley, we have dialects of Pushtu spoken in Bajaur and Dir as far as the Lahori pass leading into Chitrál. Here it meets the Kshikára, which is akin to the Dárdui and is of Prákritic origin, though many Persian vocables have been added. The pagan inhabitants of Káfiristan are of the same race as the Kshikáras and the Dárds, and speak a language having an archaic Prákritic origin.

Moving further westward into Badakshán, we come amongst a large Iranian population speaking Pushtu in the south and Persian in the north, and from Narín an Uzbek population speaking Túrki, and the nomad Hazárahhs having a dialect of their own. On the invasion of the Turks the old inhabitants retreated into the more inaccessible valleys of the great ranges, and there founded the Galcha states of Darwáz, Kurátigin, Shighnán, and Wakhán. In Darwáz and the adjoining parts of Badakshán pure Persian is spoken or understood. The vocables and grammar of the Galcha proper show a strong affinity to Dárdui, and many conjecture that, in its pure state, it must have been the intermediate link between the Iranian and Indian branches

of the great Aryan family; that the dispersion took place from the '*bám-i-dunya*,' 'the roof of the world,' the Indian branch proceeding southwards and along the Kábul river to India, whilst the Iranian branch crossed the Pámir to the plains of Turkistán. Towards the plains the Galchas are Sunnis, but in all the hill districts, except Darwáz, they are Shiáhs. The Shiáh is held in the same contempt as the infidel, and the Sunni esteems it a holy and righteous act to capture and sell his less orthodox fellow-believer into slavery. Amongst the tribute paid by the hill states to the Afghán governors of Badakshán and to the Wali of Bukhára, not the least important parts are the troops of fair-complexioned girls from the upper valleys of the Galcha states, and, when procurable, pagan boys from Káfiristán. Uzbeks are chiefly found in the country, and Tájiks in the towns along the plains below Badakshán and in Turkistán; the former invariably speak Túrki and the latter Persian, or dialects with a number of Persian words in them. In Wakhán the men wear brown woollen coats and trousers, Ladáki boots, and a scanty cotton turban, either blue or white. The women here also dress much like the men, and, as in Chitrál, wear their hair in long plaits. They have Jewish noses and are not very fair-looking. Kirghiz are found along the eastern slopes of the Tian-shán and the northern slopes of the Pámir and Kuen-lun ranges in Chinese Turkistán. The Alai Kirghiz of the Pámir have a bad reputation as robbers. East of them lie the great uninhabited steppes until we come to the country near Kanjúd, which is rendered unsafe by the raids of the people of Hanza and Nagar. Further east we have the Chang-thang highlands, occupied by the robber Bhotiya tribes of Changpás, speaking a Tibetan dialect akin to Zanskári. The dress of the male Kirghiz inhabiting the pastoral slopes of the highlands south of Yárkand and Káshgar is not different in any material degree from that of the other inhabitants.¹ Their women wear as a head-dress a white cloth rolled evenly and regularly round a skull-cap of red or other bright material, with lappets over the ears. The end of the turban is drawn down and passed over the lappets and under the chin, and the coat worn is a dressing-gown of wadded and quilted cotton. In the plains the winter dress of the females comprises a fur cap of black lamb's

¹ Gordon: *Roof of the World*.

wool with a trimming of the fur, and a crown of coloured silk or cloth worn over a square of muslin which forms a veil. The coat worn is full and long, and the boots are of embroidered leather with high heels. The hair is worn either in two long plaits or in ringlets. The men wear a close-fitting cap lined with fur and turned up at the bottom. The coat resembles that worn by the women, only it is fastened by a cotton girdle, and the boots are worn long and plain, with felt stockings. In winter, sheepskin with the wool attached is the universal material for clothing.

To Prejevalsky we are indebted for an account of the people around the lake country of the Tarim near **Kára-kurchina.** Lob-nor, to the north of the Altyn-Tágh, and of those inhabiting the eastern extremity of the Kuen-lun near the sources of the Hoang-ho. The former present "a strange mixture of facial types, some of which call to mind a Mongolian race. The prevailing characteristics are, however, Aryan, though far from pure. * * In height they are rather below the average; frame weak and hollow-chested; cheek-bones prominent and chin pointed; beard scanty and à l'Espagnole; whisker even smaller; hair on the face generally of feeble growth; lips often thick and protruding; teeth white and regular and skin dark, whence their name Kára-kurchin may be derived." Their language is said to resemble closely the dialect of Khoten. The clothing of the lake-dwellers is made from the fibres of a species of *asclepius*, and consists of a loose coat and trousers with, in winter, a sheepskin cap, and in summer one made of felt. In summer the feet are uncovered, and in winter shoes of untanned hide are worn. The coats in winter are lined with duckskins dressed in salt. Fish, wild-fowl, and the tender shoots of reeds are their principal food. All profess the Muhammadan religion. The inhabitants of the country along the Tarim towards Korla appear to be also of the same race, though comparatively more civilised.

To the north-east, towards the Yellow river, we have the Kára-Tangutans, a race apparently connected with the Tibetans proper. They are more robust in form, greater in stature, and darker in complexion, than

¹ From Kulja across the Tian-shán to Lob-nor: London, 1879, pp. 44, 166. See, further, section G. of references attached to this chapter.

the Tungutans of Kan-su. Their hair is black, but the head is shaved clean and no pig-tails are worn. The eyes are dark and large, never narrow like the Mongols; the nose straight and sometimes aquiline and also sometimes *retroussée*; the lips thick and protruding; the cheek-bones not so prominent as in the Mongol; the face long and never flat, and the skin tawny coloured. The language is akin to Tibetan. The dress in summer comprises a long, grey, woollen coat reaching to the knees, boots, and a low-crowned, broad-brimmed, felt hat. In winter a sheepskin coat is put next the skin, and the upper part of the legs is usually left bare and also the right arm and part of the right breast, the right sleeve being allowed to hang down empty. The women dress like the men, and all live in tents made of black, coarse, woollen cloth, whence the name Kára (or black)-Tangutans. They are met as far as the Murui-ussu, the extreme point reached by Prejevalsky.¹ Of the people between the Hoang-ho and Brahmaputra very little is known beyond the fact that they are of Tibetan origin and chiefly Buddhists in religion. Pamutan near Bathang, which lies in about 99° east longitude and 28° 50' north latitude, is the most westerly point attained by Mr. Cooper² in his memorable journey from Hankow towards the frontier of India in 1868; and Prun, in the Mishmi country, the most easterly point reached in his attempt to penetrate the intervening ranges to China in 1870, leaving a space of about 120 miles as the crow flies unexplored and unknown.

We have now briefly sketched the character of the countries and peoples encircling the great elevated area which from Ladák on the west to the Chinese frontier is known as Tibet, and is entirely under Chinese influence. The true name of this tract is Bod-yul or Bod-land, and the people Bod-pas, corrupted by the Indians into Bhotiyas, a name now applied to the Tibetans living on the borders between India and Tibet, whilst the people of Tibet proper are called Huniyas and the country Hundes. The eastern division of Tibet is known as Khám or Khám-yul, sometimes called Bod-chen; it extends from the frontiers of China to about 95° east longitude. Central Tibet or Tibet proper is called Bod at its eastern end, and

¹ Mongolia, by Prejevalsky: London, 1876, II., pp. 103, 301.
 a Pioneer of Commerce in Pig-tail and Petticoats: London, 1871.

² Travels of

g-Nári or *Nári* at its north-western end; the former division being the shorter of the two, but perhaps broader and more civilised and populous. A line drawn from Darjiling northwards would apparently separate Bod from *Nári*. The central part is also called U-Tsang from the two provinces of U and Tsang in which *Lhása* is situated. *Nári* is divided into the three great districts of Mang-yul, Khorsum, and Mar-yul.¹ The first marches with *Nepál* almost to its western boundary; the second extends along the British frontier of Kumaun and Garhwál and that of the independent state of *Bisahr*; and the last included western Tibet and the *Kashmíri* states of *Balti* and *Ladák*. The physical characteristics of this tract have had the effect of isolating its inhabitants, who are distinct in race and language, from all the nations we have described, and find their affinities in the *Tangutans* of the north-eastern *Kuen-lun* already noticed. They are broadly built; have dark hair, scanty beards, high cheek-bones, oblique eyes, complexion fair amongst the better classes, dark amongst the lower, who are more exposed to the weather. To the east in the warmer valleys they are agriculturists, and to the west and north follow a pastoral life. The costume varies in the different provinces and with the means of the person, but as a rule the men in the wilder parts shave the head clean like the *Kúra-Tangutans*, whilst the more settled allow it to grow long and plait it into a queue or tail like the *Tangutans* proper. To the east Chinese fashions are in vogue, and to the west the common dress is a coat and trousers of undyed woollen material with boots to the knee. The *Lamas* wear distinctive dresses, red or yellow according to the sect to which they belong, and the wealthier indulge in coloured broad-cloth garments of English or Russian manufacture. Barley porridge, tea and meat form the staple food of the people, and *chang* or beer, a simple infusion of malted barley, is of universal use all over Tibet.

Having completed our review of the nations inhabiting the
 General distribution. *Himálaya* and the surrounding country, we
 may now briefly notice the natural distribution
 of the several great races that have come under our consideration in

¹ From Captain Henry Strachey's paper on the physical geography of Western Tibet, London, 1854, still our best authority on the subject. A more detailed account will be given hereafter in the *Gazetteer* portion of this memoir: see also section H. of references attached to this chapter.

the special tracts that each now occupies. The extraordinary rigour of the climate and the physical barrier that exists between Tibet and India sufficiently explains the absence of ethnical or political relations between the two countries. Accident has given the Indian state of Kashmir political preponderance in western Tibet, but the ethnical distinction still remains, and will probably ever continue. The climatic condition of eastern and northern Tibet allows of a free migration of the inhabitants from one part to the other, which is shewn in the common origin of the people of those countries to the present day. Turks and Tartars occupy the country called Turkistán, similar in character to their original homes; but in proportion as the climatal conditions become Indian, so does the population become Aryan. The Tartar is the child of the rugged bleak steppes; and when we approach the cold and wooded mountains we come on the Hindu element at its maximum in the tract around the sacred sources of the Jumna and the Ganges, gradually diminishing as we move eastward towards the excessively moist though warmer valleys of the Nepál and Asám Himálaya, where they are replaced by races akin to the people of eastern Tibet and Siam, and on the west meeting the Iranian and the Tartar in the trans-Indus ranges. To the north of the Kuen-lun the Mongol and Chinese convert to Muhammadanism, called Tunganis, and in western Tibet the Muhammadan Baltis, divide the Buddhists from the followers of Islám. In the valleys of the affluents of the Oxus and the Kábul river we have an Iranian race of Galchas or Tájiks driven upwards by political disturbances and mixing with an indigenous mountain race of Aryan extraction. The rigorous climate and scanty cultivation which prevents the intrusion in any numbers of a southern race also debars the tribes inhabiting the higher hills from making any permanent occupation of the lowlands. Like their favourite domestic animal the yak, the Himálayan mountaineers do not thrive at low altitudes, nor can plains-bred men or animals withstand for any time the arctic cold and rarefied air of the more elevated regions. Thus, there is a clear connection between the distribution of the nations that are found in the Himálaya and the physical characteristics of the regions that they occupy; and if we had time to pursue the subject further, it might be shown that the orographical conditions of a tract have materially influenced its history, political

and religious, and the social and moral character of its inhabitants. The disposition of a people towards peace or war ; their migrations ; the diffusion of their language ; their habits, pastoral, agricultural, or commercial ; the extent of their influence—all depend more or less on the physical peculiarities of the country that they inhabit. And not only is man so affected, but the entire fauna and flora obey the same laws, so that the skilful naturalist can from a plant or even a butterfly describe the general character of the country of which it is a native, and with it the customs and manners of the inhabitants. For the nature of a country, whether mountainous or level, the direction of the great ranges, the length and line of coast, the position with regard to the equator, the relations of land and water, and the drainage systems, are all primary agents in the distribution of organic life and of the influences which govern all atmospheric and climatic phenomena.

REFERENCES.

The following list of references to books bearing on the geographical and ethnological notices in the preceding chapter is not intended to be exhaustive, but simply to furnish a guide to the bibliography of the countries within and adjoining the Himalaya, and only includes such works as are ordinarily procurable.

Section A.—Eastern Asam and Tibet.

(a)—INDO-CHINESE FRONTIER.

1. La Mission du Thibet, by Fr. C. H. DesGodins. Verdun, 1872.
 2. Pays frontières du Thibet, de la Burmanie et de Yunnan, by the same. Bull. de la Soc. de Geogr. de Paris, Oct., 1876.
 3. Territoire de Bathang, by the same. *Ibid*, Dec., 1876, and Dec., 1877.
 4. Le cours supérieur des fleuves de l'Indo-Chine, by the same. *Ibid*, Aug., 1876.
 5. Notes sur le Thibet, by the same. *Ibid*, Oct., 1877.
 6. Mandalay to Momein, by J. Anderson. London, 1876.
 7. Overland route to China *via* Asam, the Tenga Pani river, Khamti and Singpho country, across the Irrawaddy river into Yunnan, by H. Cottam. P. R. G. S., XXI., 6.
 8. Travels in Western China and on the Eastern borders of Tibet, by Capt. W. J. Gill. *Ibid*, XXII., 255, and see H., 26.
 9. Across China from Chin-Kaing to Bhamo in 1877, by J. McCarthy. *Ibid*, I. N. S., 489.
 10. Official narrative of the expedition to explore the trade routes to China *via* Bhamo under Major Sladen. Calcutta, 1870.
 11. Travels of a Pioneer of Commerce in Pigtail and Petticoats: an overland journey from China towards India, by T. T. Cooper. London, 1871.
 12. Journal of an attempted ascent of the river Min to visit the tea-plantations of the Fukhien Province of China, by G. J. Gordon. J. A. S., Ben., IV., 553.
 13. Short survey of the country between Bengal and China, by Baron Otto des Granges. *Ibid*, XVII., i., 132.
 14. Memorandum on the countries between Thibet, Yunnan, and Burmah, by Rev. T. D'Masure. *Ibid*, XXX., 367.
- (See also the writings of Blakiston, Margary, Fytche, Pollok, Phayre, &c.)

(b)—UPPER ASAM.

15. Memoir of a survey of Asam and the neighbouring countries executed in 1825-26, by Lieut. R. Wilcox. As. Res., XVII., 436, and Sel. Rec., Bengal, XXIII., 1855, and Sel. papers, Hill tracts, 1873, p. 1.
16. Account of Asam, the local geography, tea-plant, manners of the neighbouring tribes, by W. Robinson. Calcutta, 1841.
17. Report on the Eastern frontier of British India, by Capt. R. B. Pemberton. Calcutta, 1835.
18. Journals of travels in Asam, Burmah, Bootan, and Afghanistan, by W. Griffiths. Calcutta, 1847.
19. Memoir on Sylhet, Kachár, and the adjacent districts, by F. H. Fisher. J. A. S., Ben., IX., 808.
20. Notes on northern Cachár, by Lieut. R. Stewart. *Ibid*, XXIV., 552.

21. Travels and adventures in the Province of Asám, by Major J. Butler. London, 1855.
22. Koch Bihár, Koch Hájo, and Asám in the 16th and 17th centuries, by H. Blochmann. J. A. S., Ben., XLI., i., 49.
23. History of Asám, by Anundorum Boruah. Calcutta.
24. Memorandum on various phenomena in Asám, by the Rev. N. Brown, J. A. S., Ben., XII., 909.
25. Notes on the Asám temple ruins, by Capt. E. T. Dalton. *Ibid*, XXIV., i.
26. Description of ancient temples and ruins at Chárdiwar in Asám, by G. E. Westmacott. *Ibid*, IV., 185.
27. Note on ancient temples and other remains in the vicinity of Suddyah in Upper Asám, by Major S. F. Hannay. *Ibid*, XVII., i., 459.
28. Ruins at Dimápur on the Dunsiri river, Asám, by Major Godwin-Austen. *Ibid*, XLIII., 1.
29. Temple at Jayságar in Upper Asám, by J. M. Foster. *Ibid*, 311.
30. Contributions to the geography and history of Bengal, by H. Blochmann. *Ibid*, XLII., 209 : XLIII., 280.
31. Report of a visit to the hills in the neighbourhood of the Soobansiri river, by Lieut. S. F. Hannay. *Ibid*, XIV., 250.
32. Account of a visit to the Jugloo and Secsee rivers in Upper Asám, by the same. *Ibid*, XXII., 511.
33. Report of an expedition into the Mishmee hill to the north-east of Sudyah, by Lieut. E. A. Bowlett. *Ibid*, XIV., 477.
34. Journal of a visit to the Mishmee hills in Asám, by W. Griffith. *Ibid*, VI., 325, and *Sel. papers*, Hill tracts, 110.
35. Notes on the language spoken by the Mi-Shmis, by W. Robinson. *Ibid*, XXIV., 307.
36. Note on the Dophlas and the peculiarities of their language, by W. Robinson. *Ibid*, XX., 126.
37. Correspondence and journal of Capt. Dalton in a visit to a clan of Abors on the Dihing river. *Sel. Res.*, Ben., XXIII., 139.
38. On the Meris and Abors of Asám, by the same. J. A. S., Ben., XIV., 426.
39. Account of the mountain tribes on the extreme north-eastern frontier of Bengal, by Capt. Jenkins, edited by J. McCosh : containing notices of the Miris, Abors, Mishmis, Kangtis, Singhphos, Muamarias, and Nágas. *Ibid*, V., 193.
40. Notes on the tribes of the eastern frontier, by J. H. O'Donel. *Ibid*, XXXII., 400-404, and by H. J. Reynolds. *Ibid*, 407.
41. Memorandum on the north-eastern tribes, by A. Mackenzie. Calcutta, 1869.

(c)—ETHNOGRAPHY.

42. Descriptive Ethnology of Bengal, by E. T. Dalton : Calcutta, 1872. [Gives an account (with figures) of the Khamtis ; Singhphos, Mishmis, Chulikáta Mishmis ; Abor group, including Fádams, Miris, Dophlas, and Akas ; Nágas of Upper Asám ; lower Nágas, including Kukis, Mainpuris, and Koupouls ; Mikirs ; Jaintiyas and Khasiyas ; Gáros, Kacháries ; Mechs ; Kochs ; Bhotiyas ; Lepchas ; Murmis ; Limbus and Kirántis ; Hayas, and the Tipura (Tipperah) and Chittagong tribes].

(d)—GARO HILLS.

43. Observations on the inhabitants of the Garrow hills, by J. Elliot. *As. Res.*, III., 17.
44. A narrative of our connection with the Dushani and Chianni Garrows, with a short account of their country, by C. S. Reynolda. *J. A. S., Ben.*, XVIII., 1, 45.
45. Some account of the Cásiah hills. *Gl. in sec. I.*, 252.
46. Notes on the Cásiah hills, by H. Yule. *J. A. S., Ben.*, XIII., 612.
47. A vocabulary of the Garo and Konch languages, by Lieut. W. J. Williamson. *Ibid.*, XXXVIII., 14.
48. Khasia grammar, selections and vocabulary, by W. Pryce. Calcutta, 1855.
49. English-Khasia vocabulary, by Kelsall. Calcutta, 1864. See also Hodgson's papers.
50. Some account of the Pális of Dinajpur, by G. H. Damant. *Ind. Ant.*, I., 336, 371.
51. On the dialect of the Pális, by the same. *Ibid.*, II., 101.
52. The Garos, by the Rev. I. J. Stoddard. *Ibid.*, II., 336.
53. On the Khasia hill-tribes of north-eastern Bengal, by A. Morgan. *P. Lit. and Phil. Soc., Liverpool*, 30, 1875-76.

Section B.—Southern Asam.

1. Report on the wild tribes of the Chittagong frontier, by C. Ricketts. Calcutta, 1847.
2. Some account of the hill-tribes in the interior of the district of Chittagong, by the Rev. M. Barbe, J. A. S., Ben., XIV., 380.
3. Selection of papers regarding the hill tracts between Asám and Burmah and on the Upper Brahmaputra. Calcutta, 1878.
4. The wild tribes of the south-east frontier, by Capt. T. H. Lewin.
5. The Chittagong hill tracts and the dwellers therein, with comparative vocabularies of the hill dialects, by the same. Calcutta, 1869.
6. History of Arakan, by Col. J. P. Phayre. *J. A. S., Ben.*, XV., 232.
7. Account of Arakan, by the same. *Ibid.*, X., 679.
8. History of Pegu, by Sir A. Phayre. *Ibid.*, XLII., 23, 120 : XLIII., 6.
9. Notes on the Heuma or Shendoos, a tribe inhabiting the hills north of Arracan, by Capt. Tickell. *Ibid.*, XXI., 207.
10. History of the Burmah race, by Sir A. Phayre. *Ibid.*, XXXVII., i. : XXXVIII., 29.
11. The hill tribes of the north-east frontier. *Ind. Ant.*, I., 63.
12. The Lushais, by Capt. Badgley. *Ibid.*, II., 363.
13. Abstract of the journal of a route travelled by Capt. S. F. Hannay from the capital of Ava to the amber mines of the Hukong valley on the south-eastern frontier of Asám, by Capt. R. B. Pemberton. *J. A. S., Ben.*, VI., 245, and No. 3 above, p. 23.
14. Journey from Upper Asám towards Hookhoon, Ava, and Rangoon, by W. Griffiths. No. 3 above, p. 125.
15. Narrative of a journey from Ava to the frontier of Asám and back in 1836-37, by G. T. Bayfield. *Ibid.*, 134.

16. Notes on a trip across the Patkoi range from Asám to the Hukoong valley, by H. L. Jenkins in 1869-70. *Ibid*, 245.
17. Notes on the Burmese route from Asám to the Hukoong valley, by the same. *Ibid*, 250.
18. The Nágas of Asám, by Capt. Jenkins. J. A. S., Ben., V., 207.
19. Extract from the narrative of an expedition into the Nága territory of Asám in 1840, by Lieut. Grange. *Ibid*, VIII., 445 : IX., 947.
20. Extracts from a report on a journey into the Nága hills in 1844, by B. Wood. *Ibid*, IX., 947 : X., 126 : XIII., 771.
21. Extract from a memoir on some of the natural products of the Angámi Nága Hills, by J. W. Masters. *Ibid*, XIII., 707 : XVII., i., 57.
22. Narrative of a tour over that part of the Nága hills lying between the Diko and Dyang rivers, by Capt. Brodie. *Ibid*, XIV., 828.
23. Report of Lieut. Brodie's dealing with the Nágas on the Sibaágar frontier in 1841-46. Sel. papers, Hill tracts, p. 284.
24. Note on a visit to the tribes inhabiting the Hills to the south of Sibaágar in Asám, by S. E. Peal. J. A. S., Ben., XLI., i., 9 : and Sel. papers, Hill Tracts, p. 316.

(f)—LANGUAGES.

25. A rough comparative vocabulary of some of the dialects spoken in the Nága Hills, by Capt. J. Butler. J. A. S., Ben., XLII., i., App. i. Vocabulary of the Banpará Nágas, by S. E. Peal. *Ibid*, xxx.
26. Account of the valley of Manipur and of the Hill tribes, by Major McCulloch. (To which is added a comparative vocabulary of the Munniporee, Undro Bengmai, Chairel, Meeyang, Koupoee Poveeron, Koupoee Songbo, Qoireng, Khoonggoee, Phudang, Koopome, Tukamec, Muram, Murring Anal Namfau, Kookie, Burmese and Shan languages). Sel. Res., Government of India, For. Dept., XXVII., 1859.
27. Analysis of the Bengali poem 'Rájámá,' or chronicles of Tripura, by the Rev. J. Long. J. A. S., Ben., XIX., 533.
28. Progressive colloquial exercises in the Lushai dialect of the Dzo or Kúki language, by Capt. T. H. Lewin. Calcutta, 1874.
29. Hill Proverbs : a collection of wit and wisdom from the people of the Chittagong Hills, by the same. Calcutta, 1875.
30. Notes on the languages spoken by the various tribes inhabiting the valley of Asám and its mountain confines, by W. Robinson. J. A. S., Ben., XVIII., i., 183, 310.
31. A dictionary in Asámesé and English, by M. Brouson, American Baptist Missionary. London, 1876.
32. Note on the Shans, by Col. S. F. Hannay. Calcutta, 1846.
33. A visit to Xiengmai, the principal city of the Shan State, by Sir R. H. Schomburg. J. A. S., Ben., XXXII., 387 (1863).
34. Introductory sketch of the history of the Shans in Upper Burmah and Yunan, by N. Elias. Calcutta, 1876.
35. A grammar of the Shan languages, by J. Cushing. London, 1877.
36. Alphabets of the Tai languages, by Rev. Nathan Brown, with plates of those of the Ahom, Khamti, Shyán, Laos, and Burmese. J. A. S., Ben., VI., 17.

37. Interpretation of an Ahom inscription, by Capt. Jenkins. *Ibid*, 980.
38. Comparison of Indo-Chinese languages with comparative vocabularies of Bangali, Asamese, Khamti, Siamese, Aka, Abor, Mishimi, Barmese, Karen, Singpho, Jili, Gáro, Manipuri, Songpu, Kapwi, Koreng, Marám, Champhung, Lahuppa, N. Tángkhul, C. Tángkhul, S. Tángkhul, Khoibu, Maring, Anamese, Japanese and Corean, by Rev. W. Brown. *Ibid*, 1,023.
39. Description of the tomb of an Ahom noble, by C. Clayton. *Ibid*, XVII, i., 473.
40. Note on Ghargaon in Asám, by J. M. Forster. *Ibid*, XLI, i., 32.
41. Note on the Singphos, by Col. S. F. Hannay. Calcutta, 1846.
42. Report of a visit by Capt. Vetch to the Singpho and Naga frontier of Lakimpur in 1842. Sel. papers, Hill tracts, p. 255.
43. Remarks on the Indo-Chinese alphabets, by Dr. Bastian. J.R.A. S., III, ii.

Section C.—Bhutan, Sikkim, Nepal.

(a)—BHUTÁN.

1. Narratives of the Mission of George Bogle to Tibet (1774), and of the journey of Thomas Manning to Lhása (1811-12), by C. R. Markham. London, 1876.
2. Notes on Dr. Hamilton's Missions to Panákhá in 1775 and 1777.
3. Account of an embassy to the court of the Teshoo Lama in Tibet and a journey through Bootan, by Capt. S. Turner. London, 1806.
4. Poorungeer Gosseyn's residence at Lhása in 1785. As. Res., I, 207.
5. Account of Bhutan, by Kishan Kant Bose. As. Res., XII, 128.
6. Report on Bootan, by Capt. R. B. Pemberton. Calcutta, 1839. Journal of the Mission to Bhutan, 1837-38, under Capt. R. B. Pemberton, by W. Griffiths, M.D. Calcutta, 1847, and J. A. S., Ben., VIII, 208, 251.
7. Report on the state of Bootan and on the progress of the Mission of 1863-64, by the Hon'ble Ashley Eden. Calcutta, 1864.
8. The last four are reprinted in 'Political Missions to Bootan.' Calcutta, 1865.
9. Bootan, or the story of the Dooar war, by Surgeon Rennie, M.D. London, 1866.
10. Papers relating to Bootan presented to Parliament. No. 13 of 8th February, 1865.
11. Further papers relating to Bootan presented to Parliament. No. 47 of 15th February, 1865.
12. Visit to Dewangiri, by E. T. Dalton. J. A. S., Ben., XVIII, 66.
13. A Military report on the country of Bhutan, containing all the information of Military importance which has been collected up to 12th July, 1866, by Lieut. C. M. Macgregor. Calcutta, 1873.

(b)—SIKKIM.

14. Himalayan Journals: notes of a naturalist in Bengal, the Sikkim and Nepal Himalayas, &c., by J. D. Hooker. London, 1855.
15. Notes, chiefly botanical, made on a journey from Darjeeling to Tonglo, by the same. J. A. S., Ben., XVIII, i., 419.
16. Notes on a tour in the Sikkim Himalayan mountains undertaken for the purpose of ascertaining the geological formation of Kinchinjanga, by W. S. Sherwill. *Ibid*, XXII, 540, 611.

17. Journal of a trip undertaken to explore the glaciers of the Kanchinjunga group, by Major J. L. Sherwill. *Ibid*, XXXI., 457.
18. Diary of a journey through Sikkim to the frontiers of Thibet, by A. Campbell. *Ibid*, XXI., 407, 477, 563.
19. A journal of a trip to the Sikkim, by the same. *Ibid*, XVIII., i., 482.
20. Routes from Darjiling to Tibet, by the same. *Ibid*, XVII., ii., 488.
21. Itinerary from Phari in Tibet to Lasse, with appended routes from Darjiling to Phari, by the same. *Ibid*, XVII., i., 257.
22. Notes on eastern Thibet, by the same. *Ibid*, XXIV., 252.
23. Report on a visit to Sikkim and the Tibetan frontier in 1873, by J. Ware-Edgar, C.S.L. Calcutta, 1874.
24. Sikkim, with hints on mountain and jungle warfare, exhibiting also the facilities for opening commercial relations through the hills of Sikkim with Central Asia, Tibet, and Western China, by Col. J. C. Gawler. London, 1878.
25. A short account of the Cooch Behar State, by Capt. T. H. Lewin.
26. The Indian Alps and how we crossed them. London, 1876.
27. Note on the Mechis, together with a small vocabulary of the language, by Dr. A. Campbell. J. A. S., Ben., VIII., 623.
28. Note on the Lepchas of Sikkim, with a vocabulary of their language, by the same. *Ibid*, IX., 379.
29. A note on the Limboos and other hill tribes hitherto undescribed, by the same. *Ibid*, IX., 595.
30. On the literature and origin of certain hill tribes, by the same. *Ibid*, XI., 4.
31. Note on the Limboo alphabet of Sikkim, by the same. *Ibid*, XXIV., 202.
32. Ethnology of India, by Mr. Justice Campbell. *Ibid*, XXXV., 1.
33. Specimens of the languages of India, including tribes of Bengal, the Central Provinces, and the eastern frontier, by Sir G. Campbell. London, 1874.
34. A sketch of the modern languages of the East Indies, accompanied by two language maps, by R. N. Cust. London, 1878.
35. A comparative dictionary of the languages of India and High Asia, by W. W. Hunter. London, 1868.
36. A grammar of the Bong (Lepcha) language as it exists in the Dorjeling and Sikkim hills, by G. B. Mainwaring. Calcutta, 1876.

(c)—NEPÁL.

37. An account of the kingdom of Nepál, by Father Guiseppe. As. Res., II., 307.
38. An account of the kingdom of Nepál, being observations made during a mission to that country in the year 1793, by Colonel Kirkpatrick. London, 1811.
39. An account of the kingdom of Nepál and of the house of Gorkha, by Francis Hamilton (formerly Buchanan). Edinburgh, 1819.
40. Narrative of a five years' residence at Nepál, from 1841 to 1845, by T. Smith. London, 1852.
41. A journal to Kathmandu with the camp of Jang Bahadoor, by Lawrence Oliphant. London, 1852.
42. History of Nepál, translated from the Parbatiya by Munshi Shew Shankar Singh and Pandit Shri Gnanand and edited by D. Wright, M.D. Cambridge, 1877.

43. *Nepál: its people, religion, customs, zoology, botany*, by Dr. Oldfield. London, 1879.
44. *Memorandum relative to the seven Kosis of Nepál*, by B. H. Hodgson. J. A. S., Ben., XVII., II., 646.
45. *Route from Kathmandu to Darjiling*, by the same. *Ibid*, 634.
46. *A cursory notice of Nayakot*, by the same. *Ibid*, IX., 1114.
47. *Selections from the records of the Government of Bengal*, No. XXVII.
48. *Quality of the Kosis of Nepál*. J. A. S., Ben., V., 369.
49. *Papers on the colonisation, commerce, geography, &c., of the Himalaya mountains and Nepál*, by B. H. Hodgson. London, 1857.

(d)—ETHNOLOGY.

50. *Notices of the languages, literature, and religion of Nepál and Thibet*, by B. H. Hodgson. As. Res., XVI., 409.
51. *On the aborigines of the Himalaya*, by the same. J. A. S., Ben., XVI., 1235, and XVII., I., 78.
52. *Origin and classification of the military tribes of Nepál*, with a tabular view of the tribes, by the same. *Ibid*, II., 217.
53. *On the Chepáng and Kústúnda tribes of Nepál*, with a Chepáng vocabulary, by the same. *Ibid*, XVII., II., 650.
54. *Cursory notice of Náyakot and the remarkable tribes inhabiting it*, by the same. *Ibid*, IX., 1114.
55. *On the tribes of northern Tibet, Horsok and Sítán*, by the same. *Ibid*, XXII., 121.
56. *On the colonisation of the Himalaya by Europeans*. Sel. Rec. Ben., XXVII., p. 1.
57. *On the commerce of Nepál*. *Ibid*, p. 11.
58. The preceding seven papers have been reprinted in 'Essays on the languages, literature, and religion of Nepál and Tibet.' London, 1874.
59. *Classification of the Newárs or aborigines of Nepál Proper*, preceded by the most authoritative legend relative to the origin and early history of the race, by the same. J. A. S., Ben., III., 215.
60. *Comparative vocabulary of the several languages and dialects of the eastern Sub-Himalaya from the Káli or Ghagra to the Dhanari*, by the same. *Ibid*, XVI., 1244.
61. *Ethnography and geography of the Sub-Himalayas*, by the same, with a Khas vocabulary. *Ibid*, XVII., I., 544.
62. *On the Tibetan type of mankind*, by the same. *Ibid*, XVII., II., 222, 580.
63. *A brief note on Indian Ethnology*, by the same. *Ibid*, XVIII., 238.
64. *On the aborigines of north-eastern India*, by the same, with a comparative vocabulary of Tibetan written and spoken, Dhimál, Bodo and Gáro. *Ibid*, 451.
65. *On the origin, location, creed, customs, character and condition of the Kooch Bodo and Dhimál people*, with a general description of the climate they dwell in, by the same. *Ibid*, 702. Reprinted with the preceding in the *Aborigines of India*. Calcutta, 1847.
66. *On the aborigines of the eastern frontier with vocabularies of Burman written and spoken, Singpho, Naga, Abor-miri and Miri*, by the same. J. A. S., Ben., XVIII., 967.

67. On the aborigines of southern India, by the same. *Ibid*, 350.
 68. On the aborigines of Central India, by the same. *Ibid*, XVII., ii., 550.
 69. On the aborigines of the north-eastern frontier, by the same. *Ibid*, XIX., 309.
 70. Indo-Chinese borderers and their connection with the Himálays and Tibetana, by the same. *Ibid*, XXII., 1.
 71. Caucasian and Mongolian affinities, by the same. *Ibid*, XXII., 26.
 72. Comparative vocabulary of the languages of the broken tribes of Nepál: [(1) Dahi or Darhí; (2) Denwar; (3) Pahi or Pahar; (4) Chopáng; (5) Bhrámu; (6) Váyu or Háyu; (7) Kuswar; (8) Kusmá, (9) Páthya (unbroken); (10) Thakya (do.); (11) Tháru], and comparative vocabulary of the (seventeen) dialects of the Kirántí language, also a Vayu and a vocabulary and grammar, by the same. *Ibid*, XXVI., 317; XXVII., 303.
 73. On the Váyu tribe, by the same. *Ibid*, XXVII., 443.
 74. On the Kirántí tribe, by the same. *Ibid*, 446.
 75. The Magar language of Nepál, by J. Beames. J. R. A. S., IV., 1869.
- NOTE.—Articles Nos. 44, 45, 62, 64—66, 68, 69, 70—74 above have been reprinted in Miscellaneous Essays relating to Indian subjects, by B. H. Hodgson, London, 1880.

Section D.—Kali to the Satlaj.

1. Illustrations of the botany and other branches of the Natural History of the Himálayan mountains, by J. Forbes Lloyd. London, 1839.
2. Notes of an excursion to the Pindari Glacier in 1846, by E. Madden. J. A. S., Ben., XVI., 226, 596 (chiefly botanical).
3. The Tarree and outer mountains of Kumaon, by the same. *Ibid*, XVII., i., 349; XVIII., i., 603 (chiefly botanical).
4. Description of the (so-called) mountain tract of Kumaon, by J. McClelland. *Ibid*, IV., 39.
5. Notes of observations on the Bokssas, by Dr. J. L. Stewart. *Ibid*, XXXIV., ii., 147.
6. Ancient remains in the Garhwál Bhábar, by H. Pollard. *Ibid*, XXXVI., 156.
7. Narrative of a survey for the purpose of discovering the source of the Ganges, by Capt. F. V. Baker. *As. Res.*, XI., 446.
8. Pitt Rivers's wanderings in the Himala, by Barron. *Agri*, 1844.
9. Report on the Bhotia mahals of Kumaon, by G. Traill. J. A. S., Ben., II., 551. Reprinted in Kumaon Reports.
10. A few notes on the subject of the Kumaon and B.ilkhand Tará. by J. H. Batten. *Ibid*, XIII., 887.
11. Extract from the journal of J. Weller on a trip to the Bulcha and Oonta Dhoora passes, edited by the same. *Ibid*, XII., 78.
12. Journal of a visit to Mclum and the Oonta Dhoora pass in Jawáhir, by E. Maason, edited by the same. *Ibid*, XI., 1157.
13. Note of a visit to the Niti pass of the grand Himálayan chain, by the same. *Ibid*, VII., 310.
14. Notice of a trip to the Niti pass. *Ibid*, XIX., 79.
15. Account of a visit to the Biáns pass. *Gl. in Sci.*, I., 97.

16. Notes on a trip to the Kedarnath and other parts of the snowy range of the Himálays in 1853 (by J. M.) Edinburgh, 1856.
17. The Himálaya in Kumaon and Garhwál. Col. Rev., XVIII., 72.
18. Kumaon and its Hill stations. *Ibid*, XXVI., 373.
19. On the Physical geography of the Himálaya, by B. H. Hodgson. J. A. S. Ben., XVIII. ii., 761.
20. On the Physical geography of the provinces of Kumaon and Garhwál in the Himálaya mountains and of the adjoining parts of Tibet, by Capt. (now General) R. Strachey. J. R. G. S., XXI., 57 (1851).
21. Explanation of the elevation of places between Almora and Gaugri. by Lieut. H. Strachey. J. A. S., Ben., XVII., ii., 527.
22. Narrative of a journey to Cho Lagan (Rakas Tal), Cho Mapan (Manasarowar), and the valley of Pruang in Gnari Hundes, by the same. J. A. S., Ben., XVII., ii., 98, 127, 327.
23. A description of the glaciers of the Pindur and Kuphinee rivers in the Kumaon Himálays, by the same. *Ibid*, XVI., 794, 1002.
24. Notice of Lieut. H. Strachey's scientific inquiries in Kumaon and Garhwál. *Ibid*, XIX., 239.
25. Note on the motion of the glacier of the Pindur in Kumaon, by Lieut. H. Strachey. *Ibid*, XVII., ii., 203.
26. A Sketch of the mountains and river-basins of India, by Trelawny Saunders. India Office, London, 1870.
27. Trans-Himálayan missions and their results. Col. Rev., Jan. 1877, 115.
28. Geographical Magazine (London), IV., 113 (May, 1877), by C. R. Markham, on "the Himálayan system," and p. 173, by T. Saunders, on the same subject, being a reply to the preceding article in the *Calcutta Review*.
29. A memoir on the Indian Surveys, by C. R. Markham. London, 1878.
30. On the snow-line in the Himálays, by Lieut. H. Strachey. J. A. S., Ben., XVIII., i., 287.
31. On the limits of perpetual snow, by Capt. J. D. Cunningham. *Ibid*, XVIII., ii., 694.
32. Remarks on the snow-line, by Capt. T. Hutton. *Ibid*, 954.
33. Reports on the trans-Himálayan explorations in connection with the Great Trigonometrical Survey of India, 1866-67, 1867, 1868, 1869, 1870, 1871, 1872, 1873-75, 1876, 1877, viz.—
 - (1.) Route-survey from Nepál to Lhása, and thence to the sources of the Brahmaputra, made in 1865-66. As well as a route survey through western Nepál and from Niti to Gartokh and back.
 - (2.) Route-survey from Mánn to Totling, thence to Demchok and along the northern branch of the Indus to Thok Jalung and back to Milam, in 1867.
 - (3.) Surveys from Rudok by the gold fields to King-lo, and thence to the Manasarowar lake; from Manasarowar to Shigatze and from the Tadam Monastery to Muktiánath in Nepál; also along the Tingri maidan in the eastern Himálays, 1868.
 - (4.) Survey from Kabul to Badakshan and through the upper valley of the Oxus to Kashgar and back by the Karakoram, 1867-68.

- (5.) Survey from Peshawar to Fyzabad in Badakshná, 1870.
- (6.) Exploration of the Namcho or Tengri Nur lake in Tibet, 1872, and Havildar's journey from Kabul to Bukhara, 1872.
- (7.) Account of the survey operations in eastern Turkistán, 1873-74, in connection with the Yarkand Mission, by Capt. H. Trotter, R.E.
- (8.) Survey from Badakshán to Koláb, Darwaz, and Kubadian to Belakh in 1873.
- (9.) Journey from Jalálabad to Sarhadd-i-Wakhan by the Baroghil pass in 1873.
- (10.) Journey from Leh in Ladákh to Lhása, and thence by Chetang to Odalguri near Tespur in Asám, July 1873 to March, 1875.
- (11.) Indus river from Amb to Bunji; Ynasin to Mastuj and thence to Barkulti; Sar Laspur to Dodbáh near Dir; Miankalai to Pachat near New Kunar by Nawagi and Nawagi to Abazai in 1876.
- (12.) Mr. E. C. Ryall's exploration of Hundes to the north of Kumaon and Garhwál, 1877.

Account of the operations of the G. T. S., India, Vol. I. (1870), II. to IV. (1879). Dehra Dún, 1879.

Section E.—Satlaj to the Indus.

1. Journal of a tour through part of the snowy range of the Himálaya mountains and to the sources of the rivers Jumna and Ganges, by J. B. Fraser. London, 1820.
2. Travels in the Himálayan provinces of Hindustan and the Panjáb; in Ladákh and Kashmir; in Peshawar, Kábul Kundus and Bokhára, by W. Moorcroft and G. Trebeck, from 1819 to 1825, edited by H. H. Wilson. London, 1841.
3. Account of Koonawar in the Himálayas, by Capt. A. Gerard. London, 1841.
4. Notes on Moorcroft's travels in Ladákh and on Gerard's account of Koonawur, by Capt. J. D. Cunningham. J. A. S., Ben., XIII., 172, 223.
5. Journal of a trip through Kulu and Lahul to the Chumureri lake in Ladákh in 1846, by Capt. A. Cunningham. *Ibid.*, XVII., i., 201.
6. Kashmir and das Reich der Siek, by C. Von Hugel, translated into English by Major Jervis. London, 1846.
7. Notes of a visit to the valley of Kashmir in 1836, b. Baron Hugel. J. A. S., Ben., V., 184, 303.
8. Travels in Kashmir, Ladákh, Iskardo, the countries adjoining the mountain course of the Indus, &c., by G. J. Vigne. London, 1842.
9. Notice by Vigne of his travels in Kashmir. J. A. S., Ben., V., 56.
Description of the valley of Kashmir, Ghazni and Kábul, by G. J. Vigne. *Ibid.*, VI., 766.
10. Ladákh, physical, statistical, and historical, with notices of the surrounding countries, by Major A. Cunningham. London, 1854.
11. Abstract journal of the routes to the sources of the Panjáb rivers, by the same and A. Broome. J. A. S. Ben., X., 105.
12. Correspondence of the Commissioners deputed to the Tibetan frontier, by Captain A. Cunningham. *Ibid.*, XVII., i., 89.
13. Memorandum on the boundary between the territories of Mahárája Gulab Singh and British India as determined by the Commissioners, by J. D. Cunningham. *Ibid.*, XVII., i., 295.

14. Memorandum on the district of Bussahir and the pacification of the disaffected portion of the inhabitants, by G. C. Barnes. Lahore, 1869.
15. Journal of a tour through Hungrung, Kunawur and Spiti in 1838, by T. Hutton. *Ibid*, VIII., 901: IX., 489, 555.
16. Journal of a trip to the Burenda pass, by the same. *Ibid*, VI., 901.
17. Diary of an excursion to the Boorun and Shatool passes over the Himálaya, by E. Madden. *Ibid*, XV., 79.
18. Journey from Soobathoo to Shipke in Chinese Tartary, by J. G. Gerard. *Ibid*, XI., 363.
19. Report on the valley of Spiti, by the same. *Ibid*, X., 198; and by W. C. Hay. *Ibid*, XIX., 429; see also *As. Res.*, XVIII., 239.
20. Notes of a trip from Simla to the Spiti valley and Chomoriri lake in 1861, by W. Theobald. *Ibid*, XXXI., 480.
21. Illustrations of ancient buildings in Kashmir, by H. H. Cole. London, 1869.
22. Kashmir, the Western Himalaya, and the Afghan mountains, by Dr. A. M. Verchere. *J. A. S., Ben.*, XXXV., ii., 89, 169.
23. Western Himalaya and Thibet: a narrative of a journey through the mountains of northern India during the years 1847-48, by T. Thomson, M.D. London, 1852.
24. Voyage dans l'Inde, by V. Jacquemont. Paris, 1841.
25. Letters from India, describing a journey in the British dominions of India, Tibet, Lahore, and Cashmere during 1828-31, by the same. London, 1834.
26. Notes taken in 1829 relative to the territory and the Government of Iskardo, by C. M. Wade. *J. A. S., Ben.*, IV., 589.
27. Notes on Iskardo, by Vigne. *Ibid*, V., 56, 308: VI., 773.
28. The abode of snow: observations on a journey from Chinese Tibet to the Indian Caucasus, through the upper valleys of the Himálaya, by A. Wilson. Edinburgh, 1876.
29. The Jummoo and Kashmir territories: a geographical account, by F. Drew. London, 1876.
30. The northern barrier of India, a popular account of the Jummoo and Kashmir territories, by F. Drew. London, 1877.
31. A Gasetteer of Kashmir and the adjacent districts of Kishtwar, Bhadrawár, Jamú, Naooshera, Púñch, and the valley of the Kishenganga, by Capt. E. Bates, B.S.C. Calcutta, 1873.
32. Handbook to Kashmir, by Dr. Ince. Calcutta, 1868.
33. Recollections of India, by the Hon. C. H. Hardinge. London, 1847.
34. Vaseeri Rupi, the silver country of the Vaseers in Kulu, by J. Calvert. London, 1873.
35. The Himálayan districts of Kooloo, Lahoul, and Spiti, by Capt. A. F. Harcourt. Lahore, 1874.
36. Hunting in the Himálaya, by R. H. Dunlop. London, 1860.
37. Large game shooting, Thibet and the North-West, by A. Kinloch. London, 1876.
38. The large and small game of Bengal and the North-Western Provinces of India, by S. J. Baldwin. London, 1876.
39. From the Hebrides to the Himálayas, by C. F. G. Cumming. London, 1876.

40. A grammar and vocabulary of the Kashmiri language, by M. P. Edgworth. J. A. S., Ben., X., 1038
41. A grammar of the Kashmiri language, by R. Leach. *Ibid*, XIII., 397, 553.
42. Kashmiri vocabulary and grammatical forms, by L. Bowring. *Ibid*, XXXV., ii., 225.
43. A vocabulary of English, Balti and Kashmiri, by Capt. H. H. Godwin Austen. *Ibid*, XXXV., i., 228.
44. A vocabulary of the Kashmiri language, by Dr. W. J. Elmslie. London, 1872.
45. Kashmiri test-words, by the same. J. A. S., Ben., XXXIX., 95.
46. A vocabulary of the Koonawar language, by A. Gerard. *Ibid*, XI., 479.
47. Manners and custom of the Dards, by G. W. Leitner. Ind. Ant., I., 7, 84, 187.
48. The languages and races of Dardistan, by the same. Lahore, 1877.

Section F.—Afghanistan.

(a)—BALUCHISTAN.

1. Notes taken on a tour through part of Baluchistan in 1836-39 by Haji Abdun Nabi of Kábul, edited by R. Leech. J. A. S., Ben., XIII., 667, 786.
2. Some account of a journey from Kurrachee to Hinglaj in the Lus territory, descriptive of the intermediate country and of the port of Sonmeanee, by N. Hart. *Ibid*, IX., 134.
3. Map of the route to Hinglaj, by the same. *Ibid*, IX., 615.
4. Journeys in Balochistan, Afghanistan, the Panjáb, and Kalat, by C. Masson. London, 1844.
5. Caravan journeys and wanderings in Persia, Afghanistan, Turkistan, and Baluchistan, by J. P. Ferrier. London, 1856.
6. Travels in Belochistan and Scinde, by H. Pottinger. London, 1816.
7. Central Asia, Part III., Balochistan, by Col. MacGregor. Calcutta, 1875.
8. The country of Balochistan, by A. W. Hughes. London, 1877.
9. Gazetteer of Sind, by the same. London, 1874.
10. Report on the country between Kurrachee, Tatta, and Sehwan Scinde, by E. P. de La Hoste. J. A. S., Ben., IX., 907.
11. Brief history of Kalat, brought down to the deposition and death of Mehrab Khan Brahosee, by R. Leech. *Ibid*, XII., 473.
12. Memorandum on the Beloch tribes in the Dera Gházi Khán district, by Capt. Minchin. Lahore, 1869.
13. Sindh and the races that inhabit the Indus valley, by R. Burton. London, 1851. Correspondence relative to Sind, 1836-43, presented to both houses of Parliament by Her Majesty's command. Calcutta, 1844.
14. Note on the Brahosees, by N. Hart. *Ibid*, X., 186.
15. General notice of the tribes of Kujjuksyes, by the same. *Ibid*, IX., 1214.
16. Report on Upper Sindh and the eastern portion of Cutchee, with a memorandum on the Beloochee and other tribes of Upper Sindh and Cutchee, by J. Postana. *Ibid*, XII., 123.
17. Vocabulary of the Scind language, by J. B. Eastwick. Bombay, 1843, and J. A. S., Ben., XII., 1.

18. Sindhi and English dictionary, by G. Stack. Bombay, 1855.
19. Sindhi reading-book, by Dr. E. Trumpp. London, 1858.
20. Grammar of the Sindhi language, by the same. London, 1872.
21. Grammar of the Sindhi language, by W. H. Wathen. J. A. S., Ben., VI., 247.
22. Epitome of the grammars of the Brahniky, the Balochky, and the Panjábi languages, with vocabularies of the Baraky, the Pashi, the Laghmani, the Cashgari, the Teerhol, and the Deer dialects, by Lieut. R. Leech. *Ibid*, VII., 588, 608, 711, 780.
23. A Balochee grammar, by E. Mockler. London, 1877.

(b)—WESTERN FRONTIER.

24. Narrative of a journey into Khorasan and the countries north-east of Persia in 1821-22, by J. B. Fraser. London, 1825.
25. Extracts from a demi-official report on a mission into Khorasan, by A. Conolly. J. A. S., Ben., X., 116, 718.
26. Ibn Haukul's account of Khorasan, by W. Anderson. *Ibid*, XXII., 162.
27. A description of the country of Seistan, by R. Leech. *Ibid*, XIII., 115.
28. Sketch of the physical geography of Seistan, by A. Conolly. *Ibid*, IX., 710.
29. Journal kept whilst travelling in Seistan, by the same. *Ibid*, X., 319.
30. Narrative of a journey from Herat to Khiva, by Capt. J. Abbot. London, 1843.
31. Central Asia, Part IV., Persia, by Col. MacGregor. Calcutta, 1871.
32. Narrative of a journey through the Province of Khorasan and on the north-western frontier of Afghanistan in 1875, by Col. C. M. MacGregor. London, 1879.
33. From the Indus to the Tigris: a narrative of a journey through the countries of Baluchistan, Afghanistan, Khorasan, and Iran in 1872: together with a synoptical grammar and vocabulary of the Brahoe language, by H. W. Bellew. London, 1874.
34. Clouds in the east: travels and adventures on the Perso-Turkoman frontier, by V. Baker. London, 1876.
35. Journey on the Turkoman frontier of Persia, by G. Napier. P. R. G. S., XX., 8.
36. Eastern Persia: an account of the journeys of the Persian boundary commission, 1872. London, 1876.
37. The sixth great oriental monarchy, or the geography, history, and antiquities of Parthia, by Professor Rawlinson. London, 1876.
38. The seventh great oriental monarchy, or a history of the Sassanians or new Persian Empire, with notices, geographical and antiquarian, by the same. London, 1876.
39. Persia, from the earliest period to the Arab conquest, by W. S. W. Vaux. London, 1876.

(c)—AFGHANISTAN.

40. Journey from Bengal to England through Kashmir, Afghanistan, and Persia into Russia, by G. Forster. London, 1796.
41. Account of the kingdom of Cabul, comprising a history of the Afghan nation, by M. Elphinstone. London, 1818.
42. Account of the kingdom of Cabul and its dependencies in Persia, Tartary, and India: a view of the Afghan nation and history of the Dooraunee monarchy, by the same. London, 1842.

43. Narrative of a visit to Ghazni, Kabul, and Afghanistan, by G. Vigne. London, 1840.
44. Journey to and residence in Cabul, by A. Barnes. London, 1843.
45. Journey to the north of India overland from England, by A. Conolly. London, 1834.
46. *Ariana Antiqua*: a descriptive account of the antiquities and coins of Afghanistan, with a memoir on the buildings called *topes*, by C. Masson, by H. H. Wilson. London, 1841.
47. Ancient geography of India: Buddhist period, by A. Cunningham. London, 1871.
48. *Memoirs of Zehir-ed-in Muhammad Baber*, written by himself in Jaghatai Turki, translated from the Persian version by Leyden and Erakine. London, 1826.
49. Geographical notice of the valley of Jalalabad, by G. H. MacGregor. *J. A. S. Ben.*, XI., 117.
50. Notice of Jalalabad, by J. G. Gerard. *Ibid.*, III., 325.
51. Journal of a political mission to Afghanistan in 1837, by Dr. Bellew. London, 1862.
52. On the highland region adjacent to the trans-Indus frontier of India, by Col. J. T. Walker. *R. G. S. J.*, XXXII., 816.
53. *Central Asia*, Part II., Afghanistan, by Col. MacGregor. Calcutta, 1875.
54. Survey operations of the Afghanistan expedition, the Kurram valley, by Capt. G. Martin. *P. B. G. S.*, I., 617, N. S.
55. Afghan geography, by C. R. Markham. *P. B. G. S.*, XX., 4, 1876.
56. Mountain passes on the Afghan frontier, by the same. *Ibid.*, I., 38, N. S.
57. The upper basin of the Cabul river, by the same. *Ibid.*, 110.
58. The basin of the Helmund, by the same. *Ibid.*, 191.
59. The mountain passes leading to the valley of Bamian, by Lieut.-Col. Kays. *Ibid.*, 244.
60. March and operations of the army of the Indus to Afghanistan in 1838-39, by Major W. Hough. London, 1841.
61. British operations at Kábul in 1841-42, by the same. London, 1849.
62. Campaign in Sind and Afghanistan, by Col. Outram. London, 1840.
63. Narrative of the campaign of the army of the Indus, in Scind, and Cabul in 1838-39, by R. H. Kennedy. London, 1840.
64. Diary of a march through Sind and Afghanistan, by Rev. J. N. Allen. London, 1843.
65. Sale's brigade in Afghanistan, by Rev. G. R. Gleig. London, 1846.
66. The Sikhs and Afghans immediately before and after the death of Ranjit Singh, from the journal of an expedition to Cabul through the Panjáb and the Khaiber Pass, by Shahamat Ali. Calcutta, 1847.
67. History of the war in Afghanistan, by J. W. Kays. London, 1851.
68. Reminiscences of forty-three years in India, by Lieut.-Genl. Sir G. Lawrence. London 1874.
69. A retrospect of the Afghan war, with reference to passing events in Central Asia, by Sir V. Eyre. London, 1869.
70. The Panjáb and North-Western frontier of India, by Major-Genl. H. Coxe. London, 1878.

71. General Gilbert's raid to the Khaibar, by R. W. Bingham. London, 1850.
72. History of Afghanistan from the earlier period to the outbreak of the war in 1878, by Col. Malleson. London, 1879.
73. Journal of a tour through part of the Panjáb and Afghanistan in the year 1837, by Agha Abbass of Shiraz, edited by R. Leech. J. A. S., Ben., XII., 564.
74. Account of parts of the Cabul and Peshawar territories, by the same. *Ibid*, XIV., 660.
75. Reports by Burnes, Lord, and Wood, political, geographical, and commercial, on Swinde and Afghanistan. Calcutta, 1839.
76. Memoir on the climate, soil, produce, and husbandry of Afghanistan, by Lieut. Irwin. J. A. S. Ben., VIII., 745, 779, 869, 1005: IX., 33. 189.
77. Extracts from a report on subjects connected with Afghanistan, by W. Griffiths. *Ibid*, X., 977.
78. Notes on the geography of Western Afghanistan, by W. Anderson. *Ibid*, XVIII., i., 553.
79. Verification of the Itinerary of Hwen Thsang through Afghanistan and India, by Capt. A. Cunningham. *Ibid*, XVII., i., 476, and ii., 13.
80. Route from Dera Gházi Khán through the Veziri country to Kábul. by M. Honigberger. *Ibid*, III., 176.
81. Route from Dera Gházi Khán to Kandahar through the Sakhi Sarwar and Buzdar passes, with other routes, by R. Leech. *Ibid*, XIII., 527.
82. Diary of a trip to Ind Dadan Khán, &c., by A. Fleming. *Ibid*, XVI., i., 664.
83. Account of Ghazni. *Ibid*, VI., 775.
84. Itinerary from Yezd to Herat (official). XIII., 827.
85. Route from Kandahar to Herat (official). XIII., 121.
86. Report of a journey by Candahar, Kábul, and the Panjáb from Herat to Simla, by Major Todd. XIII., 339.
87. Description of Herat, by Mohan Lal, III., 9.
88. Central Asia, Part I., our north-western frontier, by Col. MacGregor. Calcutta, 1871.
89. Banna, or our Afghán frontier, by S. Thorburn. London, 1876.
90. Notes on the Bunnoo district, by Major H. B. Urmston. Lahore, 1869.

(d)—LANGUAGES.

91. Dictionary of the Panjábí language, prepared by a Committee of the Lodiana Mission. Lodiana, 1854.
92. Pukhto-English dictionary, by H. G. Raverty. London, 1860.
93. Grammar of the Pukhto or Pushto language, by the same. London, 1869.
94. Grammar of the Pashto, or language of the Afgháns, by Dr. E. Trumpp. London.
95. Dictionary of the Pukkhto or Pukhto language, by H. W. Bellew. London, 1867.
96. Grammar of the Pukkhto language, by the same.
97. Grammar of the Pushtoo or Afgháni language, by R. Leech. J. A. S., Ben., VIII., i.: XXIII., 560.
98. Is the Pushto a Semitic language? by Rev. I. Loewenthal. *Ibid*, XXIX., 323.
99. Some remarks on the origin of the Afghan people and dialect, by H. G. Raverty. *Ibid*, XXIII., 550.

100. Translation of the Pukhto text-book, Kalid-i-Afgháni, by T. C. Plowden. Lahore, 1875.
101. Notes on the Eusofzye tribes of Afghanistan, by E. Conolly. *Ibid*, IX., 924 : XVII., i., 624.
102. Report on the Yoosoozaie district, by Lieut. H. R. Lumsden. *Sel. Rec. Panj.*, I, 5, 1853. And Covert, J. A. S., Ben., V., 481.
103. An account of the early Ghiljaees, by R. Leech. *Ibid*, XIV., 304.
104. An account of the early Abdalees, by the same. *Ibid*, XIV., 445.
105. An account of the Hazarehs, by the same. *Ibid*, XIV., 393.
106. Account of Roh in Derajat, by H. G. Raverty. *Ibid*, XXVI., 177.
107. The district of Dera Ismáíl Khán, by T. W. Tolbort. *Ibid*, XI., 1.
108. History of the Gakkhars, by J. G. Delmerick. *Ibid*, XI., 67.
109. Rough notes on the Khutuks of the Terce country and Khurah below Atok, by Lieut. F. H. Pollock. Lahore, 1850.
110. Report on the eastern Khutuks, by Lieut. H. Lumsden. *Ibid*, 1851.
111. The lower Derajat frontier, by Capt. G. O. Jacob. *Ibid*, 1854.
112. Kohistan of the Jullundhur Doab, by Lieut. Parish. J. A. S., Ben., XVII., i., 281 : XVIII., i., 360.
113. The Synds of Tira and their quarrels with the Sunis, by Capt. P. Cavagnari of Kohat. Lahore, 1875.
114. Notes on the Dera Ghazi Khán district and its border tribes, by R. Bruce. Lahore, 1871.
115. Memorandum on the Dera Gházi Khán district. *Sel. Rec. Panj.*, IV., 4, 1860.

(c)—KOHISTÁN.

116. An account of upper and lower Suwat and the Kohistán to the source of the Suwat river and of the tribes inhabiting those valleys, by Capt. H. G. Raverty. J. A. S., Ben., XXXI., 227.
117. An account of upper Káshkár and Chitrál or lower Káshkár, together with the independent Afghán State of Panjkorah, including Talásh, by the same. *Ibid*, XXXIII., 125.
118. An account of the Panjkora valley and upper and lower Káshkára, by Rajah Khán of Cabul, edited by R. Leech. *Ibid*, XIV., 812.
119. Some account of a visit to the plain of Koh-i-Daman, the district of Ghorband, and the pass of Hindu-kush, by P. B. Lord. *Ibid*, VII., 521.
120. On the Siah-posh Kaffirs, with specimens of their language and costume, by Capt. A. Burnes. J. A. S., Ben., VII., 325.
121. Information regarding the Siah-posh tribe, by A. Burnes. *Ibid*, II., 305.
122. Further information regarding the same, by Mohan Lal. *Ibid*, III., 76.
123. On the language of the Siah-posh Kaffirs, with a short list of words, to which are added specimens of Kohistáni and other dialects spoken on the northern borders of Afghanistan, by H. G. Raverty. *Ibid*, XXIII., 267.
124. Notes on Kafiristan, by the same. *Ibid*, XXVIII., 817.
125. On the language of the so-called Kaffirs of the Indian Caucasus, by Dr. Trumpp. J. R. A. Soc., XIX.

Section G.—Pamir and Turkistan.

1. Memorandum on routes from the Panjáb to eastern Turkistán, by T. D. Forsyth. Lahore, 1868.
2. Report on the route to the Karakash valley and pass, by Dr. H. Caley. *Ibid.*
3. Results of a scientific mission to India and High Asia, by H. A. and R. vom Schlagentweit. Leipzig, 1861-66.
4. Route-book of the western parts of the Himálaya, Tibet and Central Asia, by the same. Leipzig, 1863.
5. Lahore to Yarkand, by G. Henderson and A. Hume. London, 1872.
6. Visits to High Tartary, Yarkand and Kashgarh, by R. Shaw. London, 1871.
7. The Roof of the world, being the narrative of a journey over the high plateau of Tibet to the Russian frontier and the Oxus sources on Pámír, by Lieut.-Col. T. E. Gordon, C.S.I. Edinburgh, 1876.
8. Kashmir and Kashgarh, a narrative of the journey of the embassy to Kashgarh in 1873-74, by H. W. Bellew, C.S.I. London, 1875.
9. Report of the mission to Yarkand in 1873 under command of Sir T. D. Forsyth. Calcutta, 1875.
10. Account of the survey operations in E. Turkistan, 1873-74, by Capt. H. Trotter. Calcutta, 1875.
11. A sketch of the Turki language as spoken in E. Turkistán, by R. B. Shaw. Lahore, 1875.
12. Les confins Anglo-Russes dans l'Asie Centrale: Etude historique, géographique, politique, et militaire sur la Kachgarie (E. Turkistán), by Capt. Kouropatkine. Translated by G. LeMarchand. Paris, 1849.
13. Benoit de Goës, missionnaire voyageur, 1603-07 (in E. Turkistan), by J. Brucker. Lyons, 1879.
14. Route from Peshawar to Bokhara, by Dr. Gerard and Lieut. A. Burnes. J. A. S., Ben., I., 189: II., i., 143, 224.
15. Memoir on the Usbek State of Kokan, called Khokend (the ancient Ferghana), by W. H. Wathen. *Ibid.*, III., 369.
16. Memoir on Chinese Tartary and Khoten, by the same. *Ibid.*, IV., 653.
17. Notes on the manners and habits of the Turkoman tribes, with some geographical notices, by E. Stirling. *Ibid.*, X., 290.
18. Russian geographical operations in Asia, by Col. J. T. Walker. *Ibid.*, XXXV., 77.
19. Note on Central Asia. *Ibid.*, XXXIV., 113.
20. Central Asia, by T. Hutton. London, 1874.
21. Central Asia and the Anglo-Russian frontier, by A. Vambéry. London, 1874.
22. History of Bokhara, by the same. London, 1873.
23. Asie Centrale, recherches, by A. von Humboldt. Paris, 1843.
24. Travels into Bokhara, journey from India to Kábul, Tartary, and Persia, and voyage on the Indus, by A. Burnes. London, 1834.
25. Diplomatic transactions in Central Asia, 1834-38, by J. Urquhart. London, 1841.
26. Journey to the source of the Oxus by the route of the Indus, Kábul and Badakshan, by J. Wood. London, 1841. New edition, London, 1872.
27. Narrative of a mission to Bokhara in the years 1843-45 to ascertain the fate of Col. Stockart and Capt. Conolly, by Dr. J. Wolff. London, 1846.

28. *Mémoire sur la partie méridionale de l'Asie Centrale*, by N. de Khanik of Paris, 1861.
29. *Notes of a journey in Russian Turkistan, Khokand, Bokhara and Kuldja*, by E. Schuyler. London, 1876.
30. *Campaigning on the Oxus and the fall of Khiva*, by J. A. MacGahan. London, 1876.

Section H.—Tibet.

1. *Geographical notice of Tibet*, by Csoma de Kőrös. J. A. S., Ben. I., 121.
2. *The physical geography of Western Tibet*, by Capt. H. Strachey. London, 1851, and J. R. G. S. XXIII., 2.
3. *Thibet nach den resultaten geographischer forschungen früherer und neuester zeit*, by K. Ganzenmüller. Stuttgart, 1877.
4. *Tibet, Tartary and Mongolia*, by H. T. Prinsep. London, 1851.
5. *Explorations in Western Tibet by the Trans-Himalayan parties of the G. T. S.* P. R. G. S., 444, 1879.
6. *Historie générale des Huns, des Turcs, des Mongols et des autres Tartares occidentaux*, by Des Guignes. Paris, 1756-58.
7. *History of the Mongols*, by H. H. Howorth. London, 1876.
8. *History of the Genghizan the Great, first Emperor of the ancient Moguls and Tartars*, by Petis de la Croix, translated into English. Calcutta, 1819.
9. *The book of Ser Marco Polo, concerning the kingdoms and marvels of the east*, edited by Col. Yule. London, 1875.
10. *Cathay and the way thither*, by the same. London.
11. *Recueil de Turkestan: comprenant des livres et des articles sur l'Asie Centrale en général et la province du Turkistan en particulier*, by V. J. Mejev. St. Petersburg, 1878.
12. *Notices of the mediæval geography of Central and Western Asia*, by E. Bretschneider. London.
13. *Notes on Chinese mediæval travellers to the west*, by the same.
14. *Itinéraire de Marco Polo à travers la région du Pamir au XIII^e siècle*, by J. B. Paquier. Paris, 1876.
15. *Le Pamir : étude de géographie physique et historique sur l'Asie Centrale*, by the same. Paris, 1876.
16. *The Russians in Central Asia, their relations with Khiva, Bokhara and Kokand, descriptions of Chinese Turkistan and Dzungaria from the Russian of Valikhanoff, &c.*, by Messrs. Michell. London, 1865.
17. *Le Kohistan, le Ferghanah et Kouldja, avec un appendice sur la Kachgarie*, by C. E. de Ujfalvy de Meso-Kővesd. Paris, 1876.
18. *Le Syr-Daria, Le Zératchéne, les pays des sept-rivières et la Sibirie-occidentale*, by the same. Paris, 1879.
19. *Résultats anthropologiques d'un voyage en Asie Centrale, communiqués au Congrès anthropologique de Moscou*, by the same. Paris, 1880.
20. *Die Russen in Central Asien*, by Fr. von Hellwald. Augsburg, 1877.
21. *Pévtsoff's expedition in North-Western Mongolia*, by E. D. Morgan. P. R. G. S., 701, 1879.

22. *The Russians on the Amur*, by E. G. Ravenstein. London, 1861.
 23. *England and Russia in the East : a series of papers on the political and geographical condition of Central Asia*, by Major-General Sir Henry Rawlinson. London, 1875.
 24. *Mongolia, the Tangut country and the solitudes of Northern Tibet ; being a narrative of three years' travel in eastern High Asia*, by Lieut.-Col. N. Przhevalsky. London, 1876.
 25. *From Kulja across the Tian-Shán to Lob Nor*, by the same. London, 1879.
 26. *The river of Golden Sand, the narrative of a journey through China and eastern Tibet to Burmah*, by Captain W. Gill, R. E. ; with an introductory essay by Colonel H. Yule, R. E., London, 1880.
-

CHAPTER II.

PHYSICAL GEOGRAPHY OF THE HIMÁLAYA.

CONTENTS.

The Himálavan river basins—The Indus basin—Subordinate systems of the Indus basin ; The Ganges basin—Subordinate systems of the Ganges basin ; The Brahmaputra basin—Subordinate systems of the Brahmaputra basin ; Systems of Tibet—Lob-Nor basin ; Oxus basin. Plains of Hindustán—Indus plain ; Indian desert ; Gangetic plain ; Anám valley ; Bhábar ; Taráí ; Siwálíks ; Dúna. Extent of Bhábar—Cause of the deposit—Hodgson's oceanic theory ; Pluvial theory ; Taráí ; Siwálíks ; Dúna ; Ganges to Brahmaputra ; Lower Himálaya. Materials of the mountains—Eocene period ; Mode of mountain formation ; Mountain-sculpture ; Ice-wedges ; Avalanches ; Glaciers ; Ráin ; Rivers.

WE have already decided to restrict our use of the word 'Himálaya' to that portion of the great mountain girdle which lies between the bend of the Indus on the west and the unexplored gorge of the Brahmaputra on the east. Hodgson, in 1849, estimated the length of this range at 1,800 miles with a mean breadth of about 90 miles, a maximum breadth of 110 miles, and a minimum breadth of about 70 miles. In fixing the breadth of the Himálaya, however, as in determining its length, we have similar difficulties to contend with. For, as we have seen that the popular estimate as to the boundaries of the range may possibly be correctly extended both on the east and on the west, so on the north, the channels of the Indus and the Brahmaputra, which are commonly assigned as the northern limit, are found in a plateau, but little lower than the passes by which the traveller crosses the first line of snowy mountains into Tibet. On the south, we have a well-marked descent to the plains of India, but on the north there is no immediate descent to a lower country beyond. On the contrary, range after range is met with to the north, many of which may compete in altitude with the snowy mountains seen from the plains of northern India.

Before proceeding to a closer examination of the form of that portion of the Himálaya lying within the province of Kumaon, it will be convenient briefly to describe the river-basins throughout the Himálaya and

River-basins.

endeavour to trace the general law underlying their arrangement. For as this depends in the main on the direction of the great ranges and the position of the great peaks, a consideration of it will conduce to a clearer apprehension of the entire mountain system itself as well as of the relations of its various parts. We find that from a water-parting about longitude 81° east and almost immediately due north of Kuinaon, the drainage of the southern part of the Tibetan plateau flows north-west in the Indus and south-east in the Brahmaputra. These two rivers maintain a course along the length of the table-land, and receive as they proceed the drainage of a large part of its breadth; the exceptions being, first, the eastern border, which apparently is drained by the Lu-tse, the Lan-tsang, and the Murui-ussu, one of the sources of the Yang-tse-kiang;¹ second, an occasional strip along the southern edge from which the water passes off² more or less directly to the south through the Himalaya; and third, the north-western part from which the water has no escape, but is collected in lakes at the lowest level it can reach.³ The waters thus accumulated in the two great streams are at length discharged by two openings in the Himalayan slope through the plains of India into the Indian Ocean. No great portion of the drainage of the table-land, so far as we know, passes in the opposite direction through the northern slope, and the area that discharges itself southward at points intermediate between the debouches of the Indus and the Brahmaputra is with one exception, that of the Satlaj, comparatively small. The waters of the northern slope with a small area of the table-land adjoining flow down to the plains of eastern Turkistán: while, in like manner, those of the southern slope with the drainage of the exceptional area along the southern border of the table-land which passes through the line of water-parting from the north, give rise to such rivers as the Jhilam, Chináb, Ravi, Jumna,

¹ From Prejevalsky's Mongolia (London, 1876) it would appear that his Burkhan Buddha range marks, in 96° longitude, the north-eastern termination of the Tibetan plateau. This range forms the southern boundary of the Tsaidam plain, which, according to native report, extends thence to Lob-Nor. The Nomokhun-gol, which rises on the southern face of the Burkhan Buddha range and joins the Balaingol, does not appear to be a feeder of the Hoang-ho or Yellow River, which has its origin outside the Tibetan plateau. To the south of the Burkhan Buddha range the elevation is from 13,000 to 15,000 as far as the Murui-ussu, the bed of which, where seen by Prejevalsky, was 13,000 feet above the level of sea with a channel 750 feet broad in January. The name 'Murui-ussu' signifies 'the tortuous river,' and, according to Yule, it is the Bri-chu of the Tibetans, the Brius of Marco Polo, and the Yang-tse-kiang or Blue River of the Chinese.

² By the Satlaj, Karnali, and Arun. ³ Examples of river-systems without an outlet to the sea are to be found in the basins of the Caspian, Aral, Balkhash, Lob-Nor, &c.

Ganges, Káli, Gandak, Kosi, and Tista. We thus see that the northern crest of the table-land or the summit of its northern slope practically forms the water-parting between the rivers that flow southwards and those that lose themselves in the plain of Gobi. In the southern crest we have a subordinate water-parting separating the rivers that fall into the Indian Ocean into two classes; first, those that rise on that slope and flow directly down it to the plains of Hindustán; and secondly, those that are collected along the table-land and are finally discharged also through the southern slope, chiefly by two concentrated channels at distant points towards the ends of the range. Captain Henry Strachey¹ has called the northern crest of the table-land the Turkish, and the southern the Indian water-parting.

If we examine the river-systems having their source in the Himálaya, we find a regularity of plan and arrangement which at first sight would not be expected. Taking the Indus basin and its system, we see that the Satlaj and all the rivers that join the Indus on its left bank have a south-westerly direction towards the Arabian Sea. On the east this system is bounded by the small inland basin of the Kaggar, which receives the drainage from an inconsiderable portion of the outer hills between the Satlaj and the Jumna, and finally loses its waters in the Indian desert. The eastern water-parting of the Indus system is found in the elevated range extending from the main Himálayan mass along the left bank of the Satlaj to Rúpur, and is continued thence in the uplands bordering the *khádír* of that river until it meets the Arvali (Aravali) range which constitutes the north-western abutment of the table-land of Central India. The character of the slope towards the south-west will be best understood from the following table of heights taken along the course of the Satlaj to Ludhiána, and thence by the Grand Trunk Road to the Jumna:—Táru, about two miles below the junction of the Panjnad and Indus, 337 feet above the level of the sea; Baháwalpur, 375 feet; Núr Sháh, 481 feet; Pír Khális, seven miles north-east of Baháwalgarh, 548 feet; Fazilka, 588 feet; Firozpur cantonment, 645 feet; Jagraon, 765 feet; Ludhiána, 806 feet; Amballa Church, 899 feet; and Madalpur, on the banks of the Jumna *khádír*, 906

¹ J. R. G. S., XXIII., p. 7.

feet. From Sodiwála in the Firozpur district southwards on the 75th meridian we have Sodiwála, 718 feet; Ahmadwála, 705 feet; Nirsá, on a mound, 737 feet; level of Sambhar lake, 1,184 feet.¹ These observations show a slight depression towards the south in addition to that towards the south-west, and would make us include the Kaggar inland basin in the Indus system. The Sotra or Hakra, the ancient river of the Indian desert, seems, however, to have once had its debouche in the Ran of Kachh,² and would therefore be still entitled to be considered separate. To the west the Rávi and the Chináb run in lines almost parallel to the Satlaj, as well as the Jhilam, from the town of that name, until it takes a bend to the south to its junction with the Chináb. To the west, the water-parting of the Indus commences at Cape Monze on the Arabian Sea, and advances nearly northwards along the Hala mountains to the east of Sohráb, Kalát, and Quetta (Kwatab). It thence continues in the same direction along the western Sulaimán range, also known as the Konak, and Kúb Jadrán range to the Safed-koh, at the head of the Kurran valley. Thence it follows the crest of the Safedkoh westwards to the hills north of Ghazni, where it separates the southern affluents of the Kábul river from the waters of the small inland basin of lake Abistida. The direction is then continued north-westerly to the ridge separating the head-waters of the Argand-áb from those of the Kábul river, and again in the range that separates the waters of the Halmand basin from the most westerly affluents of the Kábul river, whence a transverse ridge near the pass to Bámian connects the line of water-parting with the Hindu-kush. For 300 miles the line follows the Hindu-kush to its junction with the great Tighdambásh Pámr near the Baroghil pass. It then follows the Muztágh range,³ but cuts through it around by the Kárákorum pass to the north, so as to include the tributaries of the Shayok, and proceeds in a south-easterly direction by the Aling Gang-ri to its junction with the Gang-ri at Kailás, where a transverse ridge separates the head-waters of the Indus, the Brahmaputra, and the trans-Himálayan feeder of the Ganges system. The Indus has a length of 1,800 miles, and, according to Mr. Saunders, its basin has an area of 372,000 square miles.

¹ G. T. S. tables.
Cal. Rev., July, 1874.

² See Notes on the lost river of the Indian desert,
³ 'Ice-mountain': a better name than Kárákorum
(black-gravel), which should be restricted to the pass.

The Hala mountains to the south of the Mula pass are better known as the Khirthar hills and as the Pubh hills, and the drainage from them loses itself in the plains before it can reach the Indus. The same may be said of the drainage through the Mula and Bolan passes, and that by the Thal valley. To the north, the Luni, Gomai, Kurram, and Kábul rivers, each possessing a perennial stream, have an easterly course more or less parallel to each other, and break through the range bordering the right bank of the Indus by deep and narrow gorges which form the passes into the upper country. The Kábul river itself is the drainage channel for the very elevated country from the hills north of Ghazni to the Baroghil pass north of Chitrál, and from Naushera to the pass leading to Bámián, and thus forms a compact subordinate system well deserving of separate study.¹ The upper waters of the Jhilam drain the Tile and Kashmir valleys, and have a general direction between west and north to Muzaffarabad, where they unite, and, meeting a meridional ridge, take a bend southwards to the plains. The Chínáb, in the upper portion of its course known as the Chandra, has a similar direction between west and north until it meets the spurs from the range which forms the water-parting between it and the Jhilam, whence it seeks an outlet southwards towards the plains. The Ravi runs in a valley parallel to that of the Chínáb and south of it until it meets the outliers of the range that forms the water-parting between it and the Chínáb, when it also turns suddenly southwards towards the plains. Next comes the Biás, which has also a westerly direction until it meets the ridge between it and the Ravi, when it takes a bend to the south through the outer hills. The upper course of the Satlaj has a similar westerly trend until it meets the great obstruction culminating in the Leo Porgyul peak, after which the direction is between west and south until it enters the plains. A range runs between the Satlaj and the southern branch of the Indus from the meridian of Tirthapuri by Gár, to where it is joined by the ridge connecting it with Leo Porgyul, and thence into Rupshu to the north of the Tso Moriri lake, and constitutes the subordinate water-parting between the Satlaj itself and the

¹ For interesting articles on the tract between the Arabian sea and the Gilgit river, by Mr. C. B. Markham, see *Proceedings R. G. S.*, 1879 : the mountain passes on the Afghan frontier of British India, p. 38 : the upper basin of the Kábul river p. 110 : the basin of the Helmund. p. 181.

Indus. Thus we see that the great feeders of the Indus system from the west have a similar character, and that those from the east, from the Himálaya proper, have a general westerly direction, in the upper portions of their courses in the hills, until they turn southwards towards the plains, where the direction is south-west to their junction with the Indus.

Next we have the Ganges basin with its subordinate systems.

The Ganges basin.

To the north, the water-parting, as a rule, follows the ghát-line of the Himálaya and on the extreme west separates the sources of the Jádth-Ganga, one of the head-waters of the Bhágirathi, from the Hop-gadh, an affluent of the Satlaj. North of Kumaon, however, we have a phenomenon similar to that observed near the Kárákorum¹ pass, where the waters of the Shayok, Yárkand, and Kárákash rivers have a common origin in that elevated plateau at no great distance from each other. We find that to the east of the Unta-dhúra pass, north of Milam in Kumaon, the water-parting of the Ganges basin crosses to the north of the ghát-line to a place called Tara, where the sources of the Satlaj and the Karnáli lie close together, "divided by an almost level plain, across which a man might walk from one river to the other in an hour or two, without a vertical ascent or descent of 500 feet," yet the waters of one stream seek the sea at Karáchi, and of the other by Goalundo. Further east the water-parting is continued in the ghát-line to the Arun river, which has its sources to the north and forces for itself a way through the Himálaya to the plains. The water-parting then follows the eastern boundary of Nepál to the plains, where an intricate system of drainage is met with, throwing off feeders sometimes to the Brahmaputra and sometimes to the Ganges down to their junction at Goalundo. Following the line on the west from east longitude 79° 11' and north latitude 25°, we

¹ Dr. Scully describes the appearance of the country between the top of one of the Shayok gorges and the Kárákorum pass thus:—"At the top of the ascent, a strange sight met our eyes, for we found ourselves on an immense undulating plain, the Dipeang, which looked like the top of the world." * * Northwards, in front of us, we saw a few irregular flat-topped hillocks, they looked like scattered about * * I had occasion to look back in the direction of the route by which we had come. A fine snowy range of mountains met my view and looked quite continuous; but, of course, this was a deceptive appearance, as we had passed through this chain without crossing any pass." This was the Mustágh range, whilst the Kárákorum pass, distant about 25 miles ahead and forming the water-parting, lay amongst the seeming hillocks, a fact which shows that though a range may form a water-parting, a water-parting does not always form a range. Colonel Gordon gives views of the Kárákorum pass in his 'Roof of the World.'"

find in the extreme north-west that the water-parting keeps to the crest of the range running along the left bank of the Baspa, an affluent of the Satlaj, and is continued by Hattu to the ridge on which Simla is built. Thence it proceeds southward, and then eastward along the right bank of the Giri to the junction of that stream with the Jumna near Rájghát. Here the water-parting turns southwards along the line separating the drainage area of the Kaggar system from that of the Jumna, and continuing along the Arvali range and the edge of the Málwa plateau, passes through the Jabulpur and Mandla districts, separating the sources of the Nerbudda from those of the Son, and then along the range connecting the Sátপুরas with the Rájmahál hills to the plains, where it follows a course along the left bank of the Sabanreka to the sea. Mr. Saunders has given the length of the main stream of the Ganges as 1,514 miles, and the area of its basin at 391,000 square miles.

The Ganges basin, like that of the Indus, possesses several subordinate systems that may be called in their order from west to east, the Jumna-Ganges, Karnáli, Gandak, and Kosi systems. The alpine basin of the Jumna-Ganges system is bounded on the west by the well-defined range which descends from the Jamnotri group of peaks to the Satlaj river, and on the east by a similar ridge descending from the Nanda Devi group of peaks and separating the waters of the Pindar from those of the Himálayan Sarju. To the north, the ghát-line separates it from the source of the Karnáli on the east, and the source of the Satlaj on the west. Although the upper waters of both the Jumna and the Alaknanda, or principal source of the Ganges, have at first a westerly direction, they soon take a bend to the south, and form the main channels to which are directed their affluents from either side. It is remarkable that, with the exception of the Rámghanga, which unites with the Ganges in the Farukhabad district, neither the Jumna nor the Ganges before their junction receives any considerable affluent of Himálayan origin during its course through the plains. The western boundary of the alpine basin of the Karnáli is marked by the ridge extending from the Nanda Devi group between the Pindar and Sarju rivers, already noticed. The eastern boundary

is formed by a similar ridge descending from the Dhaulagiri group of peaks. To the west the Sarju, eastern Rám-ganga, Gori, and Káli enter the plains in one stream as the Sárda. Then the Swetaganga Karnáli, and Bheri form the Karnáli, and further east we have the Jhingrak or Rápti and its affluents. All unite in the plains to form the Ghogra, which joins the Ganges to the south of the Gházipur district. But this unitising law is better exemplified in the alpine basin of the Gandak, which reaches from the Dhaulagiri group to the Gosáin-thán group of peaks. We have here seven rivers named in order from west to east, the Barigár, Naráyani, Sweti-gandaki, Marsyángdi, Daramdi, Burhiya-Gandaki, and Trisúl-Gandaki. These are called the seven Gandaki by the Nepálese, and unite their waters at Tribeni within the hills to form the Gandak river of the plains. Here we have what Hodgson¹ terms an admirably defined natural division lying between two great groups of peaks. In the same manner as the Karnáli basin is bounded on the west by the spur descending from the Nanda-Devi group of peaks, and on the east by the ridge from the Dhaulagiri group, so the Barigár of the Gandak system does not receive a single streamlet from the westward of the Dhaulagiri ridge, nor does the Trisúl of the same system receive any water from the east of the ridge descending from Gosáin-thán. The alpine basin of the Kosi lies between the Gosáin-thán group and the Kanchinjinga group of peaks, and, like the Gandak system, consists of seven rivers,² known as the seven Kosis. These, named in their order from west to east, are the Milamchi or Indrawati, the Bhotiya-Kosi, Támba-Kosi, Likhu-Kosi, Dúd-Kosi, Arun, and Tamra or Tamor. The Arun has one of its sources to the north of the line of snowy peaks seen from the plains, and the Tamor is also said to have trans-nivean affluents, but all the others rise on the southern slope of the Himálaya, and unite within the hills at Varáha-Kshetra above Náthpur. The subordinate systems of the Ganges basin thus appear to be strongly characterised by a common origin within an area bounded on the north by the ghát-line, and on the west and east by well-marked groups of culminating peaks, whence ridges descend and form the water-parting between successive systems.

¹ See his article on the Ganges basin, *J. A. S. Ben.*, XVIII., 761.
XVII., ii., 648.

² *Ibid.*

The Brahmaputra basin in its full extent has not been explored, but sufficient evidence has been collected by recent travellers to show that from the water-

Brahmaputra basin.

parting between the sources of the Brahmaputra and the Indus, the northern water-parting of the former river continues in a range of lofty peaks on its left bank to the bend towards the south, by which it reaches the plains of India. This range has a direction south-east, and to the west of the 86th meridian is sufficiently distant from the Brahmaputra to allow of such affluents as the Chachu and the Charta rivers. About the 86th meridian, a line of peaks culminating in the Tārgot Ia stretch in a north-easterly direction to the Gyákhárma group of peaks, south-east of the Kyáring-cho or Kyáring lake, one of the sources of the Ná-k-chu-kha. The drainage of the southern slope of the range is sent by the Dumphu-chu into the Kyáring lake, so that the northern water-parting of the Brahmaputra must here approach much closer to the river and run in a south-easterly direction. On the 89th meridian, it descends as low as the 30th parallel in the Shiang Lahu range, which appears to be connected with the great Ninjin-thúngla range of snowy peaks to the south of the Jáng Namcho or Tengri-Nor lake with a trend to the north-east, for it gives the head-waters of the Ki-chu or Lhása river from its southern slope, as well as other important streams further east, regarding which our information is still very imperfect. To the north-east we find the Ná-k-chu-kha or Hota Sanpo, a large river that issues from the Chargut lake about north latitude 32° and east longitude 89°, and flows eastward, having its drainage area on the south, bounded by the water-parting between it and the Brahmaputra basin. This great river takes a bend to the south, and according to one of the Pandit explorers,¹ flows by Tsiamdo on the road from Lhása to Bathang, and thence through Amdu to China, under the names Máchu and Konkong. These names would connect it with the Yang-tee, but if it flows by Tsiamdo it should be one of the branches of the Lan-Tsang, the name of the upper portion of the Mekong or Kam-bodia river. Des Godins notes that the Nu-Tse or Lu-Tse is known as the Ngen-kio in Tibet, a name which may perhaps be referred to

¹ G. T. S. Rep., 1878-79, p. 63. The *Athenæum* of the 17th April, 1880, announces the arrival of Colonel Prejevalsky at Ná-k-chu-kha, about twelve marches north of Lhása, and we shall doubtless obtain some solution of the problem connected with this river from this great traveller.

the Nák-chu, and it would then be the head-waters of the Salween, whilst others claim it as the source of the Irawadi. All that can, therefore, be said is that there is a great river in eastern Tibet, between which and the Brahmaputra is an unexplored tract of country, and within it will be found the eastern water-parting of the Brahmaputra. So far as we may conjecture from the imperfect materials at our command, the range that forms the northern water-parting of the Brahmaputra takes a sudden bend southwards between the 96th and 97th meridians along the right bank of the Lu-Tse. Thence one branch proceeds westwards in the Patkoi range, and another southwards, between the Irawadi and Salween. To the south, the water-parting follows the crest of the Patkoi range, and is continued westwards in the Manipur, Lushái, and Chittagong hills, where it separates the southern affluents of the Brahmaputra from those of the Barmese systems. To the west, the water-parting is conterminous with that of the Ganges basin. Mr. Saunders estimates the course of the Brahmaputra at 1,800 miles, and the area of its basin at 361,000 square miles.

Following the systems that carry off the drainage from the southern slopes of the Himálaya and join the Brahmaputra basin. maputra in its course through the plains, we trace much the same regularity found in the subordinate systems of the Ganges basin further west. The Tista system of Sikkim is bounded on the west by a ridge descending from the Kanchinjunga group of peaks, and on the east by a similar ridge from the Chumalári group that also forms the eastern boundary of Sikkim. The alpine rivers of this system in order from west to east are the Bári Ranjít, Ratong, Lachen, Lachung, and Rang-chu, and all unite within the hills above Kalingpong to the east of Darjiling. The alpine basin of the Tarsa-Gangadhar system extends from the Chumalári group on the west to the ridge descending some fifteen miles east of the 90th meridian in the 28th parallel; and which separates the waters of the affluents of the Tarsa from those of the Manás system. The rivers of this alpine basin from west to east are the Ammo, Dor, Par, Wang, Ma, Pachu, and Tanchu, which unite within the hills to form the Tarsa and the Gangadhar tributaries of the Brahmaputra. Further east comes the Manás system, of which the western water-

parting is conterminous with that of the Tarsa-Gangadhar system. On the east, it is bounded by a ridge descending from the group of snowy peaks to the west of the Karkang pass crossed by the Pandit on the Táwáng route in 1873, and for its alpine feeders has the Mati, Manás, Kuru, Lhopra, and Táwáng streams, which apparently also unite within the verge of the mountains to form the Manás. Eastwards lies the Subansiri system between the Manás and the Dihong. The Sikang-chu, which rises to the south of the Karkang pass, probably forms one of its sources, but the remainder lie within the wild country of the Abors and have not yet been explored. The southern affluents of the Brahmaputra during its course through the valley of Asám are not so important, and may be divided into two classes—those which carry the drainage of the northern slope of the hills inhabited by the Singpho, Aror, and Nága tribes, and fall directly into the Brahmaputra, and those which carry the drainage of the northern slope of the hills of the Lushái country and hill Tipura and of the southern slopes of the Jaintiya, Khasiya, and Gáro hills to the Megna, which joins the Brahmaputra below Dakka. To the former class belong the Dihing, Disung, Southern Dhansiri, and Kopili, and to the latter the Barak, Súrma, and Dhani.¹ To the extreme east of the Asám valley is a snowy range from which issues the Lohit or Brahmakund river that gives its name to the Brahmaputra; but geographers have applied the same name to the great river flowing by Lhása, and which the best authorities identify with the Dihong that joins the Lohit in the upper valley of Asám. The more general name of the Brahmaputra, in the upper portion of its course, seems to be 'Tsanpo' or 'Sanpo,' meaning 'the river' or 'the great river,' used like 'Ganga' in the plains and 'Kiang' in China. In Asám, the name varies with the tribe inhabiting its banks or those of its tributaries, so that the designation 'Brahmaputra,' to express the entire course of the river from its source to the north of Kumaon to its junction with the Ganges, must be considered a convenient-device of geographers, and not a term based on received usage. We have seen that at its junction with

¹ Any further discussion of the systems of the Asám valley would be out of place here; they are sufficiently described in the Asám Gazetteer, to which the reader in search of farther information on this subject is referred.

the Lohit, the Brahmaputra is called the Dihong.¹ East of this junction, the Lohit receives a tributary from the north, called the Dibong, and on the south another called the Dihing, and again one called the Disung. Much confusion has resulted in Asám geography from not remembering that all these names refer to different rivers.²

We have already noticed the division of Tibet into the eastern, western, and central provinces. It is called Tibet. Si-tsang by the Chinese, and is also known under the names Tu-pu and Mu-tu, or Upper and Lower Tibet. From the accounts of the eastern province in the writings of Klaproth, Huc, Blakeston, Cooper, Des Godins, and Gill, we may assume that the drainage to the east of the 96th meridian has a southerly direction, and that the general conditions resemble much those further west. Pengshan on the Yang-tse in western Sz-chuen is 1,500 feet above the level of the sea. At Ching-tu, some seventy miles due north of Pengshan, Cooper, in March, found the fields in the neighbourhood occupied by luxuriant crops of wheat, barley, sugarcane, and opium, the latter of which demands a climate similar to that of the plains below the Kumaon Himálaya. The same traveller crossed the Yalung and Kinsha branches of the Yang-tse between Chingtu and Bathang, and beyond Tatsien-lu entered Eastern Tibet, where yaks are used in the carrying trade and a more alpine climate is met with. Bathang on the west has much the same position with respect to the elevated highland to the north that Chingtu has on the east, but lies a degree of latitude more to the south than Chingtu; so that from the meridian of Bathang, the range dividing the Tibetan plateau from the plains of China seems to take a north-easterly direction. The feeders of the Yang-tse and other rivers find their way through this range by a series of gorges similar to that of the Dihong further west. Between the Yang-tse and the Dihong, we have two

¹ Klaproth suggested the Irawadi as the continuation of the Sanpo, and Colonel Godwin-Austen was the principal advocate for the Subansiri; Progn. R. G. S., August, 1876; but the researches of the Pandit employed by Lieutenant Harman perhaps show that the Dihong is the real representative of the Sanpo, and that the bend it takes beyond Chetang is quite sufficient to leave a large water-shed for the Subansiri. The most recent advocate of the Irawadi as the true continuation of the Sanpo is Mr. R. Gordon, in his elaborate "*Report on the Irrawaddy River.*" The first volume is illustrated with hydrographical hypsometrical and orographical maps of Tibet and the neighbouring countries together with a hydrographical map of India.

² See Peal's observations on Asám nomenclature in J.A.S., Ben.

great rivers, the Ian-tsang identified with the Mekong or river of Kambodia, and the Lu-tse or upper course of the Salween. Father Des Godins informs us that the Mekong has its origin in about 33° to 34° north latitude in the mountains south of Koko-Nor, and the Lu-tse further west is known as the Ngen-kio in Tibet; but whether this name is to be regraded as one with the Nák-chu-kha of the Pandit explorer is left to future travellers to decide.

Turning now to the upper portions of central and western Tibet, we find from the Pandit explorations that the drainage to the north of the northern water-parting of the upper portions of both the Indus and the Brahmaputra flows into a number of lakes. East of the 84th meridian, these lakes appear to be connected the one with the other, and eventually with the great river Nák-chu-kha, which has an easterly direction in the upper part of its course and finds a southern outlet in one of the great rivers to the east of the Dihong. From the Pangong lake on the west to the Lonkor Cho between the 83rd and 84th meridians on the east, the drainage is collected in a series of depressions without any outlet, each of which is the centre of a subordinate minor system of its own. We do not know of any considerable stream proceeding northwards or westwards from this tract. This lake-system is a characteristic feature of the orography of north-western Tibet. The waters of these lakes are generally brackish and the margins exhibit expanses of salt-marsh. Streams of fresh water are found, but in their course towards the lakes these rapidly become brackish, and in the end little influence the quality of the lake-water itself. According to the Pandit, the country to the north of Garge and Garchethol is a great uninhabited plain.¹ It was formerly customary to travel in a north-north-westerly direction from Thok Daurákpa² for some twenty days to the range overlooking the Gobi plain in which the commercial entrepôt Nári Tháru lay.³ A two months' journey from Thok Daurákpa to the

¹ Called Jung Pháyil Puyil, meaning literally "the desert country in which the father and son have wandered," from a tradition that two men had entered it and died there from want of water. G. T. S. Rep., 1873-75, p. 55.

² In about 85° 8' east longitude, and 32° 7' north latitude. ³ Trotter suggests that Nári Tháru occupies a position at the foot of the northern boundary ridge of the Tibetan plateau, similar to that held by Polu and Sorghák, and that the stream passing Nári Tháru may be the same that passes by Charchand. Prejevalsky (p. 76) says that Charchen is situated about 200 miles to the south west of Chargnik on a river of the same name. Hence it is a ten days' journey to the oasis of Nai (200 houses), and three days further to Kiria. The position and distance of Nai would suit the Pandit's description of Nári Tháru.

north-east brought the traveller to Ajan, also a commercial centre. This road lay throughout over an extensive plain ; no large mountains were seen and no streams were crossed. Drinking-water was obtained from a number of fresh-water lakes mostly dependent on the rainfall for their supply. From these reports we learn that northern Tibet is a plateau of great elevation without inhabitants and possessed of few streams.

As we approach the west, the boundary ranges that support the Tibetan plateau between them on the north and south gradually incline towards each other, so that westwards of the Pangong lake they are little more than fifty miles apart.² Here we find the water-parting of the Indus and Lob-Nor systems in the elevated Dipsang plain, which attains a height of 18,000 feet above the level of the sea near the Kárákorum pass (18,550 feet). From the Kárákorum pass on the north flows a feeder of the Yárkand river, and from the east, an affluent of the Kárákash river, both of which belong to the Lob-Nor system. On the south-east, only eleven miles from the pass, the Daulat-beguldi encamping ground is close to one of the feeders of the Nubra branch of the Shayok river that belongs to the Indus system. In one case the waters lose themselves in the Gobi desert, and in the other they reach the Indian Ocean at Karáchi. If the statements of Kostenko and Severtsof be accepted, we have here in the west an analogue of the arrangement that has been described as characteristic of all the river-systems along the southern face of the Himálaya between the Jumna and the Brahmaputra. The alpine basin of the Yamanyár lies between two great groups of peaks, some thirty miles apart. On the south-east of the Yamanyár is the Tágharma group, of which the Muztágh-Ata³ peak attains an elevation of 25,350 feet above the level of the sea. On the north-west there is a similar group, of which the Tásh-balik peak reaches an elevation of 22,500 feet. The Yamanyár collects its waters in an elevated valley

¹ On approaching Ajan, a bare rocky range is passed and the town is occupied by Sokpas, who procure their corn and flour from Kokod and Karka, a large monastery some ten or twelve days' journey beyond the borders of their country. Trotter suggests that Karka lies between Lob-Nor and Koko-Nor, and that Ajan is one with the Anj-si of Uspenski ; but both of these identifications require confirmation.

² From the Kárákorum to Shahídullah (11,780 feet) on the verge of the plains of Yárkand is a distance by road of 78 miles, and from the same to Changlung on the south is 83 miles, and to Leh on the Indus, 163 miles. ³ 'Father of ice-mountains.'

between these groups and the water-parting range on the west, which is known variously as the Tágharma plain or plain of the Kichik Kára-kul.¹ The groups themselves are in advance of the water-parting range and are connected with it by transverse ridges. That on the south connects the Muztágh-Ata group near the Kok Mainak pass with the water-parting range, west of the Neza-tásh pass, while that on the north has not yet been accurately defined. Spurs from the groups descend northwards towards the plains and effectually divide the waters of the Yamanyár from those of the Kashgár-darya on the west and from those of the Yárkand river on the east. It has generally been supposed² that these two groups of peaks belonged to a great meridional range; but Severtsof, who has had recently exceptional means for obtaining an accurate estimate of its character, distinctly states that "these two peaks were supposed to be connected by a continuous range, while the real fact is that each is respectively the highest point of separate small high mountain knots capped with eternal snow." We have, therefore, in the west also an apparent snowy chain of mountains as seen from the plains, but which on closer examination resolves itself into groups of snowy peaks in advance of the water-parting range from which they are divided by an elevated valley. This valley gives rise to a river that makes a way for itself between the boundary groups to the plains below, while the groups themselves are connected by transverse ridges with the line of water-parting. A ridge from Muztágh-Ata to Yangi Hissar separates the drainage of the northern slope from that of the Kinkol river on the east, and a second ridge follows at some distance the left bank of the Yamanyár which shortly after its issue from the mountains is absorbed in artificial branches or canals for irrigation purposes. The whole system is a remarkable illustration of Hodgson's formula for the river-systems of the eastern Himálaya.

The alpine affluents of the Yárkand river known as the Sarikol and Zarafshán rivers also illustrate the unitising principle observed elsewhere. They

Oxus basin.

¹ Lesser Kára-kul or 'black lake.' ²The two groups form a part of Hayward's meridional Kzil-Art range, the existence of which was supported by Blaw and Trotter and denied by Fedchenko, who considered the phenomenon described by Hayward as merely representing the terminal butt ends of parallel ranges belonging to the Pámir plateau. Kostenko, however, maintained it to be a great mountain range. Views of the Muztágh-Ata peak and the Sarikol valley are given by Gordon in his 'Roof of the World.'

are separated from each other by the Kandár range, which descends in a north-easterly direction from the water-parting range to the south, and terminates where the Sarikol and Zarníshán unite their waters within the outer range of hills to the north. The Oxus system further gives an example of another of the characteristic features of the Himálayan river-systems. The water-parting follows the line of the Hindu Kush from the pass near Bámian to the Baroghil pass, and thence follows the Pámír range to the greater Kára-kul. The height of the water-parting on the Baroghil plain has been estimated by Captain Biddulph at about 12,000 feet. The Mullah states that in traversing it for a distance of five miles there was no appreciable rise or fall, and further it is said that from a point one-and-a-half miles short of the crest, the difference in height did not appear to be more than two hundred feet. The Sarhad head of the Panjah branch of the Oxus is not more than two miles distant from the Gez-kul or Oi-kul, the longest source of the Aksu branch of the same river known as the Murgháb. The principal source of the western head of the Panjah branch of the Oxus in Wood's Victoria lake is but twelve miles distant from the water-parting between it and another branch of the Aksu. So little is this water-parting marked by any defined physical feature that it was only after some difficulty that Trotter discovered it at a height of 420 feet above the level of lake Victoria. There is also reason to believe that the greater Kára-kul once gave off at one end a feeder to the Kúshgár-darya of the Lob-Nor system, and at the other a feeder to the Oxus. We have already seen that it is but a little difference in perpendicular height that determines the drainage between the Satlaj and Karnáli and between the Nubra branch of the Shayok and the Kárákash, so that where rivers have their sources in these elevated areas it may be generally stated that a slight inequality in the surface, such as it is not possible to delineate on any ordinary map, is sufficient to determine the course of springs into channels that have a very remote debouche one from the other in the plains. The terms trough, channel, basin are in such cases often misleading. Nor are the bounding ranges in these elevated regions marked with such well-defined characters as are met with elsewhere. The great Pámír is divided from the Alichúr Pámír on the west by a range having an average elevation of only 3,000 feet above the level of the

Victoria lake, and from the little Páunir by a similar range averaging about 2,000 feet above the same level. The direction of the rivers depends, therefore, on the influence of much less relatively important masses of matter than are to be found at lower levels. The slight difference in level that determines the course of the head-streams of the Kárákash and Yárkand rivers on the Dipsang plains, is sufficient to divert the former from their normal direction and drive them directly against the Kuen-lun range, where, finding a fault in the wall, they work their way through towards the plains. The influence of the law of gravity and the mechanical and chemical changes wrought by water are the sufficient causes for every form of river channel that is met with, and it is to their ceaseless action that even the stupendous gorges of the Indus, the Satlaj, and the Brahmaputra, are due.

Having concluded our review of the river-basins and their relations to each other, we shall now proceed to examine the different parts of the area under our consideration. The great plain of Hindustán which first claims our notice is a vast flat extending with an almost unbroken surface along the foot of the Himálayan slope from the upper Indus to the Bay of Bengal. Its direction is from north-west to south-east over a distance of nearly 1,500 miles, and having an area, including its western branch along the Indus and its eastern prolongation into Asam, of about 500,000 square miles. On the west it has its greatest development stretching along the Indus from the foot of the mountains to the sea, from north-east to south-east for a length of 750 miles. Its breadth from the Arvali hills to those west of the Indus is about 400 miles. The Arvali hills run in a north-easterly direction from the peninsula of Kathiawár until they lose themselves in the plain near Dehli. From this point they run in a south-easterly direction connecting with the Vindhya, and in both cases constitute the abutments of the elevated plateau of Central India. They thus form two sides of a triangle with its apex towards the north, where it separates the Indus plain from that of the Ganges. The general slope of the Indus plain is south-west, with, as we have seen, a slight depression towards the south, until the influence of the northern slope of the Arvalis is felt, when it gradually rises again. Taking

a line along the Indus,¹ we have Seliwan, 117 feet above the level of the sea ; Shikárpur, 199 feet ; Dehra Gházi Khán, 395 feet ; Segra, on the eastern bank of the Indus, opposite Dehra Ismáíl Khán, 606 feet ; and Khairabad, opposite Kálábágh, 750 feet. Following the 32nd parallel from west to east, we have Sandi on the left bank of the Indus, 629 feet above the level of the sea ; Lodri, on the left bank of the Chináb, 657 feet ; and Rámdás, on the left bank of the Rávi, 796 feet : further east, we enter the hills. Following the 30th parallel we find Máre on the 71st meridian, with an elevation of 386 feet ; Núr Sháh on the 73rd meridian, 482 feet ; and Pakka Saráwa, on the 75th meridian, 698 feet. The perfect uniformity of the surface is broken in the north-west by the small table-land between the Indus and the Jhilam, of which the salt range² forms the abutment. These hills at the Sakesir station of the survey in the Jhilam district rise to a height of 4,994 feet above the level of the sea. They extend from Khairabad on the Indus to the Jhilam opposite Chilianwála, and thence those forming the eastern flank of the table-land as well as a subsidiary range to the east of the Jhilam turn abruptly north-east and connect with the outer ranges of the Himálaya near Bhimbar (1,200 feet). The table-land itself is seldom more than two or three hundred feet above the general level of the plain, and presents an undulating though tolerably even surface broken occasionally by ridges which attain a height of from two to three thousand feet.

The Indus plain along the foot of the hills is sufficiently watered, but to the east and south at any distance Indian desert. from the rivers cultivation on an extended scale is only possible when the scanty rainfall can be aided by artificial irrigation. The latter tract known as the great Indian desert stretches through Bhatíana, Bikanír, and Baháwalpur into Sind. Tradition tells us that in former time it was a fertile and populous country studded with numerous cities and towns and inhabited by prosperous and civilised tribes. A recent writer³ states that "there is nothing in history to show that the rivers (of this tract) ever contained much more water than they do now. Some diminution in their volume may have taken place during the

¹ All these heights are taken from the records of the Great Trigonometrical Survey. ² Fleming on the salt range, J. As. Soc., Ben., XXII., 229. ³ Cal. Rev., July, 1874, p. 3.

lapse of ages, from changes in the lower Himálayan range, as well as from the destruction of forests and increase of irrigation. There is no doubt also a tendency to the obliteration of the lower part of their course; partly by the drift of sand and dust from the desert; and partly by the deposit of the silt brought down by the streams themselves, owing to the absence of the great river by which it would have been carried off to the sea." Between the Sarasvati and the Satlaj are a series of broad channels, most of them a mile in width, of which those to the west appear to terminate in the valley of the Satlaj, while those to the east, which are also the more ancient, lead to the channel of the lost river Hakra or Sotra.¹ The Kaggar now runs in an old bed of the Satlaj and was formerly an affluent of that river. The change may have taken place owing to some great cataclysm² which formed a new bed for the river and left the old one for its tributaries on the east, and there is much to be said in favour of the identification of the Sotra channel with this old bed. At the present day the water-level in wells in this tract is excessively low, being often so deep as three hundred feet from the surface. As the water when procured is often brackish, it is a matter of wonder that people are found to inhabit this country which long ago received the name of Marusthali, 'the region of death.'

The Gangetic plain extends from the debouche of the Jumna from the hills to the head of the delta of the Gangetic plain. Ganges, and lies between the great bend of the Himálaya and the north-eastern slope of the table-land of Central India, which here has a general elevation of about one thousand feet above the plain. The breadth of the plain varies from about two hundred miles at Agra to about one hundred miles at Rájmahál. The direction is to the south-east, but to the east of the Ganges the courses of the rivers exhibit more southing. A general idea of the fall in height along the course of the Ganges may be obtained from the following figures: Saháranpur, 903 feet above the level of the sea; Meerut, 735 feet; Aligarh, 610 feet; Agra, 516 feet; Cawnpore, 517 feet; Allahabad, 315 feet; Benares, 255 feet; Patna, 174 feet; Bhágalpur, 159 feet; and Bardwán, 97 feet.³ Cross sections

¹ Said to have dried up in the thirteenth century. ² Like that of 1762, when, according to General Cunningham, the river was dammed up by a landslip in the hills and rose some four hundred feet before the barrier gave way. A similar cataclysm occurred on the Indus. ³ From G. T. S. tables.

show little difference in height at any point. Bhatpura, below Mohan, at the entrance to the principal pass into the Dehra Dún, is 954 feet above the level of the sea. Following eastwards the line along the foot of the hills, we have Hardwár, 1,016 ; Najíbabad 860 ; Barhapura, 910 ; Káshipur, 750 ; Bilahri, 760 ; Sigauli in Gorakhpur, 300 ; Madanpur in Tirhut, 230 ; and Amua, 248 feet.¹ Noting these figures on any good map and following the course of the rivers, a sufficiently correct idea of the general slope of the Gangetic plain will be obtained. At the termination of the hills near Rájmahál, the plain once more expands largely to the south and again presents an uninterrupted surface from the mountains to the sea. The length of this section is about 350 miles, and its breadth from Rájmahál to the Brahmaputra about 150 miles, but increasing to about 300 miles along the coast at the head of the Bay of Bengal. The height varies from 100 feet, the level of the river Mahánadi at Málda,² to 75 feet at Jelinghi, the head of the Hugli branch of the Ganges,³ and 31 feet at Chinsurah. The Howrah station bench-mark is but 18·2 feet above the level of the sea.⁴ These portions of the great plain, often though not very appropriately called the valley of the Ganges, are intersected by the countless tributaries of that river and are under the full influence of the periodical rains. They therefore, as might be expected, comprise the richest, the most populous, and the most civilised districts of India, and in these respects form a striking contrast to the western parts along the Indus which are doomed to perpetual sterility, not from any natural deficiency in the quality of the soil, but only from the great aridity of the climate. This barrenness is no doubt, in a great measure, due to the relative position of the Indus plain to the higher ground around it and to the prevailing winds ; matters which, at first sight, appear to be of little importance, but which are the efficient causes of the extremely dry climate that it possesses. Fluvial action in erosion and deposition, productive of the alluvion and diluvium, terms so well known in the settlement records of these provinces, has clearly directed the course of the rivers in the great Gangetic plain. Mr. Fergusson, in an article quoted by the authors⁵ of the 'Manual of the Geology of India,' shows that the

¹ First five from Webb and Sigauli from Kirkpatrick.

² Prinsep.

³ G. T. S., 52 : in sc. VIII. 1866.

⁴ Hooker
⁵ p. 406. 412.

rivers of the Ganges delta oscillate in curves, the extent of which is directly proportional to the quantity of water flowing down their channels. Thus, the oscillations of the Ganges between Mungir and Rájmahál average $9\frac{1}{2}$ miles, and between Allahabad and Chunár, only 3·7 miles in length. Further, when a great river runs through a low country, its course is considerably stayed by the sluggish expanses of stationary water generally termed *jhíls* in Bengal, and it is thus compelled to deposit its silt along its banks. Hence arises the phenomenon of a river passing through a country between banks that are higher than the adjacent alluvial flats, and the gradual increase of the banks until the stream makes its way through them to some lower level. Mr. Fergusson estimates that when the slope of a river bed falls to less than six inches in a mile, a denuding river will be converted into a depositing river, and as the deposit commences at the bottom of the slope, the change proceeds upstream. Moreover, since the Ganges receives its more considerable affluents from the north, the left bank gradually increases and drives the main stream more and more towards the table-land of Central India, and makes the point of confluence of its affluents continually move upwards. This tendency is well marked in the Jumna in the Mainpuri district, where the old silted-up bed is locally known as the *bhagna*.¹ In the Tarái below Kumaon the same law prevails, and streams that in the upper portion of their course are denuding rivers in the lower portion where the check in slope occurs deposit their silt, form *jhíls*, and continually change their courses like the rivers of the Gangetic delta.

The Asám valley forms a narrow prolongation of the eastern extremity of the plain, partaking more, however, of the nature of a simple river valley liable to annual floods. It has a length of about 300 miles with a breadth of thirty to forty miles, widening at its junction with the

Asám valley.

¹ Gaz. IV., 478. The old bed in Saháranpur is called the Budhi Jumna.—*Ibid.*, II., 140. Remains have been found in the Jumna alluvion near Allahabad.—J. A. S. Ben., III., 302, 529: IV., 262, &c. Wilford also notices the discovery of bones of men and animals in the Ganges alluvion near Benares about a furlong from its present bed at ninety-five feet from the surface and thirty feet below the level of the present bed. He says the human bones were entire, but those of quadrupeds were broken and bore evident marks of their having been cut with a sharp instrument. He found no marine deposits at over 105 feet when the water-bearing strata was reached.—A. S. Ben., VIII., 294. He has also noticed the fact that the point of junction of the Kosi with the Ganges had moved up from Nawábganj opposite Rájmahál some twenty-five miles.

Gangetic delta. It is shut in on all sides, except the west, by mountains. The characteristic of a sudden and total change along a definite line at the foot of the mountains from a broken hilly surface to an absolute flat, which holds good throughout the other parts of the plain, here no longer prevails. We frequently see small isolated hills standing out like islands from the general level surface, a phenomenon observable on a smaller scale where the Vindhyan range mingles with the plain. From Sadiya, which is 440 feet above the level of the sea, the fall is gentle to Gauhati, which is placed at 163 feet, and Goálpára, which is 150 feet.¹ Taking a line along the foot of the hills, we have Titaliya, 330 feet; Rájhát at the foot of the hills near Buxa, 220 feet; and the station at the foot of the hills below Diwángiri, 670 feet: figures² which show a more sudden descent than those at similar positions under the western Himálaya.

We shall now return to the plain between the Jumna and the
 Bhábar. Sárda, and more particularly to that part
 which lies immediately below the foot of the
 Kumaon hills. We find there a narrow belt of country usually covered with forest and remarkable for the entire absence of water, a phenomenon eminently characteristic of this tract. The great rivers preserve their course with some diminution in their volume, but all the minor streams that have their origin in the lower hills on entering this belt soon lose themselves in the shingly deposit that constitutes the substratum. In time of flood, however, they often preserve a visible stream throughout their course, but this appearance lasts only so long as the cause exists. This belt of waterless forest land is called the Bhábar or *ukhar bhúmi* (waterless forest) under Kumaon, and has a breadth of from five to fifteen miles. Though no stream or spring exists, the Bhábar is clothed with a magnificent forest finding its nourishment in the few foot of alluvial matter that rests on the boulder and shingle deposit below.³ To the

¹ These figures are from the Great Trigonometrical Survey records, which make Dibrugarh, 848 feet above the level of the sea; Silmágar, 319 feet; and Baramak, near Teapur, 456 feet.

² First from Hooker, two last from Pemberton.

³ These descriptions are chiefly based on Herbert's report; and on an article entitled 'The Himálaya in Kumaon and Garhwál' by Mr. (now Sir John) Strachey, Cal. Rev. No. 36. Stewart notes that he can find no definite statement as to the breadth of the tract known as Bhábar and Taráí to the east of Sikkim, but Hooker mentions expressly that there it ranges from 8 to 12 miles in width; Kirkpatrick and Hoffmeister coincide in making its breadth opposite Nepál about 10 or 12 miles, and between the Sárda and Rámanganga it ranges from 20 to 30 miles, falling to 6 or 7 miles abreast of Garhwál and disappearing to the west of the Jumna.

south of the Bhábar, the character of the country changes into a swamp devoid of trees and intersected by sluggish streams that rise

from unhealthy morasses. This tract is
Tarái.

included with the Bhábar between the

Ganges and the Phika under the general term *ban* or jungle. East of the Phika under Kumaon, where it apparently attains its maximum breadth, it is known as the Tarái, and under Nepál as the Taryáni, with some specific addition as Morang, &c. It lies between the forest belt and the cultivated plains, with an average breadth of about ten miles under Kumaon, though varying much in different parts. Thus we have between the plains proper under Kumaon and the foot of the hills two distinct belts of country, each about ten to fifteen miles broad, known as the Tarái and the Bhábar. The Tarái is characterised by the presence of reeds and grasses showing the marshy nature of the ground. The streams carry off only a portion of the superfluous moisture and sluggishly run in tortuous channels, doubling back constantly in their course. The soil consists of moist alluvial matter without a sign of rock either in fragments or in site. In the Bhábar, on the other hand, no water rises from the ground. Throughout its whole extent not a single spring nor any water can be seen, except occasionally where one of the larger rivers takes its course. In the rainy season alone torrents cut into the ground, and the channels thus formed exhibit characteristic sections of this remarkable tract. We then find that there is but a thin covering of alluvial soil on a vast dry bed of boulders and of shingle, through which all rain that falls sinks rapidly, and which absorbs in the same way all the minor streams of the outer ranges. Instead of reeds and grasses, we have here all the magnificence usually attributed to oriental forest scenery. Gigantic *haldís* (*Adina cordifolia*) and *khaírs* (*Acacia catechu*) rear their heads above a tangled undergrowth of creepers and thorn-bushes which present a barrier to progress that an elephant alone can surmount. Towards the hills we find the *sál*¹ (*Shorea robusta*), and in Kota great groves of mangoes, while patches of cultivation appear wherever irrigation is practicable. For this

¹ The *sál* is the characteristic tree of the upper Bhábar wherever it is found. Hodgson notes that constant observation has enabled the people of the Tarái to distinguish the principal belts of the Bhábar from the trees that grow in each. The highest is the *sál* level, the next is the *khaír* level, and the lowest is the *sím* (*Dalbergia sisu*) level.

purpose, the streams of the lower hills are turned into artificial channels before they reach the shingle deposit, and even the lakes in the lower hills are dammed up to retain a sufficient supply of water for the Bhábar. West of the Kosi, however, there is little cultivation or irrigation, and the Bhábar there almost remains untouched by the plough. The actual slope of the ground between the Taráí and the foot of the hills is considerable, though not apparent to the traveller, except when he observes the rapidity of the current in the irrigation channels that line the road by which the Bhábar is crossed.

Before entering into more detail regarding the Bhábar and Taráí, there is yet a third feature characteristic of the tract below the Himálaya that must be noticed here as intimately connected with the other two, and this is the line of hills called the Siwálik¹ or sub-Himálayan. These will be well known to palæontologists in connection with the rich collection of fossil mammalian bones discovered in them by Dr. Falconer and Colonel Cautley. As a rule, they appear to rise abruptly and without any intermediate undulating slope from the apparently level surface of the flat country below to heights varying from a few hundred to three or four thousand feet. They are composed of sandstones and conglomerates, and the dip of the strata is usually towards the general mass of the mountains at a low angle. The form of disturbance of the strata is very regular, producing broad normal anticlinal flexures, the axis-plane sloping towards the mountains. Towards the plains the slope has been weathered out, so that plainswards the Siwálik exhibit a steep face from which rise the highest summits of the range, while a long gentle declivity slopes inwards and forms a longitudinal shallow valley by meeting the foot of the next line of hills. The latter, as a rule, run on a line parallel to the Siwálik, but at a distance of from five to ten miles from them.

The bottom of this longitudinal depression is, as may be supposed, by no means continuous. In some places it is cut through by the passage of the streams that drain the interior of the mountains; in others, it is quite obliterated by the near approach to each other of the two ranges that flank it, and which usually form distinct lines. This is, moreover, a

¹ Siwawála, belonging to Siva; for use of the term 'Siwálik' by Musalmán historians, see *History* *protea*.

structural feature and not due simply to denudation. In the country between the Satlaj and the Káli, these valleys are called Dúns and under Nepál, according to Hodgson, they are called Máris. They have been confounded by some writers¹ with the Tarái, which, as we have seen, is quite distinct. The lower part of the Dúns generally appears to be covered with a deposit of boulders and gravel that slopes somewhat steeply from the Himálaya towards the Siwálik, so that the whole bottom of the valley is considerably raised above the level of the plain without. In consequence of this elevation, the outer hills when viewed from the interior of the valley, as from Masúri, present a very insignificant outline. The drainage of these valleys usually collects along their longitudinal axis and either falls into some of the larger streams that cross them, or less frequently finds an independent exit for itself into the plains by a sudden bend to the south through a break in the outer range. Owing to the considerable elevation of the Dúns above the plains down to the level of which the drainage finds its way in a very short distance, the unconsolidated strata that form the floor of these valleys are constantly cut through to a great depth by water-courses. Consequently, the surface, though often presenting an apparent flat for several miles together, is frequently broken up into steps which, on the whole, are tolerably level, but at different heights, the one above the other. This phenomenon is not uncommon, and is constantly observed along rivers that are eroding their banks. To the same causes also are to be attributed the practical impossibility of procuring water by means of wells in the Dúns, a difficulty which mainly arises from the thorough dessication of the gravelly soil by the deep drainage.

We have not sufficient information to state distinctly how far the Bhábar extends both east and west along the foot of the Himálaya, but the following indications would lead us to suppose that this phenomenon is inherent in the relations of such a mass as the Himálaya with the subjacent plains. Under Nepál it is called the Jhári or Bhávar, and, according to Hodgson,² extends from the Káli to the Mechi on the east with the same general characteristics as under Kumaon.

¹ Somerville's *Physical Geography* corrected in 7th edition, 1878.
Geography of the Himálaya.—J. A. S., Ben., XVIII., 778.

² *Physical*

Eastward of the Tista, according to the same writer, the Bhábar and Tarái do not exhibit the same parallelism to the line of the Himálaya, but "show themselves plainwards, like an irregular series of salient and re-salient angles resting on the mountains. Or like small insulated plateaus or high undulated plains, surrounded in both the latter cases by low swampy land analogous to the Tarái." An example of the former is found in the plateau called the Parbat Joár on the confines of Asám and Rangpur, which is considerably elevated, quite insulated, remote from the mountains and covered with *śál*, the characteristic tree of the upper Bhábar. Again, we have undulating plains, such as those that occur around Dinajpur and to the north-west and north-east of Siligori, all of which may be identified with the Bhábar. In all these cases where the detritus bed thins out, a moist tract is met with, though in no case so marked as to the westward. Herbert¹ affirms the general applicability of his remarks regarding the submontane tract below Kumaon to the entire country between the Ganges and the Satlaj and Parish² to the tract further west between the Satlaj and the Biás. There is no well-defined line dividing the area of swamp from the Bhábar proper between the Sárda and the Rámghanga. To the east in the Tallades Bhábar, where the streams seek the Sárda directly, there is less Bhábar, and the swamps that exist are not so extensive, but at the same time are more formidable, being often surrounded by tangled masses of canebrake. The Dhyánirau Bhábar also is comparatively narrow, and it is not until we come to the Ohhakháta Bhábar that we get a breadth of eight to twelve miles that lasts until the Phíka river is reached. The Tarái exist all along the tract to the south of, and parallel to, the Bhábar from the Sárda to the Phíka. But west of the Phíka it loses its characteristics and can only be traced in the closeness to the surface of the water-level in wells. Westward of the Phíka, the Bhábar or waterless tract also narrows and the *śál* forest does not descend more than six miles from the foot of the hills, and a few miles further west it has not a breadth of two miles. The Bhábar, however, exists, and is broader than the present *śál* forest, of which much has been cut down of late years. Its presence is shown by the absence

¹ Report of the Mineralogical Survey of the Himálaya Mountains, *Ibid*, extra No., Vol. XI. ² Kohistán of the Jullundhur Doab, *Ibid*, XVII., p. 281: XVIII., p. 300.

of wells, and similarly the Taráí appears in a line of wells with water at from three to six feet from the surface, running parallel to and bordering on the Bhábar. This limitation of both the Bhábar and Taráí is conterminous with the commencement of the Pátlí Dún, which has detained the greater part of the detritus that is elsewhere spread out below over the plains. From the Phika eastwards to the Sárda, where these tracts attain their maximum importance, there are no Dúns, properly speaking; for the Kota Dún presents no great barrier to the south, and further east the Siwálíks are so blended with the outer range that a geologist alone can trace their sequence. In this fact, we have an illustration of that portion of Hodgson's theory that gives a narrow extent to the Bhábar below the Dúns and a broader range where there is no Dún to intercept the débris from the hills. The facts that we know regarding the Bhábar to the north of the Saháranpur district¹ further confirm this deduction.

In Eastern Turkistan we find a similar phenomenon. Trotter tells us that the Sagon river, which has its source at the eastern foot of the Terek pass, after it reaches the plains north of Káti Ailak wastes away and leaks through crevices in the stony ground. The hákim of the latter place assured him that wells had been often sunk but proved of no use. Trotter writes :—"This diminution in the size of rivers as they descend is one of the chief characteristics of the country, and occurs in all minor streams that have come under my notice. Of course much of this is due to irrigation, which necessarily carries off large quantities of water, but the stony soil has also much to answer for; on the other hand, the frequent appearance of large springs giving considerable supplies of water and often issuing from the open plains at long distances from the mountains may account in a great measure, if not fully, for the water thus lost in its early infancy." Here we have the existence of a Bhábar and Taráí vouched for by competent authority in the Yárkand and Kashgár country. The same phenomenon, but on a larger scale, was found by Prejevalsky to characterise the tract between Korla and Lob-Nor. A belt of country about three to four miles wide, consisting of an undulating plain covered with a pebbly or gravelly soil and totally devoid of

¹ Gazetteer, II., 140.

vegetation, runs parallel to and at the foot of the Kurugh-tágh, a low waterless and barren range. Beyond this stony margin, which appears to define the shore-line of an ancient sea, lies the great desert of drift sand amid which salt marshes exist wherever the moisture comes to the surface. The same pebbly plain was found under the northern slope of the Altyn-tágh, the north-easterly continuation of the Kuen-lun mountains between the 90th and 92nd meridians, and north of and below the stony margin the usual salt marshes occurred. The latter are also found at the foot of the north-eastern portion of the Tibetan table-land in Tsaidam. It would, therefore, appear that tracts analogous to the Bhábar and Taráí of Kumaon surround the entire Himálaya-Tibetan mass, and that they vary in character according to local influences.

Hodgson attributes the distinctive character of the Bhábar, as a Cause of the deposit: whole, "to the vast mass of diluvial detritus Hodgson's oceanic theory. which was shot from the mountains upon the plains, like gravel from a cart, at some great geological epoch, and which has been, since its deposit, variously and often abraded both in degree and direction by oceanic and, in a far less degree, by ordinary floods." Another writer considers that this theory of Hodgson's appears to be a reasonable explanation of the existence of these great beds of shingle, sand, and boulder all along the foot of the mountains. It is argued that no rivers can have laid out such a vast deposit, and we can only conclude that we see here the limits of an ancient ocean that once washed the base of the Himálaya. The boulders and shingle are spread out only for a distance of ten or fifteen miles from the mountains from which they are derived, while the finer particles of sand and clay are carried much further. Great variations in the depth and breadth of the deposit occur, due, in a great measure, to local causes. One which apparently has had a great influence is the existence or otherwise of the Siwálik range. Where there was no sandstone range to intervene between the mountains and the plains and collect the detritus within their contained Dúns, the deposit is broader and not so thick. Where there was such a barrier, it has been carried less southwards and exists in great accumulations between the barrier and the mountains. Again, where no range existed but only spurs sent forth, like bent arms, upon the plains from the mountains, Hodgson observes that the embayed

detritus is simply deeply piled and lofty within such spurs, and thinly and unequally spread without them, by reason of the action of the spurs on the current. He notices, as an example of this form, the *débris* embayed by a spur on the road to Darjiling by Pankabári, where it is accumulated to several hundred feet, and where, moreover, there is outside the spur a succession of terraces, apparently due to oceanic forces.¹ Further, "where, as from Gauhati to Sadiya, there was not room upon the plains for the free spread and deposit of the detritus owing to large and rapid rivers and to other chains both proximate and parallel to the *Himálaya*, the phenomena, created elsewhere by the more or less unrestricted spread of the *Himálayan* detritus over the plains, would necessarily be faintly, if at all, traceable. Lastly, if at the time of the descent of the *débris*, there existed a great dip in the Gangetic plain from north-west to south-east, the lithologic character, as well as the distribution of the *débris*, would be materially affected thereby, for the subsiding oceanic current would have a set from the former to the latter quarter and would continue to lash the gravel into sand and here to deposit both in a series of terraces, there perhaps utterly to displace both in the latter quarter long after the former had emerged from the waves."

The oceanic theory of Hodgson is not accepted by the majority of professed geologists. Mr. W. Blandford

Fluvialile theory.

writes:²—"There is absolutely no proof of

any sort or kind that the whole Indo-Gangetic plain has at any time been a marine area; but there is equally no proof that it has not. It has been shown that in eocene times the sea occupied the Indus valley as far as the foot of the *Himálaya*, and extended along what is now the base of the mountains, as far east as Kumaon; and also that marine conditions prevailed to the north-west throughout a great part of the tract now occupied by the *Asám* range; but it was also pointed out that, in the area between Kumaon and the *Gáero* hills, no trace of marine formations had been found. Yet it is difficult to understand, if the Gangetic plain was a sea-basin, why no marine beds occur. It is true that the northern border of the plain, throughout the most important part of the

¹ The oldest advocate of the oceanic theory is Wilford (in *As. Res.*, VIII., 393), who thus accounts for several statements made by the Pauránik geographers, and particularly the story of *Sagara*. He, however, acknowledges that the existing soil of the Gangetic plain, so far as is known, was due to fluvialile action.

² *Manual of Geology of India*, p. lx., 393.

intervening space in Nepál, is, unfortunately, inaccessible to Europeans; but still, if the Gangetic plain in any way corresponds to an eocene sea, as the Indus plain doubtless does, why are no traces of marine beds found to the south of the valley on the margin of the peninsular area, as they are in the desert to the east of the Indus? In the Brahmaputra plain, also, no marine deposits of tertiary age are found; in the plain itself only fluviatile deposits have been detected and the marine, eocene, and miocene beds are confined to the southern slope of the range forming the southern watershed of the valley." Mr. Blandford considers the post-tertiary formation of these provinces to be clearly river deposits. The latter tertiary formations belonging to the Siwálik series contain reptilia and mollusca, but not a single marine shell. "It is impossible to tell what beds may be concealed below the Indo-Gangetic alluvium, and marine strata may exist to an enormous extent without appearing at the surface; it is also unquestionable that the amount of information hitherto derived from borings is very small indeed, but so far as that information extends, and so far as the lower strata of the alluvial plain have been exposed in the beds of rivers, not a single occurrence of a marine shell has ever been observed, nor is there such a change in the deposits as would render it probable that the underlying strata are marine * * *. The only evidence known in favour of marine conditions having prevailed during the deposition of any portion of the Gangetic alluvium is the occurrence of brine springs at considerable depths in a few localities. These springs, however, are not numerous, and, without additional evidence, it is impossible to look upon them as proofs of marine deposits. At the same time it is by no means impossible that the sea occupied portions of Sind and Bengal long after the plains of Upper India were dry land." On the whole, Mr. Blandford thinks that the oceanic theory wants further support; that the fluviatile theory is the only one that fits in with the present state of our knowledge, and that the depression of the Gangetic plain is of contemporaneous origin with the disturbance and contortion of the Himálayan ranges, and that the physical features of the two areas are closely connected. No important borings have ever been made in these provinces, and nothing has ever been discovered, so far as we are aware, to show that the older theory is the correct one. The

newer theory is further supported by the discovery of the buried town of Behat in the Saháranpur district, some seventeen feet below the present level of the country and containing coins of the commencement of the Christian era, thus showing what can be effected by fluvial action in eighteen centuries.

We have already seen that the distinctive features of the Tarái are not found west of the Phika river, if we Tarái. except a small tract on the left bank of the Rámganga, the condition of which, however, is probably due to defective drainage in that particular part, and might occur in any other place. The existence, therefore, of the Tarái as a distinctive feature must be due to local causes capable of explanation, but the imperfect nature of our knowledge will only allow us to guess at them. Herbert¹ described the Tarái as "defined in its southern boundary by a rise or step which runs parallel to the common boundary of mountain and plain land." He observes the height is variable, occasionally as much as thirty feet and sometimes sudden and steep. Modern research can discover no well-defined boundary beyond the chain of springs which sometimes approach within a couple of miles of the foot of the hills and sometimes are separated from them by a belt fifteen miles wide. In no case is there any such rise or step as described. Hodgson² also accepted the existence of a longitudinal trough running parallel to the Himálaya as a characteristic of the Tarái, which he held to be a natural depression in the plain, and thus accounted for its peculiarities. This theory, however, is opposed to the results obtained by levelling operations and appears to be based on an entirely erroneous idea, the fact being that the drainage of the higher country, beyond which has been lost in the absorbent strata of the Bhábar, here breaks out again in a line of copious springs which collect into swamps in the Tarái. This feature has also somewhat plausibly been accounted for³ by the existence of an impervious stratum below the absorbent boulder detritus, and as the latter gradually thins out the finer and less permeable silt underlying it approaches nearer, and eventually reaches the surface, bringing with it the water that has been absorbed by the shingle talus and has been retained by the impervious silt.

¹ *l. c.*, art. 73.
p. 184.

² *l. c.*, p. 788.

³ Batten in Kumaun Reports,

FIG. 1.




(a) Point of re-appearance of water. | (b) Point of disappearance of water.

Although this explanation seems reasonable so far as it goes, it must be remembered that the swampy Tarai extends only from the Phika to the Tista, and we must, therefore, look for some peculiarity in this part of the plain which does not exist elsewhere, by which we may account for the existence of swamps exclusively in this particular locality.

In a recent note, Mr. Lawder gives the following section of the Bhabar and Tarai :—

FIG. 2.



His experience of this tract has led him to consider that, whatever may be the nature of the underlying formation, the surface beds are solely due to fluvial action. The mountain torrents along the foot of the Kumaon hills bring down every year a vast amount of débris which is spread out over the surface now on one side of their previous course and again on the other. This irregular deposition itself compels the torrents to change their beds from place to place until, as now obtains, the points where they debouch from the hills are marked by more or less irregular, great, fan-shaped boulder and gravel deposits. The clayey or semi-soluble particles are necessarily carried farthest and are readily deposited not only where there is a check in the slope, but where the current is impeded

by the tortuous nature of the channel which itself naturally assumes that form under these conditions. Here, during the rains, the streams saturated with clayey silt, overflow their banks, form new channels, fill up old ones, and create the Tarái. Above this deposit of clay we find one of clay combined with sand, in which, however, the latter predominates. From this bed issues the line of Tarái springs that flow uninterruptedly throughout the year, and its margin marks the northern boundary of the Tarái. Above we meet beds of sand and gravel or gravel and boulders as we approach nearer to the hills. A longitudinal section taken at the top of the Bhábur (Fig. 2 B.) will show that the hill torrents in the upper portions of their course run along a ridge formed by the débris transported by themselves, whilst a similar section of the Tarái (Fig. 2 C.) would show that, as a rule, the river channels are found in depressions below the general level of the country. The geological section (Fig. 2 A.) shows the gradients of the present ground surface on the road between Bareilly and Ránibágh, and from them it will appear that in the boulder region deposition takes place at a slope of sixty-six feet to the mile, whilst the clay is not deposited until the descent falls to about eight feet in the mile. It may fairly be assumed that these are the usual angles of deposition of the materials, and that they have obtained since the degradation of the lower hills and the resulting deposition below them commenced. If so, a series of proportionate curved lines running almost parallel to the present ground surface may be taken to represent the ground surface of succeeding periods, and such portions of these lines as may be similarly inclined with the present Tarái portion (i.e., at the same angle with the horizon) will evidently represent the Tarái or clay deposit as it then existed. A line intersecting all these beds at the several points of junction of the 'clay' with the 'sand and clay' will represent the present impermeable bottom of the Bhábur basin and account for the line of springs upon the surface where the stratum of sand and clay crops out. The upper boulder and gravel beds permit of the filtration of water freely through them to the clay, at the same time acting as a capillary reservoir to keep up the dry weather supply to the springs below.

We have further evidence in support of this theory in the fact that the Tarái proper does not extend westwards of the Phika river.

Between the Phika and the Sárda there are no Dúns, for the Kota Dún has its southern boundary broken through by rivers, and along the entire tract numerous torrents find their way directly to the flat country below. The proximity of these torrents to each other causes the accumulation of débris to exhibit a continuous appearance which seems to have suggested the theory of a marine origin. To the west of the Phika river, the drainage of the lower hills is carried off mainly by streams which collect the drainage within the Sub-Himálayan range and seek the plains in one well-defined channel. The Rámghanga is the great arterial drainage channel for lower Garhwál, and between it and the Ganges, the only considerable stream is the Khoh, which has a small strip of Bhábar below it. In eastern Kumaon the Ladhiya serves a similar purpose, and where in its course towards the Káli it approaches the plains and does not allow of any considerable stream from the southern face of the outer range, both Bhábar and Tarái are narrow, and as this influence of the Ladhiya on the east and the Rámghanga on the west decreases the Bhábar and Tarái increase and eventually attain their maximum breadth where that influence is least felt. Where rivers discharge large volumes of water like the Ganges and the Sárda, and in a lesser degree the Rámghanga and Kosi, the velocity at their debouches from the mountains is much less than that of the minor torrents, owing to their having cut back and more deeply their channels within the hills, so that only the lighter particles of eroded matter are carried onwards, whilst the boulders are left behind at their natural point of deposition. Hence, near these larger rivers it curiously happens that the width of the Bhábar and Tarái contracts in a certain ratio and in the case of the Ganges disappears.


This explanation is supported by the results obtained during the contour survey of the Tarái.¹ The second diagram² (Fig. 3) shows a portion of the country between the Dhora river and the Bhúta stream, taken from the survey maps, and will illustrate the intricate nature of the levelling operations, and show why in some

¹ See Proceedings, N.-W. P., P. W. Department, May 31, 1864; July, 1869; February, 1872.

² From the Rohilkhand remodelled canals' contour map of a portion of the Tarái surveyed in 1865-67 by Captains Thomason and F. Brown and Lieutenant Bisset. An examination of the records of this survey shows that what is stated regarding the portion noticed here is true also of the entire tract between the Phika and the Sárda.

places the streams double back on their original direction and exhibit the tortuous courses so characteristic of this tract :—

FIG. 8.



It will also be observed that there is a sudden check in the slope where the Tarāi commences ; to the north in the Bhābar the slope is from sixty to a hundred feet in the mile, and in the Tarāi it falls to about ten feet. These are the adequate causes of the existence of swamps, and though the neglect of artificial obstructions made for the purpose of utilising the water for irrigation may doubtless aggravate the natural defects of drainage, it would probably produce no effect whatever were it not for the peculiar physical conditions that exist here. To the west between the Indus and the Ganges, the great arterial drainage lines collect within the hills and run off directly from them, the general fall of the surface receiving no such check as is found under Kumaon. The same is true of the country to the east along the head of the Bay of Bengal, and in the narrow valley of Asām, the Brahmaputra runs in a deep bed at right angles to the natural course of the streams from the hills, and thus forms a perfect system of cross-drainage that does not allow of the formation of swamps.

The Siwāliks appear to have a more or less definite existence along the whole of the Himālaya from the Indus to the Brahmaputra, presenting modifications of the same general features along the entire line. To the eastward of the Tista they are wanting locally, a fact which it has been suggested is due to denudation as in the case of the partially obliterated barrier to the south of the Kota Dún. As the Siwāliks will be noticed hereafter in the chapter on the geology of Kumaon, we need not describe them here.¹ Between the Jumna and the Sārda they are found as the southern boundaries of valleys as far eastwards as the Nihāl river, and thence onwards they almost coalesce with the outer range of the lower Himālaya.

Of the Dúns or valleys, between the Siwāliks and the Himālaya, that known as ' the Dún ' or Dehra Dún, Dúns. from the town of Dehra, is not only the most remarkable but the best known. Since the physical geography of this tract will be considered in more detail hereafter, in the notice of the Dehra Dún district, it will be sufficient for our purpose here to note that the Dún, a little to the west of the town of Dehra, is

¹ See also Chapa. XXII-IV. of the 'Manual of the Geology of India,' and Gaz., II., 46.

divided by a ridge that serves as a water-parting between the Asan, a tributary of the Jumna on the west, and the Suswa, a feeder of the Ganges on the east. The tracts drained by these rivers are known respectively as the western and eastern Dúns. The two taken together have a length of about forty-five miles and an average breadth of eleven miles. The east end of the Dehra base line of the Great Trigonometrical Survey on the extremity of one of the spurs of the Gháti range, about one mile west of Mahobawála, is 1,957·65 feet above the level of the sea : Mahobawála itself is 2,096·56 feet and Dehra is 2,323 feet. The junction of the Suswa and the Ganges is little more than 1,000 feet above the level of the sea, giving a considerable fall for that stream between Dehra and the Ganges. A well sunk by Mr. Shore, when Administrator of the Dún, attained a depth of 221 feet before a plentiful supply of water was met with, and even at that depth the nature of the deposit was the same as at the surface.¹ The greatest thickness of the deposit is observed near the central ridge. It thins out to the west and east along the course of the Asan and the Suswa, and, according to Herbert, may be observed in the beds of the Jumna and Ganges resting on sandstone. Next, on the west, comes the Kayarda Dún,

¹ The following table shows the character of the stratum, and is reproduced here from Mr. Shore's notes in the Dehra archives compared with Herbert's record as one of the few notices of this character that we possess.

Feet.	Total.	Soil.	Feet.	Total.	Soil.
5	5	Fine black mould, few stones.	9	158	Sand, larger pieces of conglomerate.
4	9	Reddish earth with gravel.	4	162	Do., enormous stones
9	18	Sand, gravel, large stones.	6	168	Conglomerate and gravel.
2	20	Do., with reddish clay.	3	171	Sand, gravel, and conglomerate.
3	23	Stiff reddish clay.	3	174	Blocks of conglomerate.
8	31	Do., yellow clay.	3½	177½	Sand, gravel, conglomerate.
3½	34½	Sand, gravel, little red clay.	4½	182½	Do., gravel.
1½	36	Stiff reddish clay.	4½	187½	Conglomerate.
2	38	Sand and gravel.	2½	185	Sand and clay.
22½	60	Stiff red clay.	18½	204	Loose sand and gravel and large blocks conglomerate.
2	62	Clay, sand and gravel.	5	209	Sand, gravel, very moist.
16	78	Sand and gravel.	4	209½	Conglomerate, over half.
12	90	Stiff yellow clay with little sand.	13	211	Red clay.
35	125	Sand, gravel, few round stones.	7	218	Sand and gravel.
6	131	Sand, large conglomerate blocks.	8½	221½	Brackish clay with angular fragments of clay slate.
13	144	Do., gravel, stones.			
5	149	Do., larger stones.			

Another well in Dehra gives water at a depth of 88 feet from the surface, and one at Sati-bagh at 75 feet. There are only twenty-nine wells in the whole valley, and many of these are at favourable places near rivers, so that the difficulty of procuring water by this means is common to the whole Dún. Herbert records the

some six miles broad and twenty-five miles long, to the water-parting between the Junina and the Kaggar systems. Beyond it we have the Pinjor Dún, which has in parts a breadth of six miles and a length of about thirty miles. The Siwálíks here are neither so broad nor so high as to the south of the Dehra Dún. The Pinjor Dún is divided into two parts, the eastern and western Dúns, by a ridge similar to that observed in the Dehra Dún, and which attains an elevation of 2,402 feet¹ above the level of the sea. To the east, the drainage flows into the Kaggar, and to the west into a tributary of the Satlaj. Mani Devi, a temple in the plains just within the Dún, has an elevation of 1,263 feet,¹ giving a fall to the rivers within this Dún similar to that found to exist within that of Dehra.

East of the Dehra Dún we have the Pátli Dún, also divided into two parts, but by a depression, not a ridge, a fact possibly pointing to its being a valley of denudation. That to the west, which is drained by the Sona nadi, has a slope eastwards parallel to the Himálaya, and is known as the Kotri Dún. That to the east, which is drained by the Rám-ganga and has a slope to the west in the same direction, is called the Pátli Dún. The two rivers meet just before their waters take a bend to the south at an elevation of about 950 feet² above the level of the sea. A few miles eastward of their confluence, the valley of the Pátli Dún shows an elevation of 1,200 feet on the right bank of the stream, and thus allows a considerable fall to the Rám-ganga in a comparatively short horizontal distance. The peaks of the Siwálíks to the south along this entire line seldom rise above 2,500 feet, whilst the plains at their southern base average between eight and nine hundred feet. Further east comes the small Kota Dún between

result of borings made by him in various parts of the tract along the foot of the hills. At Káshipur, in a spot some twenty feet below the surface of the red clay, he obtained the following results:—

2' 3" superficial red clay : 1' 3" green sandy clay, water : 4' 6" black clay, extremely tenacious : 1' 6" light blue sand and abundant water. At Haldna, seven miles north, nearly similar results were obtained :—5' 6" a ferruginous sandy clay or loam, latterly becoming more stiff : 1' 6" a greenish clay, becoming blackish below : 2' a bluish-grey clay, partially sandy, not so tenacious, and quite moist. At Jaspur, nine miles north-west of Káshipur, he found :—5' surface sand, which gradually changed to a stiff red clay : 2' a red, loose sand, damp : 2' variegated sand and clay, spotted : 3' yellowish sand changing to light grey : twigs and roots were found at nine feet and water at twelve feet. At Afzalgarh 6' superficial loam with small nests of imperfectly formed lignite : 2' 6" red sand and a quicksand : 4' blackish stiff clay, and 8' 6" stiff clay, latterly sandy. Two other borings are recorded without mentioning the locality.

¹ Herbert. *l. c.* art 68.

² G. T. S.

Dhikuli and Káli-dhúngi, much broken up by torrents, and having its southern boundary cut through by the various hill streams that cross it at right angles. From Naini Tál to the Sárda the sandstone ridge that represents the Siwálíks is so close to the Himáláya that the geologist alone can discern the connection. The Dúns are here reduced "to deep longitudinal gorges and low gaps, corresponding with a leading geological boundary, that between the old slaty and schistose rocks of the mountains and the massive tertiary sandstones of the Siwálíks." Further east, according to Hodgson, the Dúns are represented by the Saliyán Mári, the Gongtáli Mári, the Chitwán Mári, the Makwánpur Mári, and the Bijayapur Mári, all under Nepál. With the exception, however, of the Dún lying on the road to Kathmándu, none of these have been subjected to scientific examination. "On this track," writes Mr. Medlicott,¹ "very complete representatives are found of the two sub-Himáláyan ranges and their intervening *dún* or *miri*. The Chúringháti range is structurally a facsimile of the original Siwálíks. At the outer base at Bichiyakoh, there are some earthy rusty beds, all greatly crushed. The dip soon settles down to 30° to north-north-west, maintaining the same angle steadily to the top of the pass. This is the typical structure of these detached sub-Himáláyan ranges, the flat inner half of a normal anticlinal flexure. The range is about four miles wide." Throughout the Sikkin and Bhútán *dúns* there is no representative of the Siwálík hills and therefore no *dúns*, but in the Dikrang basin of the Daphla hills, Colonel Godwin-Austen discovered two well-marked ranges of sub-Himáláyan hills with an intervening *dún*. We have now seen that the first characteristic features met with in advancing from the plains to the Himáláya are the Taráí, Bhábar, Siwálík range, and *dúns* or valleys, and that though not continuous and indeed occasionally altogether absent, they are, taking the whole range, characteristic of the relations of the Himáláyan mass with the subjacent plains.

Crossing the Dúns northwards towards the snows, we meet the

Lower Himáláya.

outer ranges of the lower Himáláya. They have a general elevation of about 7,000 feet above the level of the sea, while the highest summits along the line reach between 8,000 and 9,000 feet. This generalisation, though giving the nearest approach to accuracy that the state of our know-

¹ Manual of Geology. p. 543.

ledge permits, is only approximate, for we know very little regarding the outer range between the Sarda and the Brāhmaputra. There is this peculiarity in Kumaon, that the outer range first crossed by the traveller is of considerably higher elevation than the intermediate ranges crossed between it and the outlying spurs of the great snowy range itself. We have said that we hold the entire Himālaya to be but the southern slope of the great Tibetan plateau; that however rugged and furrowed this slope appears, it is homogeneous throughout. We reserve for the district notices the purely geographical description, and will here try to answer the questions that naturally arise regarding these mountains as a whole. What have geologists been able to discover regarding their history and the materials of which they are composed, and are the causes at work sufficient to produce such very varied results as are here exhibited? In attempting¹ to answer these questions, we must refer to well-known principles,² which have been found true in Europe and apply with equal or even greater force to the phenomena observable in the Himālaya. Without trespassing on the domain of the professed geologist, we may briefly summarise the facts observed regarding the Himālaya as a whole. We have seen that the outer range forms a geological as well as a physical boundary under Kumaon. The whole mountain mass may be divided into three great belts. First, the sub-Himālayan tertiary sandstones outside the Himālaya proper of geologists, and which have a considerable development west of the Jumna. To them belong the Siwāliks and the Sirmūr series on which the hill sanatoria of Kasauli, Dagshāi, and Sathu are built. Secondly, a belt of limestone and slate forming the outer range of the lower Himālaya on which the hill sanatoria of Simla, Chakrata, Musuri, Landaur, and Naini Tāl are situated. Thirdly, the crystalline rocks with granitic intrusions that form the remainder of the lower Himālayan region as well as the line of snowy peaks, and across the British frontier to the north of Kumaon, the palæozoic and secondary rocks of Tibet. Thus we have as the materials of the Himālaya the two great classes of rock known as the stratified or bedded and the crystalline. To the

¹ My acknowledgements are due to the works of Mellicott, Blandford, Tyndall, and Hekie, on which the following pages are based. ² Though known to every student, a re-statement of these principles seems necessary in a work intended for popular use in India.

former belong the limestone, sandstones, and slates that have been derived from the waste of the older rocks ; and to the latter the granite, gneiss, and schists which occur in masses.

The core or nucleus of all great mountain masses is formed of crystalline rocks, while the stratified rocks enter largely into the composition of the lower subordinate ranges. This is true of the Himálaya, Alps, Pyrenees, Rocky Mountains, and indeed of all the great mountain systems. A glance at the geological map given hereafter will show more clearly than any description the arrangement of these rocks in Kumaon. One of the fundamental principles of geology is that the sites of all the great mountain masses of the world at one time formed a part of the bottom of the sea. K2, Nanda Devi and Mount Everest, the last of which exceeds a height of 29,000 feet above the level of the sea, at one time apparently formed parts of the bed of an ancient ocean. For the crystalline rocks have arisen either from the gradual consolidation of materials which had been fused deep within the crust of the earth or from the influence of subterranean water combined with the earth's internal heat, out of earlier sediments such as sea-mud and sea-sand, which in the course of time had sunk down and been covered by many thousands of feet of later deposits.¹ Geologists tell us² that while no important movements, except small and partial changes of elevation, can be traced in the peninsular formations of India, the whole of the gigantic forces, to which the contortion and folding of the Himálaya and the other extra-peninsular mountains are due, must have been exercised since eocene times. The sub-Himálayan beds were deposited upon uncontorted palæozoic rocks ; and although a part of the Himálayan area may have then been land, the direction of the ranges is clearly due to post-eocene disturbance. It has also been shown that the movement has been distributed over the tertiary, post-tertiary, and pleistocene periods. It has been suggested that upheaval still goes on, as earthquakes are of common occurrence along the line of the Himálaya and as far westwards as Kábul. Three distinct shocks were felt at Naini Tál in April, 1880, and Srinagar in Garhwál was almost destroyed by a great earthquake in 1803. A recent traveller,

¹ Professor Geikie on 'Mountain architecture.'
of India, &c.

² Manual of the Geology

M. Severtsof, attributes the contraction of the great Kára-kul lake on the western Pámír to a similar cause, and says that the connection of the Tián-shán with the Pámír is due to an upheaval which geologically is of recent date and is still progressing. At the end of the cretaceous period and at the beginning of the tertiary period, the Tián-shán was separated from the Pámír by a strait with rocky islands, the marine deposits of which are found beyond the Tuz-Ashu pass. It is startling to the uninitiated to be told that the mighty mass of the Himálaya, as it now appears, is a formation younger than the comparatively insignificant hills of the Dukhin (Deocan) and Central India. The same, however, is said of the Alps, Pyrenees, Andes, and Rocky Mountains, all of which received their chief upheaval in tertiary times.

According to Mr. Blandford it is probable that the crystalline axis of the western Himálaya which apparently terminates in the Dhauladhár peak, Eocene period. the western extremity of the snowy range seen from Simla, coincides with the shore of the ancient paleozoic continent of which the Indian peninsula formed a portion. If this be a correct view, the cis-Himálayan paleozoic rocks are in a great part of fresh water origin, whilst the marine paleozoic formations are found throughout the extreme north of the Panjáb, Kashmír, and the neighbouring countries north of the Dhauladhár and crystalline axis. In eocene times, the peninsula of India was part of a great continent probably united with Africa. To the east was a great sea extending up the Asám valley along the southern base of the Gáro hills and thence southward throughout a considerable area west of the Irawadi in Barma. There was another sea to the north-west covering a great part, if not the whole, of Persia, Baluchistán, the Indus plain, and extending as far north-east as Garhwál, and an arm of this sea extended up the Indus valley into Ladák. The Himálaya and perhaps Tibet, wholly or in part, were raised above the sea, but there is no evidence to show that they had then attained any unusual elevation. In later eocene times, the Himálaya had risen sufficiently to send back the sea boundary to the north of the Panjáb, and in miocene times, the marine area was still further contracted. The existence of a sea in these places is attested by the presence of marine deposits and shells, and this discovery clearly shows

that not only did the sites in which they occur once form the bottom of a sea but that the difference in height now observed between them and the level of the sea must be greatly added to in order to arrive at the true measure of the upheaval that has since occurred. The work of denudation that continually goes on shows us that the present summits of the mountains must in the course of time have lost a considerable portion of their substance, and in the next place we cannot suppose that the marine shells now found lay exactly at the sea level. The bed of the great sea may also have been subject to successive periods of depression and elevation before the eocene period without greatly altering its height. Professor Geikie¹ tells us what the forces are to which these marvellous results are due:—

“The upheaval of the sea floor into land seems to have been due to a cause which has been going on from the earliest geological times and which is still in progress. It is believed that originally this planet possessed an enormously high temperature; that, indeed, it was thrown off from its parent sun with a temperature probably even much fiercer than that of the sun at present; and that it has since been gradually cooling and contracting. The external crust of the earth, varying greatly in structure and otherwise, has yielded unequally to the strain of contraction. One result of this process has been the elevation of portions here and there into long ridges, forming the continental masses and mountain chains. You may illustrate this production of lines of elevation along a generally subsiding surface by what takes place when an apple dries. Its surface contracts and wrinkles, most of the skin sinking inwards, but, at the same time, inequally and leaving intermediate ridges to stand up. So in the gradual contraction of our planet, wrinkles have arisen on its surface. It is these wrinkles which form our mountain chains. But such a subsidence of the crust could not have taken place without a very great deal of folding of the rocks. Descending nearer to the earth's centre, the various layers of the crust had a less diameter to fill. They could only accommodate themselves to their new position by being crumpled up so as to occupy less space, or by being cracked across so as to allow some parts to be pushed above others.”

We find that both these results have been produced, and the records of the Indian Geological Survey teem with illustrations of them.

One other fact is noticed, by the same writer, in connection with the crumpling up of the mountains, and that is that this process has been the means of bringing up the crystalline rocks. "Before the time of the crumpling, the whole future mountain area was covered with one continuous sheet of marine strata. But as the mountain chain began to form, the central portions came to be more and more compressed, puckered, and crystalline, some parts being squeezed up, whilst intrusive masses and veins of granite and other crystalline rocks were injected amongst the intensely altered strata along the central nucleus or core. It was during this process, doubtless, that the crystallisation of the gneisses and schists took place, when they passed from their original character of fragmentary (bedded) rocks and assumed the peculiar crystalline texture which they now present." We have already noticed that there have been successive upheavals of the Himálayan mass through the tertiary and post-tertiary and even the pleistocene periods, and the effect of these upheavals on the form of the mountain ranges must have been considerable. Supposing, with Professor Geikie, that a whole mass of sedimentary rocks has been upheaved into land as a mountain chain, we find that "on the outskirts of this elevated area, sedimentary deposits will continue to accumulate in the sea. If in the course of the slow secular contraction of the planet the upraised tract subsides, a new set of strata will be laid down upon the upturned edges of the older rocks. It is evident that in every junction of this kind, some considerable interval must have elapsed between the formation of the older series of rocks A (Fig. 4.) and the newer series B.

FIG. 4.



Section of a mountain chain showing three epochs of upheaval (Geikie).

"In the course of time, the region having once yielded to the strain from terrestrial contraction will probably yield again, and a new upheaval will take place. The series B will now be raised up together with A, and another series C will be laid down in turn

upon its edges. Subsequently, the same fate will befall the group C. These three sets of differently inclined strata would fix for us three successive periods of upheaval." This simple explanation shows how very varying must be the results of successive periods of depression and upheaval, and especially when, as in the case of the *Himálaya*, such an immense area has been subject to disturbance. We have now seen that the primary factor in mountain architecture are the great changes in the earth's crust by which mountains have been formed, and the bedded deposits have become rocks, and eventually, as the process of upheaval went on, have been crumpled, folded, crystallised, and fractured. In this process, lateral pressure has been the chief agent, and this has been exerted simultaneously from different sides in the case of the *Himálaya*, at least in the pleistocene period. No better examples can be given than those enumerated by Mr. Blandford as characteristic of the western area. Here we have amongst the mountain ridges that encircle the Indus plain and comprise *pliocene* beds, "ranges running north and south such as the *Khirthar* and *Sulaimán*; east and west as the *Marí* and *Bhugti* and the *Afridi* hills; north-west and south-east as the *Pir Panjál*; north-east and south-west as the eastern *Salt Range* and *Kharian* hills; and many intermediate directions may also be traced, independently of curved ridges." Of the extent of these lateral thrusts an example will be found to the west of the Indus. Taking the Persian area and that of the *Himálaya* and *Tibet*, "the mountain ranges fall roughly into two great curves convex to the southward; but the deeper western curve has produced the smaller mountain ranges. That a gigantic lateral movement has taken place in the apex of the western curve is, however, shown by the fact that for nearly 150 miles between *Gwádar* and *Jálk* in *Baluchistán*, the track traverses beds, all apparently of tertiary age, at right angles to their strike and that all these beds are vertical or nearly so. The contraction in breadth, or in other words, the lateral movement must have been great to have converted horizontal formations into a series of undulations, with dips so high as those seen in the *Baluchistán* ranges." The terms at our disposal to denote the relations of magnitude and adequately to depict the changes that have been wrought during such a period are utterly insufficient to convey a correct idea of what has taken place. Perhaps

Professor Geikie's simile of the dried apple is as far as we can go without entering into scientific details that belong to the professed geologist.

Only second in importance to subterranean influence in the forma-

Mountain sculpture.

tion of mountains are the sub-aerial tools of the great sculptor, the different forms of water, ice-wedges, glaciers, snow, rain, and rivers.¹ It is impossible to say what may have been the appearance of the mountain ranges when first formed, but we have every reason to believe that usually the process was gradual and that at once the denuding influence of the different forms of water came into play. Nature from the time the first atmosphere existed has ever been at work tracing lines which gradually work into gorges, ravines, and valleys, weathering peaks and rounding ridges and producing

Ice-wedges.

those alterations in the general appearance of the mountains that on a very small scale are familiar to most of us in a neglected hill-station. Both crystalline and bedded rocks abound in joints or divisional planes by which they are separable into blocks and no small part of nature's work in sculpturing mountains is thus rendered possible. Into these crevices runs the melted snow or rain, and there congeals and again expands, forcing the blocks asunder by slow degrees. The sun's rays turn the ice into water during the day, to penetrate still further and again congeal during the night, and this ceaseless process continued for many centuries shows its effect in the form of the mountains composed of even the hardest rocks. These are covered with massive boulders quarried by nature's ice-wedges in this simple fashion. Where the dislocation takes place near the edge of a weathered cliff, the mass of ruin caused by the toppling over of huge blocks is often gigantic. Gerard describes the upper portion of Purgial (Lío Porgyúl) as the "wreck of some towering peak burst asunder by severe frost." Fraser tells us that the summits near Gangotri and Jamnotri are a mere confused mass of huge crumbling boulders, and the same description applies to nearly all the peaks that have been visited by travellers.

We have next to notice snow in the form of avalanches as one of the tools employed in mountain sculpture.

Avalanches.

The winter snow, when exposed to the summer

¹ Acknowledgment is due here to Professor Tyndall's and Professor A. Geikie's mountain architecture.

sun and influenced also by the heat of the earth itself, is often detached in masses sufficient to cause great natural disturbances. It has been suggested that the change in the bed of the Satlaj in the plains to its present one has been produced by an avalanche in its upper course having dammed up the river. In time the barrier gave way and sent down an immense flood to the plains, sufficient to carve out a more direct course which the river itself has since continued to occupy.

It is, however, to the action of snow in the form of glaciers that the more important results are due. These
 Glaciers. great engines of denudation have the form of a solid river ever progressing downwards through the valleys until the point is reached where the rate of motion is balanced by the melting of the ice. In the figure of the Pindari glacier, given hereafter, it will be seen that the glacier fills the bed of the upper valley and is fed by the drainage of the snow-covered slopes on either side. Its face is discoloured with mud and stones and is utterly unlike one's preconceived idea of masses of ice. Along the sides and edges, too, are rows of earth, stones, and boulders transported and deposited in order by the ice-stream. These deposits are called moraines. Much of the material transported falls down into the crevices and gets between the bottom of the glacier and the rocky bed along which it moves, and which is thus subjected to a grinding process that reduces even the hardest rocks to powder. This fact accounts for the turbid character of glacier streams, especially near their source. The influence of glaciers, therefore, is two-fold, firstly in transporting materials and secondly in reducing them to mud or sand. The combined result is often seen in the mass of detritus heaped up towards the end of a glacier called a terminal moraine, and in the striated and smoothed appearance of the rocks that have been subjected to glacial action. As will be seen hereafter, there is sufficient evidence of a great extension of glacial action in former times that must have had a very important influence on the form of the mountains. Cunningham records three great inundations of the Indus due to the bursting of glaciers in the upper portions of its course. These had dammed up the river bed and eventually gave way, sending a flood down the channel which in the cataclysm of June, 1841, appeared as a wall of water some

thirty feet high, destroying every thing that came within its reach.¹

Rain has had even a more constant and penetrating influence on the mountain masses than any of the preceding forms, for it has furnished the materials from which the ice has been formed and is more universal in its operations. The salts and acids contained in it have also had a peculiar action of their own. Rain while falling through the air takes up some portion of carbonic acid and when it reaches a rock dissolves and carries away certain portions of its texture. The result of this process is that not only is the rock reduced in bulk by chemical action but what remains also becomes more easily operated on by the mechanical action of falling water in the next shower. For illustrations of these processes take any line along the limestone ridges about the hill sanitarium already mentioned, and it will be seen how the outer crust where exposed is crumbly to the touch, and has a rough sandy appearance. For those who have seen the long gneissic range extending from Almora to Devi Dhura in Kumaon, there could not be a better example of the influence of rain on rock than is there exhibited. Along the road on each side where the rock has been exposed to the weather, the outer layer is removable by the hand, and at the base will be found a little heap of sand that has been weathered away in course of time. Many of the more loosely formed shales, especially those that contain alum, speedily decompose on exposure to the atmosphere and it is on this account that in the midst of rocky formations in the Himálaya it is so often very difficult to obtain good building stone. Another familiar example of the influence of the rain-fall on the rocks will be seen in the stones of old buildings throughout the hills. Where protected from the weather their surface exhibits the faintest trace of the graver's tool intact, but where exposed they are worn and eaten into and the outer skin appears granulated and rough. This waste of rock material has been in progress for centuries and has produced a soil in which trees have taken root and shed their leaves to produce by decomposition and mixture with the waste the rich vegetable mould that overlies our forest-clad hills. The presence of these trees has had the further effect of retarding the removal of

¹ Ladak, 104 : J. A. S. Ben. X. 617 : XII. 183 : XVII. 230.

the newly formed soil not only by absorbing a portion of the chemical elements carried down by the rain-water, but also by breaking the force with which the rain would otherwise fall on the soft soil. Thus we find that on well wooded hills the depth of useful soil is considerable and that springs are numerous and abundant. On the other hand where the hills have been cleared of forests, the finer soils are soon washed away by the almost tropical rain. The rocks from which the soil has been formed again appear at the surface and the rainfall rapidly drains off leaving no supply for springs, and if the process be continued over any considerable area, cultivation becomes impossible and the climate is essentially altered. What deforestation has done for Almora can be seen in its scanty rain-fall, its barren slopes, and few springs, although the area affected is so small.

The action of a river in the sculpture of mountains is three-fold.

Rivers.

First, it has the chemical action of rain in dissolving portions of rock constituents : again, it has in its mountain course the grinding power of the glacier in the force with which it drives the gravel, stones, and boulders along its rocky bed : and thirdly, it has the glacier function of transporting material and laying them down in deposits elsewhere. In the case of glaciers the denuding process is the more important, and in the case of rivers the transporting function has, perhaps, more influence in moulding the features of the surrounding country. In the beds of many of our mountain streams we can detect the action of both glaciers and rivers in the striated and furrowed appearances produced by the former and the rounded forms of worn pebbles due to the influence of flowing water. The muddy colour of the water is due to mud or sand held in suspension, and it has been estimated that in this way one six-thousandth part of a foot is annually carried away from the water-shed of a great river. This waste is, however, very unequally distributed, being very much greater in slopes and valleys and less in plains. "We may be prepared, therefore," with Professor Geikie, "to find that solely by the continued erosion of running water, even the most recently upheaved mountain chains have had stupendous chasms carved out of their sides, and an almost incredible amount of material removed from their surface." Such has been the origin of the Scottish valleys which, according to the same writer, "have been cut out of the general mass of the upraised rock. The existing mountains are

what we now find them to be, because they have been left standing while the valleys have been excavated among them." Playfair in his "Illustrations of the Huttonian Theory," as quoted by the same author, writes :—" If indeed a river consisted of a single stream without branches, running in a straight valley, it might be supposed that some great concussion, or some powerful torrent, had opened at once the channel by which its waters are conducted to the ocean; but when the usual form of a river is considered, the trunk divided into many branches, and then again subdivided into an infinity of smaller ramifications, it becomes strongly impressed upon the mind that all these channels have been cut by the waters themselves ; that they have been slowly dug out by the washing and erosion of the land; and that it is by the repeated touches of the same instrument that this curious assemblage of lines has been engraved so deeply on the surface of the globe." In major and minor river systems the same principle is observed ; the lines marking the tributaries of a stream appear like the veins of a leaf all converging on the mid-rib and each forming within its own area a separate main line of a smaller system until the differences are inappreciable. But it may be asked why, if these influences are uniform in their action, the results are so varied. The answer is not far to seek and is to be found in the varied character of the materials on which the aërial forces operate. The southern flank of the Siwálíks below Dehra, consisting of soft sandstones, are weathered by the heavy monsoon rains until they are almost perpendicular. The Krol limestones give their picturesque outline to the outer Himálaya, when compared with the other lower ranges. The shales and slates have a character of their own, and the great crystalline range itself owes its form to the rocks of which it is composed. Kamet has its peculiar pyramidal shape from its cap of granite, and Nanda Devi, Trisúl, and the Panch Chúli have had their peaks defined by simple aërial action on their materials. Thus, our mountain ranges are due in the first place to subterranean disturbances, and in the second place, to the action of the different forms of water, chemically and mechanically, on the varied materials of which the rocks are composed. It is the combination of these two forces that gives such varied results, and until more accurate and comprehensive information is recorded regarding their operation, it is impossible to base our physical description on other than arbitrary grounds.

CHAPTER III.

GEOLOGY.¹

CONTENTS.

Position of the ground. Early observers : Herbert, Cautley, Falconer. The mountain system. The sub-Himálayan zone formed of tertiary rocks. Siwálíks. Post-tertiary deposits. Supposed glacial deposits. Lower Himálayan region. Simla region. Sirmur formation. Relation of Sirmúr and Siwálík series. The Satlaj valley. East of Simla. The Kumaon section. The Nepál section. Sikkim section. Slight correspondence between the rocks of the Himálaya and of the Peninsula. Frequent occurrence of carbonaceous deposits throughout the Lower Himálaya. Conjectural affiliation of the Lower Himálayan sections. The Central Himálaya. Strachey. Stolickza. Western Tibet. The principal crystalline areas. Nummulitic deposits. Post-eocene eruptive rocks. Central gneiss. Kashmír-Kishtwár region. Central Tibet. Its granites, schists, slates, conglomerates and fossiliferous rocks. Palæozoic strata, Silurian, Trias, Jurassic. Fossils. Tertiary deposits. So-called Níti fossils. General Strachey's conclusions. Stolickza's views.

THE British Himálayan districts, which form the immediate subject of this sketch, are themselves such a small portion of the immense geological region to which they belong, that we shall have to wander considerably beyond their limits to attain some idea of their place in nature. Some of the ground forms part of one of the best known, and certainly the most widely known, of our Indian rock-formations : the Dehra Dún is pre-eminently the Dún ; and the low hills separating it from the wide Gangetic plains are the original Siwálíks, a name to be found in every geological text book. The rocks of the higher hills to the north, below the snowy range, have as yet received only cursory attention, being chiefly non-fossiliferous slates and crystalline schists. On the snowy range and beyond it in Chinese territory we again come upon formations of well-established position, but of which we have little real knowledge, and are altogether dependent upon the occasional observations of a few adventurous explorers.

In geology, no less than in other sciences, it is desirable to be able to trace the stages of knowledge. Even in descriptive geology this information is interesting ; and for the student such illustrations are almost essential. Although the germinal idea of geology, that the aboriginal

¹ This chapter has been written by Mr. H. B. Medlicott, Superintendent of the Geological Survey of India, for this volume.—E. T. A.

superposition of sedimentary deposits is a sure indication of succession in time of formation, and hence that the structural relations of rocks are the ultimate criterion of age, had to be conceived before geology could have birth. The difficulty of applying this test, of observing obscure and scattered outcrops, and of putting together and discussing the features thus laboriously collected, is so great, that, from the beginning, geologists have sought for, and adopted more ready tests for the chronological classification of rocks. The history of progress in geology is in great measure made up of the failures of generalisations thus too hastily arrived at ; the total breakdown of false assumptions, and the correction of errors due to the forced application of partially understood principles ; to the neglect of the regulating laws of structure. The science is so young that its history in India affords examples of these errors. In some cases our admiration of the men and the work they accomplished is positively enhanced by our knowledge of the difficulties under which they laboured. The names Herbert, Falconer, Cautley, and Strachey call especially for mention in connection with the ground under notice. Cautley and Falconer will be imperishably associated with the palæontological branch of geology, as having with great labour brought together the unrivalled collection of fossils, the description of which was partially published in the *Fauna Antiqua Sivalensis*. Work of this kind endures, in so far as it is to a great extent a record of hard facts, having each a permanent interest, such as the existence of a certain fossil within a fixed range of strata. It is not so with facts of the first order in geology proper : the announcement that such a kind of stone occurs in any particular place conveys no information that can be said to have scientific value ; it is only when accumulated and colligated under established principles of formation that such petrographical facts come to have any geological meaning. Herbert's observations were of this order. He dealt entirely with unfossiliferous rocks, and the principles under which he had to arrange those observations were still to a great extent artificial ; his work has therefore only an historical interest.

Captain G. D. Herbert, however, must rank in merit as well as by date amongst the foremost pioneers of geology in India. As a man of great talent and of sound and extensive scientific culture, he may stand with Captain

Herbert.

Newbold, who did so much for the geology of Southern India. The advantage of the latter lay in the greater simplicity of the ground he worked in and in his being even a few years later in the field. The mineralogical survey of the Himálayan districts was one of the earliest attempts at a geological map of a considerable area made officially in India. The work was entrusted to Captain Herbert by the Marquis of Hastings; but it was left to private enterprise to make known the results. The publication was taken up by Mr. Henry Torrens, of the Bengal Civil Service, the accomplished editor and proprietor of the so-called *Journal of the Asiatic Society of Bengal*, which was then brought out at the personal risk and responsibility of the Society's Secretary, and was really the continuation of the publication started by Captain Herbert himself in 1829 with the more appropriate title (under such conditions) of *Gleanings in Science*. The report appeared as an extra number of volume XI. of the *Journal* for 1842, nine years after the death of Captain Herbert, and seventeen years after the completion of the survey. The map to illustrate the report was issued with volume XIII. for 1844. It comprises the very large area lying between the river Káli and Satlaj, more than 200 miles in length, and from the plains to beyond the snowy peaks, a breadth of 90 miles. Captain Herbert does not assume any pretensions to authority. He tells us very plainly that he made up his geology for

the occasion, but it is plain, too, from his observations and reflections, that he thoroughly mastered his authors. His suggestions in correction of current views are often very judicious, and display a truly scientific turn of mind. His work, nevertheless, can only be noticed in illustration of the history of Indian geology. He divides all the rocks of the mountains into two great 'primary' formations—one for the gneiss occupying the central region, and one for the micaceous, chloritic, hornblendic, and argillaceous schists, to which also he joins the limestones. He makes a third zone of the narrow strip of "secondary rocks, mostly, if not entirely, the Newer Red, or Saliferous Sandstone." On the strength of this purely imaginary identification borings were recommended, if not actually undertaken, along the margin of the plains, to find the carboniferous formation with its coal. This notion was not quite exploded by the discovery of the famous tertiary fauna in a part of the rocks designated by Herbert as New Red Sandstone ;

and by the latest writers, prior to the Geological Survey, the sandstone along the fringe of the Kumaon mountains, and now known as the Náhan or lower Siwálik group, are treated as secondary. The constant dip of the rocks of the southern Himálaya towards the central axis, so marked a feature in their structure, was treated by Herbert in a manner characteristic of the times. His three formations being by assumption successive in order of time, the observed structure seemed to subvert this ordained relation, making the younger apparently pass beneath the older, the schists beneath the gneiss, and the sandstones beneath the schists. He attempted first to explain this anomalous feature by faulting; but when his calculations seemed to demand a fault having a throw of eight miles, he gave up the idea in favour of a supposition infinitely more extravagant. He came to the conclusion that the apparent bedding in each of his three series is not true stratification, due to the process of deposition in water, but only pseudo-stratification, produced by some process of concretionary action; thus, for the sake of a collateral issue, he cut away the very foundations of the science of geology.

The work of Cautley and Falconer was the geological converse of that attempted by Herbert. They dealt entirely with one series of rocks, and treated them almost exclusively from the point of view of their fossil contents. The range of their operations was limited to the low fringing hills between the Ganges and the Satlaj. The structural features were very slightly touched upon, the strata being taken to belong to one unbroken formation, which was recognised as of geologically recent date, and as being distinctly made up of the débris of the Himálaya. The following abstract list of fossils will give an idea of the extent of their labours, and of the richness of the fauna they discovered:—

SIWALIK FOSSIL VERTEBRATES.

QUADRUMANA.

Scinnopithecus, Pithecus.

CARNIVORA.

Ilyana, Amphicyon, Ilyænarctos, Canis, Mellivora, Felis, Drepanodon, Lutra, Enhydriodon.

PROBOSCIDA.

Stegodon (4 sp.), *Loxodon* (1 sp.), *Euelephas*, *Mastodon* (2 sp.), *Dinotherium* (sp.).

ARTIODACTYLA.

Hexaprotodon, *Merycopotamus*, *Hippohyus*, *Sus*, *Chalicotherium*, *Sivatherium*, *Antelope*, *Cervus* (sp.), *Camelopardalis*, *Camelus*, *Bos* (sp.), *Hemibos*, *Amphibos*.

PERISSODACTYLA.

Rhinoceros (3 sp.), *Antoetherium*, *Equus*, *Hippotherium*.

REPTILIA.

Crocodylus, *Leptorhynchus*, *Varanus*, *Colossuchelys*, *Emys*, *Trionyx*, *Testudo*.

Undetermined species of *Aves*, *Pisces*, *Mollusca* and *Crustacea*.

It is to Captain Richard Strachey, of the Bengal Engineers, now General Strachey, Member of the India Council, that we owe the first sound attempt at a sketch of Himálayan geology based upon extensive observation.

An abstract of his results was published in the Quarterly Journal of the Geological Society of London for November, 1851, and the map accompanying his paper includes about the same ground as that of Herbert. A comparison of the two maps shows how great an advance had been made. It is greatly to be regretted that official and other business prevents General Strachey from making with his own hand the few corrections and additions necessary to bring his work up to date for the present publication. The annexed map is little more than a reproduction of General Strachey's, and the description also of a large portion of the ground is a reprint of his paper.¹ Acknowledgment is made by him that a considerable portion of the observations recorded were contributed by his brother, now Sir John Strachey, late Lieutenant-Governor of these provinces.

For the geographer a mountain chain is fixed by the conditions of continuity and direction. The geologist would fain make his mountain systems to

¹ On the Geology of Part of the Himálaya Mountains and Tibet : Quar. Jour., Geol. Soc., London, for November, 1851. Vol. VII.—E. T. A.

depend primarily upon contemporaneity of formation, as suggesting causal connection. From neither point of view can the Himálaya pretend to fixity of limits. On the south, indeed, no great physical boundary could be more marked than the base of the mountains between the Jhelum on the west and the Bramaputra on the east, a distance of about 1,500 miles. The line is remarkably unbroken, without projecting spurs or open re-entering valleys. It has a uniform curve, the bearing in Upper Asám being to east by north, while in the Panjáb it is nearly to north-west. Throughout this entire length the hills rise abruptly from the alluvial plains. The terminations of this boundary are also very well marked. The Asám valley ends against the ranges of the Barma-Malayan mountains, which pass up from the south, at right angles to the Himálaya chain, to coalesce with it in the elevated regions of Eastern Tibet. Similarly on the north-west the ranges of the Sulimán and Hindu Kush pass continuously into the mountain region of Little Tibet, nearly at right angles to the run of the north-west Himálaya. It is on the north that the question of a physical boundary is to a great extent arbitrary. The chain of mountains that, under the name of Himálaya, forms the northern boundary of Hindustan, is in reality the southern face of the great mass of elevated land extending through nearly 30 degrees of longitude, from the sources of the Oxus to those of the great rivers of China, while its northern face appears upon our maps as the range called Kuenlun. To the south lie the plains of India, whose greatest elevation is about 1,000 feet above the sea; while on the north is the Central Asiatic desert, which nowhere is at a less altitude than 3,000 feet. The loftiest summits known on the surface of the earth are to be found towards the southern edge of this elevated region, at least one peak having been measured whose height is upwards of 29,000 feet, while along the whole line peaks of 20,000 feet abound.

So little is known of the interior and northern parts of this region that it is impossible to offer any general account of it based upon actual observation; but as far as we may judge from those parts that have been explored, it appears that the surface is, with few exceptions, broken up into a mass of mountains, the general elevation of which, valleys as well as ridges, is very great; and there seems no reason for supposing

Tibetan plateau.

p special
 ted mass
 ortion of
 particu-
 bout the
 ough the
 order of
 e plains,
 and the

the most
 ndered at
 he origin
 -basin is
 posits is
 e opinion
 The only
 so of salt-
 sea is the
 n on dry
 disagree-
 nly to do
 The only
 sh water
 esent sea

r of these
 lue to the
 Torrents,
 umulated
 ountains
 s the *bhá-*
 it is about
 per mile,
 it. Even
 s porous
 the *turái*,
 abundant,

that either the Himálayan or Kuenluen have any definite special existence as mountain ranges apart from the general elevated mass of which they appear to be the two opposite faces. The portion of the southern chain to which the following description more particularly refers, is somewhat to the west of a central line, on about the 80th degree of east longitude, which meridian passes through the island of Ceylon, and not far from Cape Comorin. The order of notice will be in successive zones from south to north—the plains, the Sub-Himálayan zone, the Outer or Lower Himálaya, and the Central Himálaya.

While difference of opinion exists as to the formation of the most recent deposits, it is not to be wondered at that there is often much doubt as to the origin of ancient formations. The primitive idea that a water-basin is necessary to the accumulation of extensive sedimentary deposits is still widely held. To it is primarily due the not yet obsolete opinion that the plains of India are in great part of marine origin. The only direct evidence in support of this view is the local occurrence of salt-wells, which also involves a popular error, for although the sea is the great receptacle of salt, it is chiefly by rock-decomposition on dry land that salts originate. This fact is likely to force itself disagreeably upon future generations in India ; at present we have only to do with it as at least weakening a mistaken geological position. The only fossil remains found in these plains-deposits are of land or fresh water origin, and these occur even in the delta below the present sea level.

The surface-form, and the distribution of the materials of these deposits, is, moreover, just what is now understood to be due to the normal action of rivers above their final point of discharge. Torrents, streams and rain-scour from the precipitous slopes have accumulated a wide bank of coarse diluvial deposits along the base of the mountains

which is known in these provinces as the *bhábbar* or forest zone. In Rohilkhand it is about ten miles wide, and has a fall of from fifty to seventeen feet per mile, and, except in the rainy season, water is not procurable in it. Even considerable streams sink into the porous gravel-beds. Outside the *bhábbar* is the *tardi*, also about ten miles wide, in which water is superabundant,

producing swamps and excessive moisture. Formerly it was supposed that this was an area of actual depression; it has, however, a very considerable fall, averaging (in Rohilkhand) more than ten feet in a mile. The moisture is due to the copious re-appearance of the water absorbed at the head of the *bhābar*. From the *tardāi* the plains gradually decrease in slope to three or four inches per mile in the deltaic regions.

A belt of fringing ridges, varying in width and abruptly lower than the contiguous mass of the *Himālaya*, occurs throughout their entire length, with perhaps two short interruptions in Lower *Asām*, which are doubtfully and in part attributed to denudation. The pure geographer might, in some places, ignore this feature, as perhaps below Naini Tāl and Darjeeling, choosing to regard its representative there as mere spurs of the mountains. Geological observation, however, draws attention to geographical features that might otherwise escape notice, and notes that these so-called spurs will always be found affecting the form, not of spurs proper, but of ridges parallel to the adjoining mountains, and in a more or less marked degree semi-detached from them by a chain of deep longitudinal gorges and low gaps, corresponding with a leading geological boundary, that between the old slaty and schistose rocks of the mountains and the massive tertiary sandstones of these fringing hills.

From this double consideration the name Sub-Himālaya has been given to these lower ranges and to the rock-series forming them. For the most part these hills are apparent to the least observant traveller, their outer ranges being separated from the inner ones by broad flat valleys, or *dūns*. These more detached ridges have long been familiarly called Siwālīks, a name extended by Cautley from the native name (Shib-wala) of the representative range separating the Dehra Dūn from the plains. This name, too, has been hitherto currently applied to the hills and rocks here described as Sub-Himālayan, wherever distinctly recognised. The closer study of this zone has, however, brought to light distinctions involving some difficulties regarding the application of this familiar name. It has been found that there are recognisable divisions in this great series

The Sub-Himālayan or Siwālīk zone, formed of tertiary rocks.

Divisions of the tertiary series.

of the tertiary deposits of the Sub-Himalayan hills, and that the younger of these groups contains by far the largest share (if not all) of the well known Siwálik fossils. On this account it was proposed to restrict the name Siwálik to the band in which those fossils occur; but there are good reasons for preferring to continue the extension of the old name, and to indicate the separable geological horizons as Upper, Middle, and Lower Siwálíks. On the small map annexed they are coloured together as upper tertiary.

It was in the ground west of the Jumna, between the Kayarda and Pinjor díns, where the outer and inner hills are confluent for a length of about forty miles, that the separation of the original Siwálik series into distinct groups was first brought to notice. Throughout the whole cross-section the dip of the strata is inwards, towards the mountains; but along a sharply marked line, continuous through the length of these hills, there is a junction of highly contrasting rocks: the brown and yellow clays, and conglomeratic gravels at the top of the series forming the outer zones of low hills, abut against harder red clays and sandstones forming an inner zone of somewhat higher hills. The feature is nowhere better seen than in the region of the Márkanda, south of the town of Náhan, the capital of Sirmúr. As is generally the case along the junction of rocks of very different textures, the actual surface of contact is concealed by débris; and the appearance suggested by the conformable dips is that the outer rocks pass regularly beneath the inner ones.

Such was the view tacitly adopted by the discoverers of the *Fauna Sivalensi*. It was from the same ^{Overlooked by early observers.} trans-Jumna region, south of Náhan and of this rock-junction, that a very large proportion of the great Siwálik fossils were procured. No particular notice was taken of the striking feature just described; and Cautley accounted for the absence to the east of the Jumna of the highly fossiliferous beds known to the west, by supposing a lesser upheavement of the ground to have occurred in that direction; thus distinctly implying that those beds underlie and are older than the rocks of the cis-Jumna Siwálik range, which he had himself,

from fossil and petrological evidence, identified with the rocks at Nahan.

Herbert's deliberate rejection of the elementary fact of stratification, to make way for a theoretical difficulty, is scarcely a less remarkable datum than the ignoring of so striking a stratigraphical feature by the original explorers of the Siwaliks. The facies of the two contrasting groups, as seen along the boundary, at once suggest that the outer and apparently underlying rocks are really the younger, and this is immediately confirmed by finding that the conglomerates of this group are principally made up of the débris of the contiguous inner strata. Although this latter fact would be somewhat against the supposition, the steep abutting rock-junction would next suggest a fault, along which the lower strata were upheaved on the north, and thus brought into contact with younger beds of the same series. Patient search, however, revealed an exposed section of the actual contact showing the relation of the strata to be quite different from that implied by faulting. The conglomerates were found to rest against a denuded surface of the older group of rocks; the junction being in fact that of an original steep edge of deposition to which an actually overhanging, inverted pitch had been given at many points by subsequent lateral compression.

The feature, as thus described, involves much more than simple successive deposition of the groups. It exhibits strong unconformity between them, requiring the older group to have been upraised and deeply denuded before and during the formation of the younger one. Such a relation generally implies a considerable break and lapse of time between the formations, with a corresponding change in their fossil fauna. It is precisely the history of such changes which it is the business of geology to unfold. But to the unfortunate neglect of the simplest stratigraphical observations, fossils from both groups were mixed together in the magnificent collection that lay ready to the hand of the early discoverers. On account of the character and extent of this stratigraphical feature, suggesting that an unconformity of such depth must have a very wide range, it was proposed by the first observer of it to restrict the name Siwalik to the younger formation, and to designate

the older rocks, of the inner zone, as the Náhan group. It is the name by which Cautley identified a certain horizon in the series, although apparently assigning to it a position the reverse of the correct one.

Large as is the gap absolutely required between the ages of the beds actually in contact along this boundary, it was matter of surprise from the first that
Corrected view.
no trace of so great an unconformity could be found in the immense thickness of deposits to the south of it. From the conglomerates at the junction, southwards to the plains, one crosses a descending section of several thousands of feet of strata without a trace of unconformity: showing either that even at the base of this section the beds of the Náhan group are not represented, or that the disturbance which produced the unconformity along this line of abrupt contact was of such a nature as to admit of continuous deposition within so very short a distance. Ruling ideas at the time were certainly against the latter supposition, that extended observation seems to confirm it. Going westwards along the Pinjor Dún, we find at the Satlaj, on the very strike and extension of the Náhan range, a continuous conformable sequence from the beds of the Náhan horizon into the softer sandstones, clays and conglomerates at the top of the series. The line of disturbance which in the Náhan region resulted in a denuded scarp against which the topmost beds were deposited by overlap, produced in the Satlaj region an anticlinal flexure which must have been so gradually evolved that the deposits accumulating along its southern base were sensibly conformable throughout, although now we find the uppermost conglomerates almost vertical, with a southerly underlie, at the edge of the Dún. Thus it is evident that a well-defined break is not a general feature in the Sub-Himálayan rock-series, and that it would be premature so far to sever such a portion of it by a separate name from the time-honoured Siwálíks. The name Náhan is already current in print, and may at present be understood to indicate lower Siwálík rocks. It is important, however, to point out that the real inference from the unconformable junction is of more interest than the *prima facie* one; without it we should not have had distinct proof that slow disturbance of great amount took place in the Sub-Himálayan zone during the formation of the Siwálík deposits.

Although the marked separation of groups suggested by the peculiar feature of the hills between the Kayarda and Pinjor duns

A division maintained. is not maintainable westwards, there is a uniformity of change throughout the series from the rocks and fossils : upper Siwálíks. base to top whereby approximate horizons are

assignable. Conglomerates and gravels prevail at the top, variably associated with brown sandy clays. In many clear sections the thickness is quite 4,000 to 5,000 feet. It is not, however, to be understood that the deposits were ever strictly superimposed to that depth vertically. The mode of deposition in successive banks, each trailing upon and thinning out beyond its predecessor, as pointed out above for the section on the Satlaj, must greatly modify the familiar meaning of the word *thickness* as regards *space*; although where such deposits become tilted up by lateral pressure, and exposed along a comparatively shallow section, the appearance is quite the same as if vertical superposition of the whole series had obtained. Nevertheless, as regards *time*, the fullest thickness must be taken into account, for each bed is truly successive to that below it. Even when raised to the vertical, those upper Siwálík strata have so fresh an appearance as to be scarcely distinguishable from the most recent deposits—from the beds of the torrent shingle or of sandy alluvium now accumulating in the duns or on the plains. The complete justification for their distinction as an upper Siwálík group is found in the few fossils they have yielded, some of which, as *Bubalus palaindicus*, would connect them with the pleistocene deposits of the Narbada valley rather than with the pliocene Siwálíks.

The main fossiliferous zone of the Siwálík series constantly occurs beneath the thick mass of deposits
Middle Siwálíks. noticed in the last paragraph, and it has a fairly characteristic rock-facies of its own. Massive, clear, gray; soft-sandstone is decidedly the prevailing rock; but brightly tinted clays are also often in great force. The large vertebrate remains, although mostly found in sandstone, are certainly more abundant where there are associated clays. Several thousand feet of thickness must also be assigned to this middle-Siwálík group. Falconer considered this Siwálík fauna to be miocene, but palaeontologists are now decidedly in favour of its pliocene affinities.

In conformable sequence beneath the fossiliferous zone we find rocks of the same type, but having a decidedly different aspect—strong sandstones, but of a darker hue, and often highly indurated, with hard clays generally of a deep red or purple colour. Throughout the Himalayan range, east of the Satlaj, they form the flanking ridges close under the higher mountains, and inside the duns; or at least they mostly occupy that position, for it cannot be said that middle Siwálik beds do not occur there too, as will be seen from the remarks upon structure. No fossils can be quoted from those beds, but it is believed that some existed in the original Siwálik collections. They would probably be of miocene age.

One of the most interesting features in these Siwálik deposits is the variation they exhibit in relation to the position of the great river-gorges. This is most marked in the case of the upper portion of the series. The accumulation of coarse conglomerates is immensely greater in the immediate vicinity of the large rivers of the Himalayan system, and, moreover, it is only within the range of those streams that we find the beds of large rounded blocks of quartzite and other hard rocks such as are now brought down by those great torrents. In the intervals between the rivers such conglomerates as occur are formed almost exclusively of the debris of the adjoining hills, the same as are found in the minor streams now flowing from those hills; but in this position sandy clays often form the bulk of the formation. The same influence is observable in the middle group of the series, which is often conglomeratic and gravelly, or almost exclusively sandy, near the main rivers, while away from them the clays are often in great force. These facts are very observable at the Satlaj; in the gorge above Bubhor the whole of the upper group is coarsely conglomeratic, and the middle one is more or less pebbly throughout, while at seven miles to the north-west the brown sandy clays, in which the fossil *bubalus* and *camelus* were found, form three-fourths of the entire thickness of the upper group. The apparent exception in the case of the Jumna is even a more marked illustration of the fact under notice. The river now flows through the Siwálik range at many miles to the west of the gorge where it leaves the mountains; the fact being that in the elevation of the outer range the

Jumna could not erode a passage through the great accumulation of conglomerates it had formerly discharged in front of that gorge, and which now form the highest summit of the outer range. It had to work round them not in the direction of its near neighbour the Ganges, but towards the region of lesser deposition. Thus the contrast between the Siwálik strata to the east and west of the actual river-passage is so great that it has been questioned if the formations can really be the same. To the east the upper group is made of the hard shingle conglomerates, while on the same strike to the west the conglomerates are composed of local, principally lower Siwálik, *débris*. In the *cis*-Jumna Siwáliks the middle group is formed of thick masses of soft sandstones that have yielded very few fossils, while to the west clays occur largely on the same horizon, and fossils abound. The facts indicated in this paragraph are of great importance, as bearing upon the question of the mountain-formation; showing, as they clearly do, that although these deposits, to a thickness of 8,000 to 10,000 feet, are now in many places turned up to the vertical, and even inverted, yet the main features of the higher mountains must have been during the Siwálik period sensibly similar to what they are now.

So far we have briefly considered the original characters of the Siwalik strata: it is necessary now to notice the features induced by disturbance. This has taken place on the grandest scale. On the right bank of the Ganges above Hardwár the gray sandstones of the middle group have a high southerly dip; and this rises gradually, through an enormous thickness of strata, to a nearly vertical underlie in the conglomerates at the outer edge of the range. A section of the same type is splendidly exposed in the gorge of the Satlaj above Bubhor, in the second range of the Sub-Himálayan hills. There is much method in the form of these flexures: they very generally affect the form known as *normal*, i. e., bends in which the dip is greater on one side of the axis, and so called because of more common occurrence than the symmetrical flexure—when both dips are equal—or than the folded flexure, in which the strata on the side of the steeper dip have been pushed beyond the vertical, and so partially inverted. As an almost universal rule in this region, the

steep side of these normal anticlinal flexures is turned from the mountains. From this there results the familiar conformation of the Sub-Himálayan hills, presenting a scarped face to the plains and a long slope towards the interior valley. These dűns, or at least the flat longitudinal valleys which are the typical dűns, are thus structural features, not mere valleys of denudation ; they rest upon the comparatively little disturbed strata in the hollow of the syndinal flexure. The range separating the dűn from the plains is formed by the anticlinal, the steep (outer) limb of which is generally broken up and denuded away, hence the south face of the range presents the scarped outcrop of the beds on the north side of the axis of flexure.

In the inner ranges, where the disturbing action was greater, the normal flexure often becomes folded, with, of course, inversion of the strata. There is an instance of this fairly seen in the Nűn stream under Mussooree : below the narrow gorge, through massive sandstones having a steep northerly underlie, there is a continuous section in the low banks showing the sandstone becoming pebbly, then interbedded with thin conglomerates, then with thicker and coarser beds, all having the same high northerly dip. This is undoubtedly an ascending section though apparently, according to the dip, it is a descending one, *i.e.*, the whole series is inverted. To any one who has understood these simple explanations, it will be apparent that if these conglomerates are upper Siwálik, and unless there is a fault somewhere about the mouth of the Nűn gorge, the inner Sub-Himálayan range under Mussooree must be in great part made up of middle Siwálik, and not of Náhan beds ; and indeed their character would support that view. If these conglomerates of the Nűn are not upper Siwálik they would form a new sub-division of the Náhan group, which could then hardly be classed as lower Siwálik. Similar conglomerates have lately been observed in this inner Sub-Himálayan zone east of Naini Tál. It is however to be remarked that these folded flexures often are attended by great faulting, on such a scale that all appearance of flexure is lost and we only find a section of upper Siwálik dipping against, and apparently passing under, beds of the Náhan type. There are several grand instances of such faults in the broad area of Sub-Himálayan rocks beyond the Satlaj in the Kángra

region and on through the Jamu hills. They run quite straight or in very open curves for several scores of miles, and as the dislocation lessens, the unfaulted flexure is gradually disclosed. The section in the Nún shows us that to the east as well as to the west flexures may take the place of the peculiar unconformable overlapping boundary between the upper and lower Siwálik described in the Náhan region, where there is no dún. The difference of structure would be such as might result from the presence of some unyielding mass of rock underground in this latter position, preventing the formation of flexures. This Náhan region is in other ways remarkable, as we shall see in following sections.

The form of the structural features prevailing in the Sub-Himálayan zone indicates plainly a thrust from the adjoining mountain mass ; and the magnitude of the total movement is astonishing in connection with the reflection made in a previous section, that it must in great part have occurred since the mountain mass had assumed somewhat of its present form ; or, at least, that the main drainage system had remained the same throughout.

From what has already been said, a general idea might be formed of the distribution of the different groups of the Siwálik series ; but the effects of denudation and other influences remain to be indicated. The chief expanse of the Sub-Himálayan hills and rocks is beyond the Satlaj. From about Pinjor, the lower Himálayan ranges trend northwards towards the lofty gneissic ridge of the Dhauladhár, overlooking the Kángra Dún. The outer range of the Sub-Himálayan is not affected by this change of direction, so that the zone of tertiary rocks becomes permanently widened to about three times the breadth it exhibits along the whole mountain range to the east. In the Kángra region there is a succession of three dúns occupied by conglomerates, separated by ridges of sandstone brought up along great faults. To as far as the Ganges the upper Siwálik are still well represented in the outer hills ; but east of this river, along the whole of Rohilkhand, only remnants of these outer Siwálik have been observed. They seem to have been mostly denuded away, and one comes at once upon the older sandstones of the ranges immediately flanking the high mountains. This is the case under Naini Tál. The change, however, is not a permanent one ; for, again, far to the east, on the road to

Káthmándu, there is a broad outer Siwálik range, formed of soft sandstones and conglomerates, and separated by a dún from an inner zone of lower sandstones. But, again, along the Sikkim and Bhután border, there is no sign of the outer Sub-Himálayan range.

It has been said above that the uppermost Siwálik strata are considered to be of newer pliocene age, and Post-tertiary deposits. we have seen modern deposits being laid down against them along the base of the hills in the *bhábar* region. In these a fossil village was dug out by Colonel Cantley in the excavations for the Eastern Jumna Canal. They no doubt passed down into beds of the *prehistorical* or *recent* period. Still there would be a great gap left in the sequence of formations—the whole of the pleistocene period, represented in Europe by the drift, and the cave deposits, which for years past attracted so much attention in connection with the discovery of human remains. For some middle portion of that period representatives are found in the Sub-Himálayan zone. They are unmistakably exposed on the Satlaj, above Bubhor, where the hills on either side of the river, to a height of some 500 feet, are capped by clays and coarse conglomerates, resting quite undisturbed on the edges of vertical Siwálik strata, both of the middle and uppermost groups. It is clear that a long time of disturbance and denudation must have intervened between the deposition of those totally unconformable deposits; also that those high-level conglomerates are separated from recent deposits by at least the time it has taken the Satlaj to excavate its gorge to its present depth. These two limiting tests of age are quite as cogent as those applied to corresponding deposits in Europe. Beds of the same age are well exposed above the sanction already described in the Nún river under Mussooree. They are here quite on a level with the summits of the Siwálik hills to south of the Dehra Dún.

The most interesting deposits of this age are those to which a Supposed glacial deposits. glacial origin has been assigned. It was long since observed that the glaciers of the Himálaya had once extended to a much lower level than they do now. Unmistakable moraines are found in Sikkim to within 8,000 feet of the sea level, the present limit of glaciers being about 14,000

feet. More recently a glacial deposit has been described in the Kángra valley. Enormous blocks of gneiss are freely scattered in the low ground, at an elevation of 2,500 to 3,000 feet, along the whole base of the Dhauladhár range, resting on the Siwálik strata. One cannot well assign a limit to the mass that may be moved by a rush of water on a considerable slope ; but the distribution of these blocks makes it very difficult to account for them by any action of this kind through the existing gorges ; for they occur along the slope of the flanking ridges, seemingly quite out of reach of any possible sweep of the torrents. They are three possible assumptions to account for their position : by supposing the scarp of the gneiss now forming the mountain-ridge to have once extended a couple of miles in advance of where it is now, and so as to bring those blocks within the range of its talus, but in this case the blocks should be found over the intermediate heights, which is not the case ; or to admit that the blocks were ice-borne, and not by glaciers, for the blocks are not arranged in moraine fashion, but by floating ice in a lake of that period ; or to suppose that the whole valley and the main gorges were formerly choked up with detrital accumulations to such a level as may have brought these lateral positions within the range of the spill from the gorges, ever overtopping the lower flanking hills. This is by no means a gratuitous supposition, for we find those coarse superficial deposits capping the heights above Kángra fort, on the south of the valley, and not derivable from the Siwálik conglomerates on which they rest. The necessary slope of deposition for such materials, from this position to the mountain range, would give elevation enough there for any observed distribution of the great blocks, by simple diluvial action. This is probably the real history of the case ; but the possible influence of ice action to aid the process is not to be lost sight of. When the Himálayan glaciers reached to 8,000 feet lower than at present, ice-work must have been very active in the Dhauladhár, and its effects very marked in the deposits at the base of the range. All the facts suggest that those high-level gravels are of the same age throughout the Himálayan border, and probably of the same age as the former extension of glacial action ; and it is impossible not to notice the near coincidence of this age with that of the glacial period of Europe. There is no independent evidence of changes of level since the age

of these deposits ; and it is doubtful how far mere difference of rainfall could count for the change ; we should thus be driven to entertain the idea of an ice-age.

The confirmation of this physical evidence of a great phenomenon having affected synchronously so large a portion of the earth's surface would be of the highest importance to geological science. Since the abandonment of the primitive idea that all similar rocks were of contemporaneous origin, we have been at a loss for any test of absolute time-horizons. Within continuous land areas some approach to a judgment can be made by closely comparing series of adjoining sections, but for any distant or detached area we have to trust to palæontology for the homotaxis, or comparative classification, of formations. Palæontological homotaxis, however, implies difference as well as correspondence in actual time relations ; and the problem of settling, from fossil evidence only, in which direction the difference should be counted, is an exceedingly complex undertaking. Thus palæontology itself was the chief sufferer by the natural limitation of age-tests for the stratigraphical foundation upon which it was based, and of which it is as yet far from being independent. And as the history of life upon the globe is the object of the highest interest in geology, the check to progress was a very serious one. The occurrence of even one semi-universal phenomenon, leaving such peculiar and well-marked stratigraphical characters as those of an ice-period, would afford an invaluable test whereby to check the direction of growth and distribution of organic forms in all the formations nearly connected therewith.

The change from the Sub-Himálayan hills to the outer region of the mountains is always, as has been shown, a more marked feature than the mere difference of height would suggest. The hills of the Náhan zone range from 3,000 to nearly 5,000 feet, while the summits of the adjoining mountain-ridge vary from 6,000 to 8,000. From this to the great snowy range there lies a tract more than fifty miles wide, of deep valleys and narrow ridges, the average elevation of which would be scarcely over that of the border-zone of mountains. This is the region appropriately known as the lower or outer Himálaya. The main watershed of this broad tract of mountains lies

well to the north of the line of snowy peaks, and the great rivers traverse the lower hills in very tortuous courses. The configuration stamped upon the area by the denudation from rain and rivers gives very little clue to the rock-structure. Although the general strike of the formations is parallel to that of the range, the composition of the strata is too complex, and the cases of local distortion too frequent, to admit of anything like the regularity of feature that has been described in the Sub-Himálayan zone. One character may, however, be noticed as constant throughout the western part of the lower Himálaya to as far east as the Nepál frontier : along the outermost zone there occurs a strong limestone formation, producing ridges of more rugged outline and having a greater elevation than the hills for some distance to the north of it. Our knowledge of this immense stretch of mountains is so fragmentary that no connected account can be given of it. We can only give a sketch of the four sections that have been even cursorily observed, and offer some conjectures as to their connection. The four sections occur in the Simla region, the Kumaon region, Nepál and Sikkim.

The first thing to note of the Simla region is that it constitutes the termination of the Lower Himálaya as characterised above. In describing the Sub-Himálayan zone it was noted how from about Pinjor (on the main road to Simla) the boundary of the mountains trended round to the north for about eighty miles, up to the base of the Dhauladhár ridge, which is exactly on the line of the great snowy range, and is structurally its equivalent. Thus, of course, the continuous broad area of lower mountains is cut off. It is important to notice that this is not a freak of denudation, a great bay worn into the mountains by a tertiary sea : were this the case, we should find the successive formations of the area striking out along that curve and abruptly cut off there. The fact is not so. The strike of the rocks bends regularly with the direction of the boundary, thus showing that the entire feature is an original character of the mountain-structure. We have here, in the extinction of the Outer Himálayan region, the beginning of the north-westerly decrease of the Himálayan elevation. The mountains beyond the Ravi, which, in a superficial sense—as being south of the range of maximum elevation—might be held to represent the Lower Himálaya, do not

reproduce its characters. Kashmir has structurally more relation to the Central Himálaya than to anything south of the snowy range. This fact of diminishing original elevation comes out very clearly in the peculiar distribution of some of the formations that occur in the Simla region. These have been roughly classified as below :—

Sirmur ... { Kasauli.
Dagshái.
Subáthu (nummulitic).

Krol (? triassic).

Infra-Krol.

Blaini.

Infra-Blaini.

Schists and gneiss.

The marked change in the surface configuration from the Sub-Himálayan to the Lower Himálayan hills introduces for the most part a total change of rocks. Below Mussooree and Naini Tál, and throughout the whole range to eastward, one steps at once and for good from the upper tertiary sandstone to the much older slaty rocks. It is not so, however, in the region between the Jumna and the Satlaj. The high ridge on which stand the stations of Kasauli and Dagshái is formed of rocks very similar in character to those of the Náhan zone close by ; but the marked boundary separating them is continuous with that forming to the east the separation of the Náhan from the slates. Indeed even here, under Kasauli, the slates often appear along the boundary beneath the sandstones of the ridge, which have been upheaved upon a basement of their supporting rock. Resting upon the slates in this position we find thick beds of dull brown, gray, and olive indurated clays with bands of limestone, in which there occur abundantly fossils characteristic of the nummulitic period. These beds are well seen about Subáthu. They are overlaid conformably and with alternating transition by red clays with hard purple and gray sandstone, well seen about Dagshái. In the ascending section sandstone prevails to the exclusion of the red clays, as is well seen on the ridge at Kasauli. In these top beds numerous leaves have been found indicating the proximity of an abundant sub-tropical vegetation. The aggregate thickness of this threefold formation (Subáthu, Dagshái,

and Kasanli) may be from 2,000 to 3,000 feet. Collectively it may be known as the Sirmur series, a considerable part of its area occurring within that State.

Here, then, at last we have a formation the horizon of which is

fixed by a well-marked marine fauna. The
 Its distribution.

Subáthu group at the base of this series is certainly cocene ; and considering the perfectly transitional character of the three groups, we may provisionally consider the whole series to be of this age. The study of its position and relations seems to throw much light upon the history of the mountain-system. It forms an almost isolated outlier, caught up on the edge of the mountain-area. Its greatest width, east of Dagahái, is about ten miles. In that direction it stops out along the crest of a ridge at about fifteen miles west of the Jumna ; the mode of termination showing that it was effected simply by greater elevation to the east, and consequent denudation. The only other known occurrence of these rocks within the south Himalayan boundary to eastwards is a small patch of Subáthu beds on the top of the ridge east of the Ganges close to the village of Bon in Garhwál. In the far east, however, the rummulitic deposits at the south base of the Gáro hills have a very striking resemblance to the Subáthu beds ; but it is doubtful if they were ever connected. To the west, at the Satlaj, the outlier is attenuated to a band a few yards wide—a bottom remnant of the Subáthu beds. These rocks again appear in some force in the ridge beyond, but become gradually depressed in that direction, so that before reaching the Biás the most characteristic bottom group has disappeared. The band as represented by the upper groups is well marked, though very narrow, at the Ravi. The hill station of Dharmasála stands upon it.

Some points of interest have been made out regarding the rela-

tions between the Sirmur series and the
 Its unconformity to the older rocks. contiguous older formations. There is very deep unconformity : the Subáthu beds do not rest upon the next youngest rock, which is the great Krol limestone, but upon beds which underlie that limestone. This is fully made out. As a natural concomitant, we find that the surface of contact of the two contrasting formations is a very uneven one. On the ridge at Subáthu there are not more than 50 feet of the typical Subáthu

beds below the red rocks ; while in the valley alongside there must be 600 to 800 feet of this bottom group. These facts suggest proximity to the edge of deposition of the nummulitic sea ; and the succeeding deposits, ending in the plant-beds of the Kasauli zone, tend to confirm the view.

It seems, too, that little or none of the contortion which now affects all the rocks had occurred before the deposition of the Subáthu beds ; for at Subáthu itself a characteristic bottom layer is clearly seen, resting continuously throughout a considerable synclinal flexure upon approximately the same bed of the supporting slates. In agreement with this observation, we now find these rocks to have undergone equal contortion with the slates. The inner boundary of the eocene area is a very broken one ; and outlying shreds of the nummulitic clays are found caught up in folds of the slates, as may be seen on the road north of the Haripur rest-house.

All these facts would tend to prove that although some general elevation of the mountain area, involving deep denudation of the rocks, had occurred here prior to the tertiary period, none of the special disturbance characteristic of the existing mountain-system, and so specially marked in this fringing zone, took place till after the deposition of the eocene rocks. The Sirmur series exhibits more intense and varied disturbance than is at all general in the Náhan group.

The relation of the Sirmur to the Náhan group cannot be so definitely made out, as they are only seen near each other along a single steep line of boundary ; and the question is, as to the nature of this boundary. The argument for the total separation of the groups in this region appears, however, to be pretty conclusive. Although, as has been said, the rocks of the upper Sirmur groups have a strong lithological resemblance to those of the Náhan group—so much so that the type of the Sub-Himálayan deposits may be said to have set in with the eocenes—the facies of the Náhan and Sirmur groups are so distinct, in close proximity, that there is little ground for considering any parts of them, as represented in this Jumna-Satlaj region, to be equivalent. It is only on the supposition of the boundary

between the areas (which is also the chief boundary along the base of the mountains) being a great fault, that the question of correspondence can arise at all. But as this supposition is the *prima facie* one—the one that would be applied from the accepted interpretation of like boundaries in other mountain regions—it is necessary to state the evidence against it.

Throughout the whole range of the Náhan zone in this region

no trace of the very characteristic Subáthi
Probable unconformity. beds has been found, either at the apparent

local base of the section, or as a remnant adhering along the supposed fault-ground. In the former position it might be said we should be more likely to find the top rocks of the older group ; but this too can be answered negatively : the lower we get in the Náhan group, we find clays to occur more frequently, whereas the Kasauli beds are almost exclusively sandstones. It is also to be noted that no remnant of the Náhan rocks has been recognised capping the Kasauli beds, or otherwise, within the eocene area. No conglomerate has been observed in the Sirmur group. The only position from which the original continuity of the groups, with separation by faulting, could be maintained, would be to assert that, as the actual base of the Náhan group has never been seen in this region, the whole eocene group may be buried beneath it in conformable sequence, or otherwise ; and correspondingly, that any trace of the Náhan deposits had been washed away from the present eocene area. So much for the direct evidence. As to the indirect, there is really little, except the fact of abruptness, in the character of the boundary itself to countenance the supposition of a great fault. A straight line drawn from the west end of the eocene area in Sirmur to the small nummulitic outlier east of the Ganges would touch the Siwálík hills south of Dehra ; and along the great bay-like course of the boundary, north of that line, there are many sharp changes of direction, such as are not supposed to be compatible with dislocations of such dimensions as would be required in this case. The great faults in the Sub-Himálayan zone were seen to be remarkably straight. The alternative and most probable supposition is that the eocene area was upraised, and something approaching to the present steep edge of contact eroded out of it, before the deposition of the Náhan rocks.

It is, however, to be mentioned that far to the north-west there is complete transition throughout the tertiary series, from the Subáthu to the top Siwálik. The partial obliteration of the Sirmur series, as forming a sharply defined zone at the base of the mountains, commences at some fifty miles beyond the Ravi about Udampur. The zone is there more than twenty miles broad, and rocks of Siwálik aspect occur within it. Still as a zone of greater upheaval it is traceable to beyond the Púñch; but before reaching the Jhilam it is quite effaced, the whole tertiary series sweeping across it in an anticlinal flexure. These facts do not in the least disturb our conclusions regarding the relation of the lower and upper tertiary series in the Lower Himálayan region. They only form part of the concurrent evidence that towards the middle of the Himálayan system the elevation was greater, and commenced earlier than in the terminal region. The same fact is emphatically shown in the comparative relation of the Sirmurs to the older rocks in these two positions. In this case there can be no doubt of the deep unconformity in the Simla region—a relation of the same kind as that here adopted for the Sirmur-Náhan relation in the same area. But this feature too is quite changed to the north-west: in the great inliers of old limestone that occur within the tertiary area of the Jamu hills, the Subáthu group, with the same characteristic bottom-bed as noticed at Subáthu, is everywhere observed in parallel (conformable) superposition with the old limestone. It is not indeed proven that this rock represents the Krol formation; nevertheless, the contrast of the stratigraphical relation is most striking.

We have again to refer to the Jumna-Satlaj ground for the best sections of the next older rocks to the
 Older rocks of the Simla region. best sections of the next older rocks to the
eocene. The peculiarities of this region come out stronger as we recede in time. Its character in the plains, as the present main watershed of Hindustan, is really its least permanent feature: there is some reason to think that the Jumna once upon a time may have flowed towards the Indus through western Rajputána. It can certainly be affirmed that such a course was within the range of the diluvial conditions that formerly obtained in upper India. The peculiar unconformity found in this position between the upper tertiary rocks is not,

like the plains' watershed, accidental or temporary, but structural. In the well-preserved sections of the Sirmar group we found in this region the only representative of the eocene period within the Southern Himalaya; and again here we find the best preserved remains of the older formations. This ground too has been examined in more detail, so it will serve as a standard of comparison.

The Solan rest-house on the new road to Simla stands between three picturesque mountains of limestone, more or less isolated in each case upon a base of supporting rock. It is a blue gray stone, several hundred feet in thickness. From its position here we are safe in taking it as the youngest group of the series with which it is connected. It has now for some years been spoken of as the Krol limestone. There is often a band of coarse sandstone at the base of it, which seems to vary a good deal in thickness, often at the expense of the limestone. Below this there is well exposed in this neighbourhood a thick band of black carbonaceous slaty shales, which pass down into similar non-carbonaceous flaggy beds, forming the whole base of the mountains in this zone, down to the lowest levels. At a thousand feet or more from the base of the Krol limestone there occurs in those slaty flags a thin band of compact limestone of clear pink, yellow, or gray tints, often accompanied by a bed of conglomerate, and a white quartzite. This band, though a thin one, seems to be very persistent to great distances; it is therefore important as a well-marked horizon. It has been identified on the flanks of the hills under Simla; far up the valley of the Tons, at the crossing of the Simla-Mussooree road; on the Ganges near Tapuban, and again on the hillside north of the Kota Dún in Kumaon. It is called the Blaini group. The Krol group is traceable eastwards almost continuously, and in about the same condition as at the Krol, through Deoban in Jaunsár to Mussooree and Naini Tál.

From certain lithological resemblances to the rocks in Spiti, which he had classified from fossil evidence, ^{*} Conjectured affinities. Dr. Stolickza has conjectured that the Blaini and *infra*-Blaini beds may correspond with members of his Muth and Bhábe series, of upper and lower silurian age; that the Krol group represents his Lilang series, which is triassic; and that the *infra*-Krol may correspond with his Kuling series of the

carboniferous period, in the central Himálaya. But no recognizable fossil has yet been found in these rocks in the lower Himálayan region.

The relation of this series, which forms so continuously the outermost zone of the lower Himálaya, to the metamorphic rocks on the north is very puzzling, yet essential to the explanation of the mountain structure. Some important hints towards it are found in the Simla region. Unless one chances to stumble upon an outcrop of the Blaini group, the flaggy slates of the *infra*-Krol and *infra*-Blaini horizons are indistinguishable. Immediately north of the Krol there is a compressed anticlinal flexure, with elevation to the north of it; so that the slaty rocks occupy the whole ground till we reach the quartzites of Tara Devi and Boileauganj, which dip towards their common synclinal axis at the gap south of Simla: limestone occurs above them on Jatog. These may represent the Krol group. However this may be, there can be no doubt about the identification of the Blaini group on the spur under the Yarrows and under Chota Simla, on the opposite sides of Jako. The thickness of strata above these outcrops, to the top of Jako, would quite carry the section up to the base of the Krol group.

An interesting feature of the section is that the rocks of the Jako and Boileauganj hills are highly metamorphic—mica-schists and garnetiferous hornblende-schists with abundance of vein-quartz—while the flaggy slates above and below the Blaini group, all round the west, north and east base of the hill, are as little metamorphic as on the base of the Krol. This is a crucial instance of a phenomenon that meets us far and wide throughout the Himálaya, the superposition of highly metamorphic upon non-metamorphic strata. The metamorphism is often apparently greater than here—we find gneiss instead of crystalline schists at top—while the case for superposition is less distinct; so that it is possible to doubt the fact of its being a normal ascending section; and accordingly this has always been a chief stumbling-block in the interpretation of the lower Himálayan sections. It is the feature that so fatally puzzled Herbert fifty years ago. We shall have to return to the subject presently, and would only remark here that the very instructive

Typical instance of over-lying metamorphism at Simla.

instance we have just seen occurs in a much frequented position, where many English people pass many idle hours.

The same flaggy slates seem to continue for a long way north of Simla, for the most part with a moderate north-easterly inclination. There are several lines of crush and strain, generally found at the gaps, but apparently unattended by great dislocation, for no new rock appears along these lines. The Blaini group has been identified on the ridge north of Theog bringing in the Krol beds towards Matiani, where there are symptoms of gradual general metamorphism. At Nágkanda this change is very decided, and here those slaty schists seem to pass into the flanks of Hatu, the top of which is formed of massive beds of gneiss, lying nearly horizontally. This rock is the same as the so-called *central gneiss* forming the southern basement of the great snowy range, where its chief characteristic seems to be the prevalence in it of ramifying veins of albite-granite. These observations have suggested that the Krol beds had here overlapped the slate series, and are in original contact with the gneiss.

We have now seen a general section up to the great mountain range. It is taken for the most part along the watershed in a north-east-by-east direction from Simla. The apparent simplicity of it is very encouraging; but we have not far to go to dispel this illusion. It may be noted that no limestone appears on this section beyond Simla. The Sháli mountain, however, only a few miles to the west of Matiani, is made of strong limestone, not unlike the Krol rock, with an accompanying sandstone, and underlaid by flaggy slates like those about Simla. If they are the same, and indeed in any case, their position is somewhat puzzling, for from Sháli they dip eastwards under the schists of Matiani and northwards down to the Satlaj, where the limestone seems to pass under the gneiss of the Jalori ridge.* It is not merely a case of dipping *towards* these metamorphic rocks; the V-shaped outcrop of the junction along the steep sides of the Satlaj gorge points up the valley, and is more or less parallel to the dip of the strata in both rocks. In the Satlaj valley above and below Suni, close to the north of Simla, the confusion of the limestones and slates is indescribable; and there is profuse trappean intrusion: whereas, on the whole section through Simla, from

the plains to Nágkanda, only one small dyke has been observed. The absence of fossils in these limestone and slate rocks makes it almost impossible to settle their stratigraphical relations with any certainty.¹

The distribution of these limestones and slates is as irregular to the east as to the north of Simla. The Chor mountain—directly between Simla and Mussooree, and remarkable for being the highest summit (12,000 feet) occurring within such a short distance of the plains—is an isolated mass of gneiss. To east of it the limestone stretches again far into the mountains, along the valley of the Tons, forming Deoban hill; along the Mussooree ridge it occurs frequently, as on the Abbey and Camel's-back hills. On the top of Landaur it is mixed with sandstones, and appears again by itself on the Tapuban point. The Blaini limestone and conglomerate are well seen on the flanks of the Sarkanda summit, and again in the Ganges at the confluence with the Hiunalgár.

In the Kumaon section we still find the limestone and slate rocks fairly represented. The ridge of Naini Tál is a broad synclinal range, with many local fractures and contortions, like its type the Krol range. The strong limestone that forms the summits about the lake is very like the Krol rock. Here, however, and also at Mussooree, there is a good deal of trappean intrusion. In the Syámkhet valley, north of Naini Tál, trap-rock is in great force, and immediately to the north we come upon crystalline schists. Along the heights of Sunthala and Gággar these are gneissose. The dip throughout is at a moderate angle to north-north-east, and about Almora one or more bands of granitoid gneiss occur in these rocks. Its general mode of appearance is that of interstratification with the schists; but in one place it has been described as intrusive, which would establish its character as a true granite. North of this, for some way, there is a reversal of the dip to south-south-west up to a line of trappean intrusion, which has been traced for many miles along the strike. The rocks to the north of this band are of a more varied character; some are slaty; and limestone is of frequent occurrence, often steatitic in the vicinity of the trap. The dip is less constant in these rocks, and their relation to the crystalline schists of the snowy range is not well defined.

¹ Much detail with suggestive conjectures regarding this region is given in *Memoirs of the Geological Survey of India*, Vol. III, Part 2.

Where we next get a section of the lower Himálaya through the Nepál valley, there is little outward resemblance to what we have seen to the west.

The Nepál section.

The Churia Gháti range, between the plains and the Etounra Dún, is a pattern specimen of the Siwálik type. North of the Dún there is an equally characteristic representative of the Náhan range flanking the mountains. But inside this we no longer find the border mountain-range of slaty rocks capped by plain blue limestone that is so constant to the west of the Káli. We come at once upon schistose rocks. These, however, are not mica schists of the ordinary type, such as those north of Naini Tál. First there are earthy (slaty) schists, some quite black and with carbonaceous layers; then flaggy quartzose schists, passing up into strong schistose quartzites; and these are succeeded by a great mass of dense highly crystalline white limestone within three miles from the Náhan boundary. All are more or less vertical and folded with a prevailing northerly underlie, the strike being 15° south of east. Beyond this steep ascending section there is a broad band of still greater disturbance, apparently a synclinal; for the limestone is variously repeated, and the underlying rocks brought in again. It would seem to be followed in the Chessa-garhi ridge by a crushed anticlinal, about the axis of which there are thick bands of porphyritic gneiss associated with flaggy quartzites. From here there is again a general ascending section through similar flaggy quartzitic schists to the Chandragiri range, bounding Nepál on the south-south-west. This ridge and all those to west-north-west and east-south-east of the valley are made of folded repetitions of one set of rocks, in which a calcareous element is more or less prevalent; varying from strong beds of pure limestone to earthy schistose limestone, and prominently a thickly bedded fine-grained quartzite, with scanty calcareous bond. The Shiupúri ridge, on the north-north-east of the valley, is of massive gneiss; schists appearing again to the north in the valley of the Tádi and the Trisúl Ganga.

There is sufficient resemblance in the two bands of limestone and

Possible affinities.

the underlying flaggy quartzites of this section to suggest that they are repetitions of the same series; and this has certain characters of resemblance to the rock-series at the Krol, which admit of our conjecturing their identity. The whole of this Nepál section exhibits an intensity of disturbance

throughout, such as has not been observed in any other section of the lower Himálaya. Only one instance of a doubtful trap-rock has been noticed in the above section, in the schists near the outer boundary.

We have one more section of the lower Himálaya to take note of, far to the east, in Sikkim, and still more unlike than that in Nepál to the sections of the north-west Himálaya. It is of the highest interest, because we find here in a recognizable state a formation well known to us in the peninsula of India, thus establishing almost the only link between these separate geological provinces. There are, indeed, the nummulitic deposits resting against the edges of the Deccan trap at the base of the western gháts to compare with the nummulitics of Subáthu, and showing that the Himálayas are younger than the Deccan plateau. But this is a comparatively superficial connexion ; whereas in Sikkim we find a bottom formation of one of the great rock-series of the peninsula, intimately connected with the rock forming the mountains. It is now thirty-one years since Dr. Hooker discovered Damuda fossils near Pankabári, at the foot of the Darjeeling hills ; but it was only in 1874 that an examination of the mode of occurrence of the rocks was made by Mr. F. R. Mallet, of the Geological Survey, who was sent to investigate the prospect of a useful coal being found. He traced the band of Damuda rocks from Pankabári to Dalimkot ; at the Tista it is nearly a mile in width. In the western Duárs it does not occur, but on or about its horizon there is a new formation, not found in Sikkim, consisting largely of massive dolomite, and called by Mr. Mallet the Buxa series. In 1875, Major Godwin-Austen, who accompanied the expedition into the Daphla hills, found the Damudas again in force at the base of the mountains in Upper Asám, the Buxa series being there wanting. It is, of course, possible that the Damudas may recur to the west also, in Nepál, though they do not appear at the Sikkim end of the frontier, and certainly they are not specifically represented in the section through Káthmándu.

Lithologically as well as by fossils the formation resembles its prototype in the Damuda valley, consisting of strong sandstones, gray shales, and coal seams. In some few spots the rocks are so little altered that the resemblance is complete, the coal itself being the only rock that has

Its condition and position.

not escaped modification. It is always crushed to powder, and could only be utilized by being made into bricks. Very often, however, the whole group is as much altered as the contiguous rocks to the north, the sandstones being converted into foliated quartzites, and the shales into splintery slates or carbonaceous schists. The important point to settle is the stratigraphical relation of the group to the other rocks of the mountains. These are, first, a zone of slate rocks, some greenish and slightly unctuous, some ordinary clay state with bands of flaggy quartzite, rarely hornblendic and calcareous, and also rarely carbonaceous. These form the Daling series of Mr. Mallet. Next comes the gneiss of the upper hills : it is distinguished as the Darjeeling gneiss. The dip in all these rocks is into the mountain, and hence the immediate inference that they underlie each other in the above order. Here, then, we find again an instance of the puzzle noticed in the section at Simla, and on which Herbert made shipwreck. Mr Mallet was unable to find any escape from the position ; the Daling beds pass most regularly with parallel interstratification into the gneiss by increase of metamorphism, appearing underneath it all up the gorges of the Tista and the Ranjit to the north of Darjeeling, but in a more altered condition. Also he found in several clear sections most completely conformable and transitional junction between the Daling and the Damuda beds. Their junction forms a re-entering angle up the Tista valley ; the Damudas, however, are not known to rise to the surface again in the interior of the mountains. The conviction was forced upon him that they are indeed what they appear to be,—the lowest and oldest rocks of the Himalayan series in this position.

Yet so great is the prepossession in favour of gneiss as necessarily a fundamental rock, that doubt still prevails amongst those who have not seen for themselves. It may then be well to mention some independent considerations which tend to remove this obstruction to the *rationale* of Himalayan geology. It has long since been shown experimentally that the silicious minerals entering into the composition of crystalline rocks can, through the medium of moisture, be produced under very moderate temperature and pressure ; and, hence, that the hydro-metamorphism to which gneiss and even granite were due need not be a very plutonic operation. Also it can be urged that

although when gneiss occurs extensively on the flat, we may be entitled to regard it as a fundamental rock, due to such hypogene action as would require any underlying rock to exhibit as great a degree of metamorphism as itself, yet when we come to mountain formation the case is very different. Here a special concentration of forces has manifestly occurred which may be adequate to the production of this apparently anomalous result. In a recent and very thorough discussion of this branch of geological dynamics we find a direct explanation of our difficulty ; that if a mixed mass of strata were subjected to compression, those portions which by position or texture were least capable of yielding, whether by shrinkage or contortion, would have to bear the brunt of the pressure, and to undergo in some other form its effects, prominently in the development of internal heat.¹ In some such way overlying massive strata may have been converted into gneiss, while softer beds below underwent no crystalline metamorphism. If it should be shown, as in the Simla region, that the Darjeeling gneiss is the same as the central gneiss, the above interpretation of the Sikkim section would have to be abandoned.

Geologists in India have been long on the look-out for a connecting link between the rocks of the Peninsula and of the Himálaya. So many of the formations in the former region are unfossiliferous, it was hoped that some clue to their homotaxis might be obtained through their representatives in adjoining regions ; and although the elevation of those mountains may have occurred in tertiary times, the rocks so upraised might, of course, be of any age. Thus this discovery of the Damudas, as apparently the oldest formation in the Lower Himálaya, at least in the east, comes rather as a surprise. Although the Damuda series is, according to the most recent estimate of its fossils, of lower mesozoic age, its appearance in the field amongst other Indian formations is one of comparative youthfulness. It is, for instance, immensely younger than the great Vindhyan formation, which cover such large areas in undisturbed stratification, but in which as yet no trace of life has been observed, and below which there are several groups of slaty and sub-metamorphic rocks before we come to the fundamental gneiss of the region. However, in these matters, what is, is best ; the object being to know what

Slight correspondence between the rocks of the Himálaya and of the Peninsula.

ing link between the rocks of the Peninsula and of the Himálaya. So many of the formations in the former region are unfossiliferous,

¹ See R. Mallet on Volcanic Energy, Phil. Trans., Vol. CLXIII., page 147.

really has occurred. If the Himálayan sections would reflect to the geology of the Peninsula some light in return for that it has now received therefrom, by clearing up the doubts that still exist about the age of the Damuda formation, it is all we could expect.

There is but one character found almost constantly throughout the Lower Himálayan sections—the frequent appearance of carbonaceous matter. It may be of no great significance, but it is worth notice. Coal-mining had been attempted near Subáthu in the carbonaceous shale of the *infra*-Krol group, or rather where this rock has been compressed and glazed in fault-ground. And this appearance of carbon (some of it as volatile hydro-carbons) with fault-rock is very common throughout the Lower Himálaya. It occurs at Simla, below Nágkanda, at Mussooree, and east of the Ganges. Among the more highly metamorphosed rocks it is represented by graphite, as about Almora. Its appearance in the Káthmandu section, at the outer fringe of the mountains, next the tertiary sandstones, is about the only specific similarity between this section and that in Sikkim, where carbonaceous matter occurs in the Daling and the Buxa, as well as the Damuda horizon. It may provisionally be taken as a suggestive link of affinity between all these rocks. In this connection it is important to recall that in Dr. Stolickza's independent classification the *infra*-Krol horizon corresponds with his Central Himálayan Kuling series of carboniferous age, which is also the age assigned by some to the Damuda formation.

From what we have seen of the rocks of the Lower Himálaya, there is not much prospect of a near solution of this question. The main hope is in the present manifest want of information. Immediately west of Sikkim comes Nepál, taking up the whole middle region of the Himálaya for 500 miles in length, and which through the unaccountable forbearance of our Government, is as much a forbidden land to Europeans as is Chinese Tibet. Except on the single track to Káthmándu, which has lately been traversed by a geologist visitor to the Resident, no Englishman dare set foot in Nepál, whose people are entirely dependent upon our good will for communication with the outside world, and freely enjoy it. In the section to Káthmándu, which is only thirty miles from the plains, we have seen a general

resemblance to the rocks of the Simla region, the contrast being chiefly the universally high dips and general metamorphism. Except in this latter character it has little in common with the Sikkim section ; specific resemblance in the two rock series cannot be made out. The great limestone has no equivalent in kind in the whole of Sikkim. Where distances are so great, much allowance must, however, be made for probable original change in the nature of synchronous deposits. The fact that the present base of these eastern Himálayas adjoined, or indeed formed part of the Damuda land surface, suggests a difference in the deposits near it from those at a distance. We have also had to notice all through our sections a disposition in the Krol limestone to pass into a sandy rock. In some such manner it may yet be shown that the Darjeeling gneiss is on the same horizon as the fine calcareous schists of the Nepál valley. Every geologist will understand the very precarious nature of such a speculation. As an object to confute, it may be of some service.

The Central Himálaya.—In treating of the Lower Himálaya we have had almost entirely to depend upon local names for the various rock groups, and to be satisfied with conjectural identifications of them in different parts of the ground. This unsatisfactory result is owing to the want of fossils. For a great part of that ground it must be a permanent difficulty, owing to the high state of metamorphism of the rocks ; but there are large tracts where this condition does not obtain, and where we may hope that fossils will yet be found. At the snowy range and over a large part of Tibet the case is quite different. There numerous zones of strata are identifiable, not only locally, but in the established scale of formations, by the presence of well-known fossils.

Our information of those regions is, however, in a very fragmentary state, and must long remain so, owing to the inaccessibility of the ground and the rigours of the climate at such great elevations. Numerous travellers have crossed the mountains in various directions, and have brought back a few fossils and isolated observations of the rocks, but only two observers have given a connected geological description of any considerable area. One account is of a portion of Central Tibet in

Strachey, 1848-49.

Stoliczka, 1864-65.

Chinese territory, north of Kumaun visited by Captain R. Strachey, R.E., 1848 and 1849. The description is quoted in *extenso* from his paper, the map attached to which is reproduced to illustrate this chapter. The second and fuller account is that of Western Tibet, by Dr. Stolickza, published in the fifth volume of the *Memoirs of the Geological Survey of India*. In the summers of 1864 and 1865 he explored the region between Spiti and Drás and the Indus. It will be recollected that he died in June, 1874, after crossing the Kárákorum pass, on the return journey with the mission to Káshgár, having lost his life through his zeal for scientific research. His observations on this expedition complete a rough section across the whole Tibetan mountain region from the Panjáb to the plains of Khotan.

Western Tibet. Dr. Stolickza's list of formations, observed in Western Tibet, is as follows :—

	<i>Age.</i>	<i>Group.</i>	<i>Fossils.</i>
IV.—RIVER AND LACUSTRINE DEPOSITS.			
III.—TERTIARY.—	Nummulitic ...	Indus or Shingo Beds ...	<i>Nummulites.</i>
II.—SECONDARY.	Cretaceous ...	Chikkim Beds	... <i>Rudista</i> and <i>Framinfers.</i>
"	Jurassic (Upper).	Gleumal Sandstone	... <i>Aricula eckinata.</i>
"	Jurassic ...	Spiti Shales (Braun Jura).	<i>Ammonites macrocephalus Parkinsoni, triplicatus, etc.</i>
"	Middle Lias ...	Upper Tagling Limestone.	<i>Trochus opulus, Chemnitzia undulata, Terebratula sinemuriensis.</i>
"	Lower Lias ...	Lower Tagling Limestone.	<i>Terebratula groenlandica and pyriformis, Rhynchonella Austriaca, Belemnites.</i>
"	Ethetic ...	Para Limestone	... <i>Megalodon triquetor, Dicerocardium Himalayense.</i>
"	Triassic (Upper).	Lilang series	... <i>Halobia Lomacii, Ammonites floridus etc.</i>
I.—PALÆOZOIC —Carboniferous, Kuling series			
"	Upper Silurian, Math series		... <i>Productus semireticulatus, Spirifer Kothavii, etc.</i>
"	Lower Silurian Bhábeh series		... <i>Tentaculites, Orthids etc.</i>
GNEISS, METAMORPHIC SCHISTS, etc.			
			... <i>Orthids?</i>

The distribution of these formations, though subject to many local irregularities, exhibit well the general structural features of the region. There are two main synclinal rock-basins, along the centres of which the younger members of the sedimentary series, with one important exception, are found. These geological features have no superficial relations to any geographical basins. The Indus receives a great part of the drainage from both areas in this region, flowing nearly along the intervening anticlinal axis. The southern basin is best exhibited in the Spiti valley, where the fullest section of the upper secondary formations is found. The northern synclinal-basin forms the Kárákoram range. This latter ground has been very little searched, and as yet neither oolitic nor cretaceous strata have been observed in it. Dr. Stolickza describes the Kárákoram pass as formed of liassic rocks resting upon trias.

These long rock-basins are bounded by parallel areas of crystalline metamorphic rocks. The southernmost of these is that already spoken of as the central gneiss. Its peculiarity, as compared with the other crystalline ridges, is that we seem to have here what may be locally called the fundamental rock. On Dr. Stolickza's type section at the Bhábeh pass, and in General Strachey's ground, 200 miles to the east, the *infra*-Silurian rocks in a non-crystalline state, and of great thickness, are represented as overlying the gneiss. There is no doubt a parallelism of strike in the two contiguous rock-systems, but the condition suggested or implied is that the conversion of the gneiss is of pre-Silurian date; although, in the east at least, granite seems to penetrate both formations. It was in this sense of *basal* that Dr. Stolickza applied the word *central* to this gneissic axis.

The gneiss and schists forming the middle region of the mountain mass, from 70 to 80 miles wide at the Pangkong and Moriri lakes, are described of a quite different type, as being largely syenitic, and as more or less made up of metamorphosed Silurian rocks. Along the north outcrop of the southern synclinal basin even the zone of carboniferous rocks is hardly recognizable, and all below it is converted into crystalline schists and gneiss, the south-westerly dip continuing in these rocks up to the Indus: similarly along the north-east side of this gneissic

mass the carboniferous formation is the first that is clearly recognizable; all below it being strongly foliated and mineralized, passing with a north-easterly dip under the Kárákorum synclinal basin.

The gneissic axis of the Kuenlun is also described as formed chiefly of syenitic gneiss and quartzose and chloritic schists, the relation of which to the adjoining slates has not been made out. Carboniferous rocks with fossils have been observed on both sides of the range. On the northern flanks of the Kuenlun, triassic and cretaceous deposits are the only secondary formations noticed by Dr. Stoliczka.

Some minor features in the distribution of the several formations will be mentioned presently. We must first notice the remarkable exception, already mentioned, in the general arrangement as above sketched. The nummulitic formation occurs in great force in the Central Himálaya of Ladák, but not even approximately in sequence with the next oldest group. The cretaceous deposits are found capping the sedimentary series in the centre of the southern synclinal basin; but no trace of nummulitic beds has been noticed near them. These occur in force along the valley of the Indus, in the centre of the middle gneissic area. Dr. Stoliczka remarked how strongly the rocks resemble these of the same eocene age at the south margin of the mountains, especially those of the Dagshái and Subáthu groups. It may also be noticed how similar the stratigraphical conditions are in both positions. In the Simla region we saw that the Subáthu beds were deeply unconformable to the contiguous formations, the youngest of which is thought to be triassic. In the Central Himálaya the unconformity is even more striking, because the upper secondary period is well represented, yet the succeeding lower tertiary rocks are in a totally independent basin of deposition. Stoliczka estimates their thickness at 5,000 feet.

There is yet another formation to be mentioned in the Central Himálaya. We have seen granite in connection with the central gneiss. Stoliczka frequently notices greenstone as locally associated obscurely with the Silurian rocks; and intrusive trap (much of it of presumably tertiary age) is occasionally, as has been said, very abundant in the

lower Himálayan rocks. There is, however, one exhibition of eruptive rocks in the Central Himálaya that calls for special notice. It occurs in the middle gneissic zone, locally forming the axis of a mountain range. It is more than ten miles wide at the Hanlo valley, and is continuous thence, with a variable thickness on the south edge of the eocene rocks, to Kargil. Stoliczka describes it as an epidote, diallage and serpentine rock. It appears to be, at least in part, the same rock which he speaks of as syenite to the west of Kargil, where the nummulitic rocks stop out. The eruptive rock there gets entangled in silurian schists, and becomes quartziferous. This igneous rock strongly affects the eocene strata and is therefore of younger date, and is probably connected with the similar rock described by General Strachey on the same strike far to the east. We were able to draw some very instructive warnings from errors of the early school of Himálayan geologists, so we must not let the moderns escape the same ordeal. On Stoliczka's first visit he did not recognise the rocks on the Upper Indus as nummulitic. In that position they are considerably altered, and he took them to be a very old formation. In then writing of the contiguous eruptive rock, he remarked (*l. c.*, page 128): "From their dark colours these rocks have sometimes been referred to *basalts*, but they have certainly nothing to do with these more recent volcanic rocks." His next season's work proved that they cannot be older than middle tertiary, and therefore much younger than the great basaltic formation of the Deccan. The idea of the lithological criterions of age in eruptive rocks has still a strong hold upon the German school of geologists.

It cannot be supposed that the rough cross section we have sketched near the west end of the Tibetan mountain mass can be taken as a type for the immense region to the east. Already within known ground, some interruptions can be pointed out to the longitudinal extension of the several structural zones. Of the continuation of the Kárákorum and Kuenlun, and even of the middle gneissic range, we may be said to know nothing. Stoliczka describes the nummulitic band as completely stopped out against the syenite at Kargil; and although this obtruding rock is at least in part of later date, it is suggested that the termination of the eocene beds here is probably aboriginal. The eastern

Range of the nummu-
lites.

extension of these deposits is quite unknown, save that nummulitic strata occur in the far east north of Sikkim. Some doubtful observations of them in the Changchenmo valley and about the Pankgong lake are recorded.

The southern synclinal basin of secondary rocks, continuous for 200 miles from Spiti to the north-west, is also interrupted in the Kargil region against an upheaval of metamorphosed palæozoic rocks. Any recurrence of regularity in this strike cannot be expected, for the whole Himálayan mountain system becomes confused there, towards the transverse gorge of the Indus, where the stratigraphy is complicated by other systems of flexure. The short break that occurs at the transverse gorge of the Satlaj between the secondary basin in Spiti and the perfectly homologous one to the east, as described by General Strachey, is also due to a transversely obtruded mass of partially metamorphosed palæozoic rocks, through which the stratigraphical continuity is so far maintained.

Although the main elevations, constituting the chain of Himálayan peaks, are sometimes, at least in this western region, formed of the older stratified rocks along the outcrop of the southern synclinal basin, the underlying, pseudo-conformable, central gneiss must be taken as the stratigraphical axis of the range. Stoliczka's type-section of it at the Satlaj and the Bhábeh pass is on the actual continuation of that described by General Strachey to the east, in a more central portion of the great Himálayan chain, where these gneissic rocks are much more prominent. The connection of this gneiss of the main chain with that forming the core of the ridges to the north-west of the Satlaj has not been proved. There are three such ridges. On the direct line of the great chain there is the Dhauladhár ridge, having an axis of coarse gneiss, with slates (probably silurian and lower) resting high on its northern shoulder and passing down into the valley of the Rávi in Chamba. But the Dhauladhár is cut off from the Bhábeh section by the deep valley of Kulu, on the upper Biás, where no massive gneiss has been observed. The Dhauladhár ridge absolutely terminates at Dalhousie, the slates sweeping round the end of the gneiss at the bend of the Rávi. Again, Stoliczka observed his 'central gneiss,' though greatly reduced, north of the

Chínáb, at the southern base of the Zánskár ridge, below the Báralácha pass. But on the south of the Chínáb valley, on the northern flanks of the Rotáng ridge, he observed slates which he conjectured to be continuous with those of the Bhábeh pass; in which case the gneiss of the Zánskár ridge can hardly be continuous with that of Bhábeh section. This ridge of the Rotáng pass is apparently from the maps the structural continuation of the Pir Panjál range, in which a coarse gneiss is again prominent. These three ridges are in a manner confluent in the mountain region of Lahul and Vazíri-Rupi, from which flow the head-waters of the Chínáb, the Rávi, and the Bías. The Bhábeh gneiss strikes into it from the east. Thus it would seem as if the main Himálayan axis broke up into three minor features of the same type in its extension to the north-west.

The gneiss of the Pir Panjál passes beyond Kashmír towards Kaghán. But north-west of Kashmír, at the
 The Kashmír-Kistwár region. Zojila, the gneiss of the Zánskár ridge is extinct, the whole range being there formed of the palæozoic schists and triassic limestone, which thus roll over from the Tibetan area into Kashmír. Carboniferous limestone occurs in the valley, striking through the Marbal pass into Kistwár. Eruptive rocks have been frequently observed in Kashmír, but none of later than silurian age. The whole Kashmír-Kistwár region, between the Pir Panjál and Zánskár ranges, is very little known. Its geological affinities are with the Central Himálaya rather than with the region specially designated as the lower Himálaya, east of the Satlaj.

General Strachey's description of the Central Himálaya, towards
 Central Tibet. the sources of the Indus and Satlaj, is so brief that it can be given in full in his own words, with his summary of conclusions upon the mountain-formation. A few notes are added, giving additional information or suggesting other opinions :—

“Entering the region of the crystalline schists¹ of the great line of peaks, we find the strike still remaining the same, with the dip pretty constantly to the N.-N. E. Along the lines on which

¹ The term *crystalline schists* is often used, as above, to include gneiss. General Strachey hardly uses this latter word, which is now so much applied to distinguish the felspathic and often massive form of metamorphic rock. It is probable that some, at least, of the granite of General Strachey's description is really massive gneiss.

the points of greatest elevation are found in this part of the range, we invariably see for a breadth of several miles, veins of granite

Granite.

in great abundance penetrating the schists, often cutting through them, but perhaps most frequently following the bedding of the strata, between which they seem to have been forced. The great peaks are, I think, in almost every case composed of schistose rock, but the granite-veins may be most clearly seen on the faces of the mountains to very great elevations. Kamet, one of the highest of the peaks in this region, seems, however, to be among the exceptions of this rule; its summit, which is upwards of 25,500 feet above the sea, appearing to consist of granite alone. This line of granite seems to be subdivided into several branches, distributed generally along the strike, but otherwise not very regularly (see map). It appears to consist, where I have seen it, almost entirely of veins of moderate size, and such is probably its general character in the portion of the mountains between the Satlaj and the Káli; but the veins occasionally expand into masses of considerable magnitude, and more rarely large outbursts are met with that constitute whole mountains. In the vicinity of the peak to which I have just alluded, Kamet, the granite-area is very large (see map), and a similar development of it also occurs in the vicinity of Gangotri, at the source of the sacred branch of the Ganges. The vein-granite is usually large-grained with schorl-crystals. It is very hard and durable, neither it nor the schists that accompany it being at all liable to decay. The felspar of all granites that I have seen in these mountains is white, and kyanite is of frequent occurrence in the veins.

“The schists that accompany this granite are very hard and crystalline, and comprise all varieties of mica-schist and gneiss. Beds of highly crys-

Schists.

talline limestones, some pure, others hardly to be distinguished by sight from mica-schist, are of frequent occurrence, and a band of such rocks seems to traverse the country near the line of greatest elevation. The strata, where penetrated by the granite, are often very much contorted, and the dip appears on the whole to increase as we approach the granite, where it reaches an angle of 45°, which it does not often exceed. Thermal springs are met with in many of the valleys along the line of granite, and in several

that I am acquainted with the temperature seemed pretty regularly to be about 128° Fahr. The whole of the appearances presented by the granite and crystalline schists of the great line of peaks in this part of the mountains seem to be universally repeated throughout the whole length of the chain when we reach the region of maximum elevation ; and as we extend our examination, we still continue to find additional reasons for concluding that the general geological phenomena of the range, and the causes that have produced them, remain very similar over great distances.

“ In immediate succession to the crystalline schists penetrated by granite veins, we here come at once upon slaty beds overlying them, along the bottom of which, near the mica-schists and gneiss, is a line of granite-veins differing somewhat in appearance from those of the larger eruption, and not producing any great alteration in the slaty beds themselves, as is shown by the occurrence of a coarse conglomerate, the component parts of which are perfectly distinct, only a few feet above the granite. Sufficient change, however, has taken place to prevent our distinguishing much more than that the constituents of this rock are chiefly quartzose, and that it contains rounded stones of all size. I have met with this conglomerate in a similar position, and with much the same general appearance, thirty miles or so farther to the east. Above these are slaty beds, in all perhaps 9,000 feet in thickness, consisting of coarse slates, grits, and limestones, all more or less affected by slaty cleavage, and all devoid of fossil remains.

“ It is after reaching the top of these strata, which is rarely done at a less elevation than 14,000 feet above the sea, that we at length enter again a region of fossiliferous rocks, which extends as far as my examinations have been carried. And it is not a little wonderful to find at this immense elevation a regular succession of most of the more important formations, from the Silurian to the Tertiary Periods. The Palæozoic beds met with immediately above the slaty rocks I have just mentioned seem to have a thickness of about 6,000 feet, but it is quite possible that organic remains may extend lower than I supposed ; indeed, from the very difficult nature of the country, the precise thickness of the deposits and the limits of the different formations cannot be determined properly without a much more

careful examination of the country than I was able to give it. The lower portion of these strata are undoubtedly of lower Silurian age, and I am indebted to Mr. Salter for the following list of the species that he has been able to recognize on a somewhat cursory examination of my specimens.

"Among the Trilobites are—*Cheirurus* (the Silurian form of the genus), *Lichas*, *Asaphus* (only as yet found in Lower Silurian beds), *Illanus*, *Calymene*, *Protopiscus*, *Sphaerexochus*.

"Of Molluscs are—*Strophomena*, a strongly ribbed *Orthis*, *Terebratula*, *Leptæna* very like *L. depressa*, *Lingula*, *Orthoceras*, *Cyrtoceras*, *Lituites*, *Theca*, *Bellerophon*, *Murchisonia*, *Pleurotonnaria*, *Raphistoma*, and *Ctenodonta*.

"Of Polyps—*Ptilodictya*, *Chaetetes*.

"Also *Encrinites* and *Cystidæ*, *Tentaculites* and other Annelids and Fucoids.

"I had also an opportunity of showing these fossils to M. Barande, who appeared to have little doubt, from their general character, that some of the beds from which they came were certainly of Lower Silurian age.

"The lowest beds of these Palæozoic strata consist of dark-coloured thickbedded limestones, in some places filled with corals. They are succeeded by limestones mixed with slates, in which were found the strong-ribbed *Orthis*, *Terebratula*, *Lingula*, *Bellerophon*, and fragments of *Encrinites*. Above these come flaggy limestones with grits, that contain the greater part of the *Trilobites*, *Strophomena*, *Leptæna*, *Lituites*, *Ptilodictyon*, *Cystidæ*, and Fucoids. The beds then become more argillaceous, and shales and slates mixed with an impure concretionary limestone follow. In these beds are found *Cyrtoceras* and *Orthoceras*, and amongst the nodular concretions of limestone a *Chaetetes* is common. Next in order come dark-red grits, sometimes marly, containing only a few fragments of Encrinital stems. Above these, pale flesh-coloured quartzite, and finally a white quartzite, in neither of which I ever found any fossils, and which form the highest peaks of the ridges composed of the Palæozoic rock. The whole of these strata are in various degrees affected by cleavage and joints, which penetrate all the beds without regard to their

mineral character, although in a somewhat less marked degree in the limestones and quartzites. That the general sequence of these strata is pretty regularly maintained, I have seen over a longitudinal extent of about fifty miles, but it appears highly probable that their development has a far greater range, as we shall also see to be the case with some of the other groups of the fossiliferous rocks.

“ Before passing on, I must observe the very remarkable similarity of general mineral appearance that
 Compared with England. subsists between the Silurian rocks of the Himalaya and of England. The peculiar pale tint assumed by many of these rocks answers most exactly to the descriptions given by Sir Roderick Murchison of the Silurian districts of Wales, and the characters of the concretionary limestones of both countries appear equally to correspond. Even in hand specimens the texture and appearance of the rocks and of the fossil impressions are so similar that they might most readily be mistaken one for the other. In pointing to these resemblances, however, I would not have it supposed that I should wish in any way to set up mineral character as a criterion by which to decide on the age of any rock. Nevertheless, the facts, if they are to be relied upon, would appear to indicate that as we see the conditions of the existence of organic matter to have been generally similar over large areas, or even over the whole earth, during the same epoch, and to have changed with the progress of time, so likewise has it been with the conditions under which the mineral constituents of the earth have been aggregated.

“ The Palæozoic strata that I had an opportunity of examining
 Silurian predominates. in detail *in situ*, which I have just been describing, appear to be exclusively Silurian, but the existence of rocks of Devonian or Carboniferous age seems to be shown by some of my specimens, not found *in situ*, which contain *Productus*, *Chonetes*, *Athyris*, *Orthis*, *Aviculopecten*, *Spirifer*. I may here be allowed to repeat that the higher portions of the Silurian rocks being usually found at elevations of 17,000 or 18,000 feet, their examination is not a very easy task, and the difficulties occasioned by the great altitude are infinitely aggravated by the confusion into which the beds are thrown by the vast dislocations. that have accompanied the elevation of these mountains. In concluding my remarks on the Palæozoic beds I would observe that,

as a general rule, to which, however, there are no doubt many exceptions, these rocks are to be found forming the summits of the highest passes between the British provinces of Kumaon and Garhwal and Tibet, which probably average 18,000 feet in elevation, and that the highest points of the ridges on which these passes are found not unfrequently reach nearly 20,000 feet in altitude.

“In proceeding along the section, we shall next observe some beds very remarkable from their apparently close similarity to the Trias of Europe. I can now only regret that, not having been sufficiently aware of their importance, their exact relation to the beds below them has not been better made out ; but their position in the series immediately above the Palæozoic rocks is at least certain. In one place these strata were found *in situ* intermediate between the Palæozoic and Secondary rocks, but the greater part of my specimens were obtained from fragments lying on the north slope of the Palæozoic ridge, which appears to terminate with a line of fault, to the north of which a cliff of Oolitic age suddenly rises. From these strata I have obtained not less than twenty-five species of fossil shells, which is a remarkable circumstance, considering the small bulk of the specimens that I was able to bring away with me. Mr. Salter, who has been so good as to examine these also, tells me that we have *Ammonites* several, *Ceratites*, *Orthoceras*, *Natica*, *Exogyra*, *Halobia*, (*Avicula*), *Pecten*, *Lima*, *Athyris*, *Waldheimia*, *Rynchonella*, *Spirifer*. The Triassic beds were chiefly dark-coloured limestones and, where seen *in situ*, were associated with shales and dark-red grits, the latter of which seemed very similar to those found near the top of the Palæozoic series. The line on which they were seen was, however, a very bad one for determining such matters, for it was in one of the great valleys, and consequently on a great dislocation where accumulations of *débris* almost always greatly predominate over rock *in situ*.

“In our progress northward, we next come upon the strata that form the representatives of the Jurassic group. As in the Palæozoic beds, so we here find the general dip to be to the north ; but it is impossible for me to offer any opinion as to the degree of conformability of any of these deposits one to another, owing to the great disturbances to which they have everywhere been subjected. It appeared to me, however, as

probable that in the parts of the mountains that I examined, a great line of fault intervened between the Oolitic and Palæozoic series. The mountain-ridge of Silurian age most carefully examined by me lies generally parallel to the line of strike, and along its north-east face runs a stream separating it from the Secondary rocks, which rise in an almost impassable precipice beyond. The section here exposed must be at least 5,000 or 6,000 feet in thickness, but the difficulties of the route prevented my extending my examinations into the lower beds. The lowest that I reached were of black limestones and shales, with very few organic remains, and those very imperfect. Above these lie several thousand feet of limestones of various descriptions, the rock in some places being almost made up of fragments of shells. Professor Forbes, who has kindly looked over my specimens from these beds, is inclined to identify some of the species with certain forms that occur in the Fuller's Earth and Cornbrash of England; and it appears that there is here no representative of the Lias.

“ Continuing to ascend in the series, we reach next a large development of dark-coloured shales which abound with remains of *Ammonites* and *Belemnites*, the former usually imbedded in spherical nodules, apparently of much the same nature as the shale itself, but exceedingly compact. The shale is for the most part, on the other hand, very rotten, and the band of country along which it is found is often depressed so as to form a valley, apparently in consequence of this disintegration of the rock. This shale Professor Forbes pronounces to be without doubt of the age of the Oxford Clay, a conclusion indicated by the peculiar forms of the *Ammonites*, two of which seem to be identical with species found in beds of the same age in Kachh and Sind, which have been figured and described in the Transactions Geol. Soc.¹ The existence of these beds in the northern parts of the Himálaya was pointed out by Sir Roderick Murchison some years ago, as proved by the occurrence of some of these *Ammonites*, which he had seen. There is indeed direct evidence of the existence of these Oxford Clay strata for a distance of about 200 miles to the westward of the places where I have myself seen them, and their prolongation along the north of the

¹ N. Ser., Vol. V.

mountains for 200 miles more in an easterly direction is rendered highly probable by the well-attested recurrence of the *Ammonites* in the eastern parts of the kingdom of Nepal. Although we find stratified deposits apparently lying conformably on the Oxfordian strata, I cannot say anything definite regarding them, as they appear to be almost entirely devoid of fossils. They are very hard and compact, consisting of grits, shales, and limestones, and have not improbably been converted into their present state by the action of eruptive rocks which are of common occurrence in this region.¹

[Subsequent to the publication of General Strachey's papers in the Journal of the Geological Society, a description of his collections of fossils was drawn by Messrs. Salter and H. F. Blanford, and printed for private circulation. The following complete lists are taken from that work :—

SILURIAN FOSSILS.

Asaphus emodi.
Illænus brachyonisus.
 Do. *punctulosus*.
Cheirurus mitis.
Protopiscus mimus.
Sphaeræxochus idiotes.
Lichas Tibetanus.
Calymene nivalis.
Tentaculites sp.
Serpulites sp.
Nautilus ? involvens.
Cyrtoceras centrifugum.
Lituites uliformis.
Orthoceras striatissimum.
 Do. *Kemas*.
Theca lineolata.
Bellerophon Ganessa.
Strophomena trachealis.
 Do. *chæmerops*.
 Do. *umbrella*.
 Do. *aranea*.
 Do. *nubigena*.
 Do. *bisecta*.
 Do. *halo*.
 Do. *lineatissima*.
Orthis Thakil.
 Do. *Tibetica*.

Murchisonia Himalensis.
 Do. *pagoda*.
Pleurotomaria turbinata.
Raphistoma emodi.
Trochonema hamifusa.
Cyclonema rama.
 Do. *subtersulcata*.
Holopea varicosa.
 Do. *pumila*.
Otenodonta sinuosa.
Cyrtodonta ? imbricatula.
Lingula Kali.
 Do. *ancyloides*.
Leptæna Himalensis.
 Do. *repanda*.
 Do. *cratera*.
 Do. *nux*.
Orthis compta.
 Do. *monticula*.
 Do. *uncata*.
Ptilodictya ferrea.
 Do. *plumula*.
Sphaerospongia mellifera.
 Do. *inoculana*.
Chæstetes ? Yak.
Heliolites depauperata.

CARBONIFEROUS FOSSILS.

Productus Purdonii.
 Do. *Flemingii*.
Chonetes Vishnu.

Athyris Roissy.
Aviculopecten hyemalis.

¹ In 1873 Mr. T. H. Hughes, of the Geological Survey, brought some fossils from the Milam Pass. They were identified by Dr. Waagen as representing cretaceous, jurassic, triassic, permian, carboniferous, and silurian formations. This brings the series here into fuller agreement with that described by Stoliczka in Western Tibet.

TRIASSIC FOSSILS.

Ammonites floridus.
 Do. *Aon*.
 Do. *Winterbottomi*.
 Do. *planodiscus*.
 Do. *diffusus*.
 Do. *Gaytani*.
 Do. *Ausseneanus*.
 Do. *Blanfordii*.
Ceratites Jacquemonti.
Orthoceras pulchellum.
 Do. *salinarium*.
Natica subglobulosa.

Exogyra sp.
Halobia Lommeli.
Pecten scutella.
Lima Stracheyi.
Athyris Deslongchampsii.
 Do. *Strohmeyeri*.
Waldheimia Stoppani.
Rhynchonella retrocita.
Spirifer Oldhami.
 Do. *Stracheyi*.
 Do. *Rajah*.

OOLITIC FOSSILS.

Belemnites sulcatus.
Anopites acucinctus.
 Do. *alatus*.
 Do. *bifrons*.
 Do. *biplex*.
 Do. *communis*.
 Do. *concavus*.
 Do. *Eugénii*.
 Do. *Gerardi*.
 Do. *Griffithii*.
 Do. *guttatus*.
 Do. *Hookeri*.
 Do. *heterophyllus*.
 Do. *Himalayanus*.
 Do. *Hypbasis*.
 Do. *Jubar*.
 Do. *Medea*.
 Do. *Nepalensis*.
 Do. *octagonus*.
 Do. *robustus*.
 Do. *scriptus*.
 Do. *Spittensis*.
 Do. *strigilis*.
 Do. *tenuistriatus*.
 Do. *torquatus*.
 Do. *triplicatus*.
 Do. *Thouarsensis*.
 Do. *umbro*.
 Do. *Wallichi*.
Turritella montium.
Pleurotomaria ? sp.
Turbo invitus.
Chemnitzia sp.

Anatina Avaginula.
Myophoria Blanfordi.
Cardium truncatum.
Cyprina trigonalis.
Astarte major.
 Do. *unilateralis*.
Modiola sp.
Nucula chneiformis.
Cucullæa virgata.
 Do. *leionota*.
Inoceramus Hookeri.
Lima acuta.
 Do. *gigantea*.
 Do. *mitiloides*.
Monotis concentrica.
Avicula echinata.
 Do. *iniquivalvia*.
Pecten iniquivalvia.
 Do. *comatus*.
 Do. *bifrons*.
 Do. *monilifer*.
 Do. *lævis*.
 Do. *Sabal*.
Ostrea flabelloides.
 Do. *acuminata*.
Terebratula numismalis.
 Do. *carinata*.
 Do. *globata*.
Rhynchonella variabilis.
 Do. *concinna*.
Acrosalenia ?
Pentacrinites sp.]

“ But the most striking feature of the geology of these mountains is probably that which I have next to mention, viz., the existence of a great Tertiary deposit at an elevation of from 14,000 to 16,000 feet above the sea, still preserving an almost perfectly horizontal surface. On crossing the watershed-ridge between the streams that flow to the south into the Ganges, and those that fall into the upper part of the Sutlaj to the north, which here constitutes the boundary between the British territory and Tibet (see map), we find ourselves on a plain

120 miles in length and varying from 15 to 60 miles in breadth, that stretches away in a north-westerly direction. Its western portion is everywhere intersected by stupendous ravines, that of the Satlaj being nearly 3,000 feet deep. The sections afforded by these enable us to see that this plain is a deposit of boulders, gravel, clay, and mud of all varieties of fineness, laid out in well-marked beds that run nearly parallel with the surface, and that hardly deviate from a horizontal position. The discovery of the fossilized remains of several of the larger mammalia distinctly marks the Tertiary age of this deposit. The existence of such fossil remains in the northern parts of these mountains had been long known, but we were altogether ignorant of the precise locality whence they came, and had no facts before us from which any conclusions could be formed as to their geological import. The Niti Pass, from which it was said that the bones had been brought, was not the place where they were found, but one of the routes only by which they came across the great Himálayan chain from unknown regions beyond.

“Mr. Waterhouse, who has been so obliging as to examine the specimens that I procured from these so-called Niti fossils. beds, informs me that he recognizes amongst them the following :—Metacarpal bone and distal end of tibia of *Hippotherium* ; patella of small horse ; distal end of radius of a larger species of horse ; distal half of tibia of a horse of very large size ; part of metacarpal of a horse ; upper end of tibia of bovine ruminant ; dorsal vertebra of a ruminant. Portion of head of an undescribed animal allied to goat and sheep, having, like them, prominent orbits, and the horns above the orbits ; but which differs in the peculiar form the bony core of the horns. The horns are remarkable for being placed very near to each other at the base (their upper portions are broken off). There is a specimen in the British Museum, however, from the same locality, of an animal very like this, in which the horns are seen to be short, stout, and slightly bent outwards at the apex. Right wing of the atlas vertebra of rhinoceros ; phalanx of one of the outer hind toes of ditto ? ; and portion of tooth of elephant ? Specimens of the bones of ruminants, pachydermata, and other animals from this district, presented to the Society by Sir Thomas Colebrooke and Dr. Traill, are in the Museum of the Geological Society, London.

“The bones that we have hitherto obtained from these strata are almost all very miserable fragments, so that Actual site as yet not identified. it is difficult even for the very learned naturalists that I have mentioned to do more than distinguish the genus to which they belong. It is therefore, I am afraid, at present impossible to come to any decided conclusions as to the identity or otherwise of the species here found with those of the Siwálik hills, a question of the greatest interest with reference to all our speculations on the geology of these mountains. The fossil bones I have not seen *in situ* nor indeed, curious to say, could I, in spite of every attempt, learn a definite locality in which any one knew positively that they had been found. But of the general position where they occur there can be no doubt, for, besides the common account of their being found in some of the ravines that traverse the plain, on many of the specimens quite enough of the rock in which they are imbedded has remained to enable me to recognize a fine-grained calcareous conglomerate exactly identical with beds such as I have seen intercalated with the boulder and gravel beds that constitute the mass of the deposit. Hills of limestone rise here and there above the general level of the plain, and it appears as though the calcareous matter derived from them had cemented together portions of the sands and gravels that were deposited near them.

“The existence of such animals as I have mentioned as being found in these beds being a physical impossibility in the present state of the country, Probably of marine origin. there can be no doubt that the strata have been elevated to their present height from some lower level since the time of their deposition. There is no direct proof that these beds are marine, no shells having been obtained from them, but I think on the whole that the probabilities appear to be in favour of this plain having been a true sea-bottom rather than of having been occupied by a detached body of fresh water. The general extension of some of the older fossiliferous rocks along the northern face of the Himalaya over a great longitudinal distance is a fact of which we have tolerable proof, and it thence follows that the line on which they occur, distant about twenty or thirty miles to the north of the great line of peaks, has probably been a sea-margin from the remotest ages of the earth's history until as late as the Oolitic

period at least. So far, therefore, there is nothing adverse to my supposition; nor is the present interruption of the plain any proof that it did not once have a far greater extension. This is sufficiently proved by my having traced these tertiary beds to the very top of the watershed-ridge in the vicinity of the Niti Pass, where they reach an elevation of upwards of 17,000 feet; the summit of that pass being strewn with boulders that appear to be derived from the white quartzite capping the Silurian strata of the neighbourhood. Further, two or three miles to the south of the pass, a detached portion of this deposit is to be seen on the declivity of the mountain, which must have been separated from the general mass by the dislocations that have upheaved the whole country. It is, moreover, to be noticed that there seem to be grounds for supposing that plains, such as I have mentioned, are found in other parts of the chain under somewhat similar circumstances, which may not improbably have once formed portions of the same sea-bottom. The plain of Pámír, so long known from the accounts of Marco Polo, and the existence of which is fully corroborated by Lieut. Wood of the Indian Navy, in his Narrative to his Journey to the Source of the Oxus, may be its representative to the west; while to the east the plains described by Turner as having been passed over during his embassy into Tibet, as well as others mentioned by Kirkpatrick as existing to the north of Nepál, the descriptions of which are quite confirmed by Dr. Hooker, are not improbably of a similar nature. Another argument in favour of the marine origin of this deposit is, I think, also to be derived from the very regular way in which the beds of gravel and boulders are laid out, for which I should conceive that some action like that of the tides would be requisite.¹

¹ No notice has been taken of these great detrital accumulations up to this, because they occur on the grandest scale in the regions described by General Strachey; but every explorer in the Ladák country has noticed deposits of the same kind. In many cases they are very recent. The process of their formation has been very well described by Mr. Drew in the Quarterly Journal of the Geological Society for 1873. From the fossils enumerated above, it is, however, certain that some of them are very ancient. The fact that General Strachey did not observe the actual site of these bones leaves it open to conjecture whether they may not occur in some disturbed beds more or less covered by these horizontal gravels; for it is very difficult to conceive how any great dislocation or upheaval of the mountains can have occurred without disarranging such incoherent materials. On the other hand, the fact that such large animals as the Yak, the Kiang, the Ovis Ammon, &c., now flourish in the wild state in those bleak sterile regions suggests that a moderate change of climate, without any great change of elevation, might make them habitable for the fauna now found fossil there. No evidence for marine action later than the nummulitic period is known within the Himalayan border.

“ I have already mentioned the occurrence of eruptive rocks in the Tibetan plateau. A great outburst, in which are found hypersthene and bronzite, besides syenitic and ordinary greenstones, and various varieties of porphyry, occurs in the vicinity of the lakes which are found at the eastern extremity of the plateau (see map). The greenstone is known to extend considerably to the west, and forms, at an elevation of about 17,600 feet, the summit of Balcha, one of the Himálayan passes into Tibet which I have crossed.

“ Having thus given a general description of the geology of this region, I shall, as shortly as possible, enumerate the chief conclusions to which I have been led with regard to the physical forces that have been called into action in the formation of these mountains :—

(a.)—The general extension of the chain along the direction of the strike of the strata is a phenomenon necessarily connected with the action of an upheaving force along a line. This longitudinal action is further evinced by the parallelism of the lines of eruptive action with that of the *strike*. The continuance of action of the upheaving forces along the same general line for a vast period of time, with occasional intervals of repose or of subsidence, is indicated along both the north and south faces of the Himálaya. The great depth at which the forces have originated seems to be proved by the regularity of the action along the entire length of the chain, as shown by the elevation of such a ridge as the Siwálik hills.

(b.)—The granites appear to constitute lines of elevation, not of rupture ; but there seems to be no specific action produced by them on the dip of the strata, which they appear to leave generally unchanged.

(c.)—The greenstones, on the other hand, usually follow lines of dislocation of the strata, being sometimes apparently contemporaneous, and at others intruded through rocks already consolidated.

(d.)—The cause of the general north-easterly direction of the dip is obscure, although its occasional sudden reversal to south-westerly seems to indicate some connexion with the action of an upheaving force from below, or of violent lateral thrust.

(e.)—The lines of fracture of the strata are constantly either parallel or perpendicular to the direction of the upheaving force. The positions of the rivers appear to be altogether dependent on the configuration of surface produced by these fractures ; while the configuration of surface, on the other hand, seems to be but slightly affected by the action of the streams, of which there is rarely any visible sign at 200 feet above the present level of the waters, and never to my knowledge above 300 feet.

(f.)—The fact of the granite of the great snowy peaks being seen in veins, penetrating the schists up to 20,000 feet, makes it highly probable that the granite must have been injected long before the mountains received any considerable development. That this granite is older than the Silurian period is rendered probable by the comparatively unaltered state of the lower beds of the Azoic slates at the foot of the Palæozoic series, where almost in contact with it.

(g.)—The conglomerate bed near the bottom of these same Azoic slates shows the proximity of land at the time of its deposit, and indicates that some upheaval of land had already taken place near the present line of great peaks, possibly occasioned by the granite in question.

(h.)—The occurrence of pebbles of greenstone in the sandstones along the southern edge of the mountains shows that the exterior lines of greenstone are older than those beds of sandstone.

(i.)—The frequent occurrence of boulders of the quartzites, slates, and greenstones of the outer ranges of mountains among the Tertiary deposits of the Siwálik hills shows that the Tertiary ocean washed the foot of those mountains.

(j.)—The regular slope of the plains of Northern India up to the Siwálik hills, which rise suddenly from the flat ground, leads me to infer that the sea must have continued to reach at least as far as the foot of the Siwálik hills for some time after their upheavement.

(k.)—The rise of the Tibetan plain has not been caused by the granite eruption of the line of snowy peaks. That the greenstone rocks that abound in many parts of it have equally not caused it, is proved by the peculiar nature of the valleys among the hills to

the west of the lakes, which must have been laid out level under water ; from which it is to be inferred that these eruptive rocks are older than the tertiary beds of the plain. The same thing is shown by the occurrence of worn pebbles of greenstone in the surface of the plain in the vicinity of some of the detached hills of that rock.

(l.)—The former extension of the glaciers far beyond their present limits is a phenomenon that may be noticed almost everywhere in these mountains, and may give rise at first sight to an idea that there may here also have been some special period of cold corresponding to the glacial epoch of Europe. But it seems, I think, more probable that this is here only the result of a change of climate consequent on the upheaval of the great plains of Northern India.

(m.)—The existence of ancient moraines on the tertiary plain of Tibet proves that the extension of the glaciers is post-tertiary. Now, if we conceive that after the rising of this plain to nearly its present elevation, the sea still continued to wash the foot of the Siwálik hills, as I have already said that I considered likely, it is clear that the climate of the Himálaya would have been far more moist, and that the quantity of snow that fell on the highest parts of the mountains would have been greatly in excess of what now falls there, causing a great extension of the glaciers beyond the limits to which they have now receded."

There are several points in the foregoing summary that might be objected to in detail, but it is hardly necessary to point them out. With the much more extensive information now available than was at General Strachey's disposal, one ought to be able to give a more precise account of the phenomenon under discussion. But one lesson of experience is caution. From all sides the geologists of Europe have been for long years hammering at the Alps ; yet the mode of formation of those mountains is still a subject of very vague speculation. What then can we expect from our fragmentary knowledge of Himálayan geology ? In one respect we seem to have the advantage : the much grander scale on which the phenomenon took place, and perhaps also the less advanced stage of the process, have resulted in a somewhat less complexity of structure.

The process of investigation is to find out as far as possible from these remnants of the formations, and their relations to each other, what were the conditions of the surface at the time of deposition of each, and what successive changes of conditions occurred. Thus General Strachey's observations that the metamorphism of the rocks forming the crystalline axis and the introduction of the granite are of pre-silurian date; that this axis of elevation was a shore of deposition at that time, would be facts of prime importance in the early history of this region. It would give a prodigious antiquity to the beginning of the Himálayan mountain system, for all subsequent disturbances have conformed in elevation with that which produced the gneissic axis. It may be remarked that this view is apparently in immediate opposition to what is now a very favourite theory of mountain-formation—that which connects these areas of special contortion and elevation with a preceding long-continued accumulation of deposits, and accompanying depression, in the same area; whereby through the gradual rise of temperature in the sediments thus sunk to a considerable depth, expansion occurs, and also a softening of the rocks, including a yielding to the horizontal thrust in the earth's crust, thus producing the compression and up-squeezing into mountains of the accumulated sediments.¹ In the simple application of this process the position of greatest elevation should approximately coincide with that of the preceding greatest deposition, and not with the limit of the deposition, as we find it according to General Strachey's observation.

To this objection, and in defence of the theory, it may be very fairly argued, that the line of actual maximum elevation is due to denudation having removed the softer and more broken strata and left the more massive rock; that according to the observation under discussion, the sedimentary series never passed across this primitive gneissic axis; and that the position of greatest elevation (in the active sense of upheaval) occurred about the middle gneissic axis formed of metamorphosed palæozoic rocks, all the once overlying strata having been removed; that but for this result of denudation we should have the crest of the Himálaya to the north of the upper valley of the Indus.

¹ Notwithstanding the numerous recent claimants to this theory, the mechanical elements of it are essentially those given by DeBeaumont in his *Système de Montagne*, p. 1318: the other ideas in it being due to two other equally eminent philosophical physicists, Herschel and Babbage.

The interpretation of the drainage system gives direct support to this view. It is an evident postulate of physical geology that along any line of elevation the drainage is originally transverse. The manner in which this primitive system becomes largely converted into longitudinal drainage lines is explained in every text-book on geology. Now, making the fair assumption that the initial line of elevation coincides with the maximum line of upheaval, the main watershed of the future continent is determined by that initial line, and it is presumably a very permanent feature. Every geography book notes the fact that the Himálayan watershed lies far to the north of what is more particularly described as the Himálayan range, but the line of reasoning we have just indicated would suggest that the real axis of maximum elevation in the Himálayan system may coincide more or less with the watershed. We should thus have two magnificent examples of the process of drainage-conversions above alluded to: the Indus and the Sanpo (Brahmaputra) now flow from about the same central position, having gradually worn back along the line of easiest erosion, cutting off in succession the originally transverse drainage along the whole line.

The applicability of this theory of mountain-formation to the Himálaya system does not, however, depend upon the correctness of General Strachey's view regarding the exceeding antiquity of the first gneissic axis. The gneiss there in early Palæozoic times

Stoliczka's views. may have been a floor of shallow deposition without being a range limiting that deposition.

Stoliczka does not adopt this latter view though, leaving it an open question. His own provisional identification of lithologically similar Silurian rocks in equal force to the south of the axis would, perhaps, suggest their original continuity across that axis. He hazarded very few remarks upon the general geology of the Himálaya, wisely postponing such considerations until the data would warrant something definite. He points out that the deposits of carboniferous age, filling only broken ground in variable thickness, represent the close of a general geological epoch. The Permian and lower Trias are not represented. He considers that after the Trias extensive upheavements occurred, laying dry large tracts that have not since been submerged. The Jurassic basin was then approximately defined. The evidence for these conclusions is

not given, and they seem difficult to reconcile with the general regularity of succession of the rocks. On his figured sections the Para limestone (Rhætic group) is the only one that exhibits great inequality of distribution, being of considerable thickness on the north side of the basin, and altogether wanting to the south. His remarks would, however, assign at least an early Mesozoic age for the origin of some of the prominent features of disturbance now stamped upon the Himálayan system.

The most notable features in the sequence of formations in the Himálaya is the position of the nummulitic deposits, as already mentioned. They are in greater thickness than any of the older formation except the lower Silurian. Stoliczka speaks of them as having been deposited in the narrow basin where they now lie in the very centre of the mountain region. This view strongly confirms that previously arrived at from the consideration of the same deposits at the south edge of the mountains: that long and extensive denudation of the Himálayan area had preceded the Tertiary epoch. It seems to have been greatest, as would naturally occur, along the centre of the area of upheaval, wearing down to the metamorphic rocks along what may then have been the back of a broad flat tubercence of the earth's crust. A comparatively slight settlement of the area, submerging only the more deeply eroded parts, would then have sufficed for the accumulation of those eocene deposits, and it seems possible that the production of the synclinal basins of the Central Himálaya—as was shown for the contortion of the infra-Krol beds of the Lower Himálaya—did not occur, or, at least, was not strongly developed till the great compression upheaval in middle Tertiary times.

CHAPTER IV.

THE HIMÁLAYA.

CONTENTS.

The outer Himálaya. Mountain systems; Jumna; Ganges; Káli. Subordinate boundaries. The Nayár. Western Rámungga; Káli. The snowy range. Axis of highest elevation. The Tibetan plateau. Glaciers. Glacier of the Pindar; feeders of the glacier; moraines; crevasses and ice tables. Glacier of the Kuphni. Motion of the Pindar glacier. Snow-line. Observations in the eastern Himálaya. Observations in Kumaon. Bishahr. Across the snowy range. Kumaon; Garhwál; Kunaor; Western Tibet. Lakes.

WE have seen that the term 'outer Himálaya' has different meanings according to the sense in which it is used. Geographers understand by it all the mountain systems lying between the snowy range and the plains of India, whilst in the western Himálaya geologists restrict its use to the limestone and slate formations that lie between the outlying tertiary series and the central crystalline axis. To the geologist Simla and Naini Tál are situate on the outer Himálaya, and Kasauli and Subáthu are not; whilst to the geographer all these hill sanitaría are on the outer Himálaya. To obtain a correct appreciation of the physical relations of the ranges of the outer Himálaya we cannot ignore their geological affinities, but we have as yet no geological details on which we can rely for the greater portion of their area. For the tract between the Tons and the Káli we have the record given in the preceding chapter and the map that accompanies it. A glance at this map will show us that the main line of gneiss and granite, which is almost conterminous with the region of perpetual snow, is met with between the eightieth and ninetieth mile from the foot of the Himálaya, and at a distance of from twenty to thirty miles south of the Indian water-parting. Between this line of perpetual snow and the plains there are numerous well-defined ranges, some seemingly spurs from the snowy axis, and of which the geological relation is not well established, and others having an apparent separate and independent existence. As we have already noticed, the traveller from the plains meets first an outer range, which has a general elevation of about 6,000 to 7,000 feet above the level of

the sea, and which in China above Naini Tál rises to 8,568 feet, and in a peak, on the Tirhi road, east of Mastúri, to 8,565 feet. This band or outer range has a general direction parallel to the plains, and is pierced by the greater rivers, such as the Tons, Jumna, Ganges, Rám-ganga (western), Kosi, and Káli. Between it and the ridges descending immediately from the snowy range we cross a number of subordinate ranges which are, as a rule, of considerably less elevation than the outer range. Between China and Dhákuri Bináyak, on the road to the Pindari glacier, a distance of forty-five miles as the crow flies, there is no peak having an elevation of 8,000 feet, and very few attaining to 7,000 feet. But further westward, in the same parallel, we have groups of peaks attaining an elevation of over 10,000 feet, notably those connected with the Duduka-toli range in Garhwál. When the river valleys close to the snowy range are reached, the increase in elevation is rapid and marked. The flanking ranges seldom fall below 10,000 feet, and are crowned with peaks rising still higher, until the culminating ridge crowned with perpetual snow is met.

If we carefully examine the great sea of mountains lying between the outer Himálaya and the snows,

Mountain system.

we shall find that the dominating ranges are spurs from the great groups of peaks, remarkable alike for their elevation and the position they fill as the boundaries of the several river-basins. On the west, the western boundary of the Jumna system is found in the elevated ridge that has its origin in the group of peaks crossed by the Shatúl and Burenda passes. This ridge follows the left bank of the Satlaj in a south-westerly direction to Hatu (10,700 feet), where it bifurcates: one branch continuing the normal direction to Biláspur, and the second proceeding in a south-easterly direction

Jumna system.

by Chor (12,081 feet), where it forms the water-parting between the Giri and the Pábar branch of the Tons. A second great ridge, descending from the Jamnotri groups, and marked by the Deoban (9,847 feet), Chakráta (7,300 feet), Chilmeri (7,160 feet), and Bairát (7,423 feet) peaks in British territory, separates the affluents of the Tons from those of the Jumna. The eastern boundary of the Jumna system is formed by a great ridge having its origin in the same group of peaks, and which joins the outer

Himálaya near the Sarkanda peak to the east of Masúri (Mus-sooree). The eastern boundary of the Gangee system is found in the great ridge descending in a south-westerly direction from

Ganges system. the Nandakot peak, and which passes along

the left bank of the Pindar to its junction with the Alaknanda, and thence along the left bank of the Alaknanda to Deoprayág. It admits of two great bifurcations: one at the head of the Katyúr valley and one at the head of the Lohba valley. From the group of peaks at the head of the Katyur valley a branch passes in a south-easterly direction through Binsar and Dol and along the right bank of the Ladhiya to Barindeo on the Sárda, and

Káli. from the head of the Lohba valley a branch

runs south-by-east to Gujargarh, whence it passes almost due west to the Ganges at Kharak and Chándi. The tract to the south of these two arms is in shape a great obtuse-angled triangle, with its base towards the plains and its apex in the group of hills to the north of Dwára Hát. It forms the mountain basin of the western Rúnganga. The eastern boundary of this system forms the western boundary of the Káli system, of which the eastern boundary is found in a great ridge descending from the Api peak in Doti of Nepál. It is clear from the above brief description that it is the spurs from the snowy range that bound the river basins, and if we examine further the affluents of each system, we shall see that the ramifications from these spurs form the water-parting between each minor system.

The principal affluents of the Jumna system are the Tons and

Subo inate boundaries. the Giri. The Púbar, Rúpin, and Súpin unite to form the Tons, and are separated

from each other by transverse ridges descending from the great boundary ridge. United they drain a delta-shaped basin having its apex at Kálsi. To the south-west the Giri drains a similarly shaped basin having its apex near Kálsi, and to the east the Jumna drains one having its apex at Kálsi. We find that the point of junction of the apices of these three deltaic basins lies within the Siwálíks, the outer range of geographers, and that the union of these three main affluents forms the Jumna of the plains. This basin, as well as the minor systems within it, is bounded by spurs from the great snowy range or transverse ridges descending from them. If we

further examine the relations of the minor feeders of the three great constituents of the Jumna system, we see that, as a rule, they flow at right angles to their recipients, and that the affluents of these minor feeders obey a similar law. The ridge separating the Tons from the Jumna gives off feeders on the west to the Tons and on the east to the Jumna, at right angles to its direction. From the western slope the Dháragád, Binol, Shaula, and Manjgaon streams flow to the Tons, and from the eastern slope the Ralena, Kútñi, and Silo seek the Jumna. Each of these minor feeders is separated from the other by lateral spurs, descending usually from some peak or knot of peaks, and all, as the veins on a leaf seek the midrib, flow towards the mid-depression and give it their moisture. The two great rivers that unite to form the Ganges are themselves the centres of subordinate systems. The Bhágirathi is divided from the Bhilang by a great ridge descending from the Gangotri group of peaks, whilst a second ridge having a similar origin separates the Bhilang from the Mandákini, an early affluent of the Alaknanda. The Bhágirathi unites with the Bhilang near Tirhi, and the two rivers drain a delta-shaped tract having its apex at Deoprayág. The Sáraswati and Dhauli, which form the head-waters of the Alaknanda, are separated from each other by a ridge of snowy peaks, and its more southern affluents, the Nandákini and the Pindar, are divided from each other by a great ridge descending from Trisúl. The entire basin of the Alaknanda to its junction with the Bhágirathi at Deoprayág is thus a great delta-shaped tract, cut up by the minor feeders into subordinate systems that are bounded by great ridges descending from the snowy range. Between Deoprayág and Hardwár, the Ganges receives from the east the Nayár and the Hinnal, and from the west the Súsua that drains the Dehra Dún. All unite within the Siwálíks to form the Ganges of the plains.

We shall now take up the compact system of the Nayár in southern Garhwál, which at first sight^e would appear to be an exception to the general rules. We find that the boundary ridge that marks its extent is a continuation of the great ridge that, descending from the snowy peak of Nandakot, runs along the left bank of the Pindar to the head of the Lohba valley. Here, as already noticed, this ridge bifurcates; one branch proceeding in the normal direction along

The Nayár.

the left bank of the Alaknanda to Deoprayág, whilst the second branch passes south to Gujargarh and then west to the Ganges at Kharak. These two branches mark the axis of highest elevation in the tract through which they pass. Following the western branch, we have the Dubri peak, 9,862 feet; Dobri peak, 9,862 feet; Gandkhola, 7,553 feet; Devidatta, 7,034 feet; Kankwala, 6,651 feet; Gurdari, 5,893 feet; Jhangarh, 5,878 feet; and a peak about two miles from Deoprayág, 5,030 feet. Following the southern branch through Gujargarh, we find the second, Duda-ka-toli peak with an elevation of 10,180 feet above the level of the sea; Barma-dúngi, 9,190 feet; Nandatopa, 8,086 feet; Khamek, 7,152 feet; Gujargarh, 7,969 feet; Khatti, 8,270 feet; Utain, 6,901 feet; Bukrari, 6,267 feet; and Naugarh, about a mile above Kharak, 6,065 feet. Close to the point of bifurcation, the boundary ridge sends a lateral spur southwards, that divides the basin of the Nayár into two parts, that drained by the western Nayár or Chhiphalghát river, and that drained by the eastern Nayár or Kainyúr stream. This great spur preserves the superior elevation of the boundary ridge, and is marked by the following peaks:—Barári, 8,499 feet; Bandani, 8,278 feet; Panjing, 8,810 feet; Devitank, 8,849 feet; Matikhál, 7,688 feet; and Chhatargah, 6,790 feet, when it is lost in the valley of the eastern Nayár near Kandui. Short ramifications from this spur or the boundary ridge itself determine the course of the minor feeders of the two great channels of the Nayár system. Thus a transverse ridge from Dubri, marked by the Tara-ka-kand (9,000 feet) and Banjkot (8,203 feet) peaks, divides the Chhiphalghát river near its source from its feeder at Paitháni on the left bank, whilst other ridges from Gandkhola, Devidatta, and other peaks, separate the other feeders the one from the other. A similar rule obtains along the course of the eastern branch, and we thus see that there is no real difference in principle between the arrangement of the drainage system of this apparently abnormal minor basin and that of the other greater systems already noticed.

In the system of the western Rámanga, however, we have an arrangement for which we find an analogue in that of the Bágmati in Nepál. As we have already seen, the basin of the Rámanga is in shape a great

Western Rámanga.

obtuse-angled triangle, with its apex towards the snowy range and its base towards the plains, thus filling up the gap between those systems that have their apices towards the plains and their bases towards the snowy range. Of its affluents, the Kosi alone has a considerable course within the hills, the remainder having their origin in or close above the elevated tract lying along the foot of the hills, and joining the Rānganga at some distance southwards in the plains. Here, although the main boundary ridges are still the spurs descending from the snowy range, the several streams do not unite within the outer range, but like the Bāgmati and its affluents, well beyond it in the plains. The Kāli is known as the Yānkti near its source, as the Kāli during the greater portion of

Kāli.

its course through the hills, as the Chauka or Sārda in the Bhābar and Tarāi, and as

the Sarju and Ghogra in Oudh to its junction with the Ganges, to the south of the Ghāzipur district. The Kāli basin is bounded on the west by that of the Ganges, and on the south-west by that of the western Rānganga. It receives from the west the Gori, Sarju, and Ladhiya, and from the east some small streams from Nepāl, all of which unite within the hills to form the Sārda at Barmdeo. The Sarju is divided from its affluents, the eastern Rānganga, by a great meridional ridge, extending from their sources to their junction, whilst a second ridge, running in a south-easterly direction from the same group of peaks to Askot, separates the latter river from the Gori. The great Pancha-chuli range, running south-east from the line of water-parting, divides the basin of the Gori from that of the Dārma Yānkti, and a similar snowy range having a like origin separates the Dārma Yānkti from the Kutli Yānkti. The great mass of peaks comprising Trisul, Nanda Devi, and Nanda-kot thus send forth great boundary ridges from their entire southern face to the west between the Rānganga and Nandākini: to the south-west between the Nandākini and the Pindar, and between the Pindar and the Sarju; to the south between the Sarju and the eastern Rānganga; and to the south-east between the last river and the Gori. These indications are sufficient to mark the salient features of this portion of the Himālaya, and we reserve the details for the Gazetteer articles of this notice.

The mountain system lying between the snowy range and the plains may therefore be said to consist of
 The snowy range. an outer range parallel to the snowy range and connected with it at wide intervals where it meets the great ridges descending therefrom. These latter bound the river basins, and the ramifications from them determine the course of the minor feeders of each system. We have now to consider the snowy range itself, and in these provinces at least it is a well-marked feature, lying from ten to twenty miles to the south of the Indian water-parting. The line of snowy peaks seen from Naini Tál and Masúri all lie to the south of the passes by which travellers cross into Tibet. The route by Nilang in foreign Garhwál through the Jádth valley crosses north of the Jamnotri group; that by Mána in British Garhwál lies to the north of the Kedárnáth group; that by Milam in Kumaon behind the Nanda Devi group; and that by the Lanpiya-dhúra, also in Kumaon, behind the Panchachúli group. The Jádth Ganga runs in a valley parallel to the snowy range and the line of water-parting, at an elevation of 15,000 feet above the level of the sea, near its source at Sumla, but gradually falling to below 9,000 feet at its junction with the Bhágirathi, near Bhairongháti. The horizontal distance between these two points is a little over 17 miles, and the fall is therefore over 350 feet in a mile. The Vishnuganga or Sáraswati descends from the Mana Pass at 18,000 feet above the level of the sea to Vishnuprayág, a distance of some 35 miles, where it is little over 5,000 feet. Here we have an average fall of about 370 feet in the mile. The increase in elevation from the place where the river valleys enter the area of greatest elevation is equally marked in the valleys of the Dhauli, the Gori, and the Dárma river. Dhárchúla on the Káli, before the line of snowy peaks is reached, is only 2,750 feet above the level of the sea; whilst Golam Ja, to the north and above that line, and about 1,500 to 2,000 feet above the bed of the Káli, is 8,000 feet above the level of the sea. The bed of the Káli at Changru is 10,000 feet, and in the twenty-five miles between it and the Lanpiya-dhúra Pass the fall is over 8,000 feet, giving an average fall of 400 feet to the mile. These facts well illustrate the law that the river beds¹ to within a distance of ten miles in a direct line from the snowy peaks seldom

¹ See page 8.

exhibit a rise of more than four or five thousand feet ; but when we cross the line on which the great peaks are situated, the ascent very rapidly increases, and a very few miles carries the river-bed up to an altitude of nine or ten thousand feet, thus showing that the sudden increase of height of the mountains along this line is not confined to the peaks alone, but is a general elevation of the whole surface.

This sudden and steady rise in elevation when the line of snowy peaks is reached appears to be a well-marked feature of the entire Himálayan range. The snowy peaks, however, do not occur in a continuous ridge, but in masses separated the one from the other by deep depressions that form the line of drainage for all the surplus moisture of the tract between the snowy range and the line of water-parting to the north. These depressions are, so far as we know, a more distinctive feature of the Indian than of the Turkish slope of the Himálaya-Tibetan mass, a fact doubtless due to the greater rainfall received by the former. The influence of the monsoon on the southern slope is shown in its supporting a more dense and varied vegetation, and even on individual ranges and hills the southern exposure is similarly strongly marked in Kumaon. It is the ceaseless action of water that has furrowed out these valleys from the great mass, and naturally this has been accomplished on a greater scale and with more results along the southern slope that receives the full force of the periodical rains. The mass of peaks known as the Gangotri, Kedárnáth, and Badrináth groups, are separated from the next group to the east by the valley of the Sáruswati, and this group again from the Nanda Devi mass by the Dhauli river. The last is again divided from the Panchachúli group by the valley of the Gori, and the latter from the succeeding group by the Dárma valley. All these valleys are inhabited and cultivated during the summer and rains, and some of them are comparatively of considerable extent. The masses of snowy peaks are more like the terminal ends of huge spurs descending from the line of water-parting than a continuous ridge. They, however, occur in regular sequence along the entire line of the Himálaya, and, as seen from the plains, have the appearance of a connected chain. The following table, showing the principal peaks that occur throughout some eleven degrees of longitude, will give some idea of their number and importance.

*Points on the Hindūlaya mountains arranged in order of longitude by
Mr. Trevelyan Saunders, from the records of the G. T. S.*

Number.	Name of peak.	Latitude.	Longitude.	Altitude.
I.	Chinnalari	27° 19' 37"	89° 18' 13"	23,911
II.	Gipnachi	27° 16' 23"	88° 56' 37"	11,518
III.	Pacham (Dunkar)	27° 56' 52"	88° 53' 5"	23,186
IV.	Chumundin (Chob)	27° 27' 28"	88° 49' 38"	17,325
V.	Black rock (Guatam)	27° 31' 7"	88° 48' 39"	17,572
VI.	Narsing...	27° 39' 36"	88° 19' 28"	19,116
VII.	Pandun	27° 31' 31"	88° 15' 35"	22,017
VIII.	Kanchanjunga	27° 11' 26"	88° 11' 50"	27,815
IX.	"	27° 12' 5"	88° 11' 26"	28,156
X.	Katun	27° 36' 26"	88° 9' 15"	24,015
XI.	Jannu	27° 10' 52"	88° 5' 13"	25,301
XII.	"	27° 39' 55"	87° 7' 56"	...
XIII.	Sihpur	27° 53' 18"	87° 7' 51"	27,799
XIV.	Chamlang E.	27° 16' 27"	87° 1' 21"	21,020
XV.	Mount Everest	27° 59' 12"	86° 58' 6"	29,002
XVI.	Chamlang W.	27° 15' 16"	86° 51' 56"	22,215
XVII.	Sankosi	27° 15' 11"	86° 39' 57"	22,826
XVIII.	"	27° 52' 16"	86° 31' 57"	21,987
XIX.	"	27° 8' 13"	86° 28' 32"	23,570
XX.	"	27° 57' 18"	86° 22' 12"	23,117
XXI.	"	27° 57' 21"	86° 9' 8"	19,560
XXII.	Jalajil iya	28° 7' 36"	85° 51' 42"	21,853
XXIII.	"	28° 21' 3"	85° 49' 21"	26,305
XXIV.	"	28° 10' 20"	85° 48' 17"	22,891
XXV.	Dayabang	28° 15' 17"	85° 33' 35"	23,760
XXVI.	Akn	28° 23' 25"	85° 10' 12"	21,317
XXVII.	"	28° 20' 38"	85° 7' 21"	23,013
XXVIII.	Yassa	28° 25' 58"	84° 11' 0"	25,818
XXIX.	"	28° 30' 7"	84° 36' 31"	25,729
XXX.	"	28° 32' 55"	84° 36' 9"	26,680
XXXI.	"	28° 39' 39"	84° 21' 56"	...
XXXII.	"	28° 39' 21"	84° 20' 32"	...
XXXIII.	Barathor	28° 29' 18"	84° 13' 56"	22,917
XXXIV.	"	28° 32' 0"	84° 9' 52"	26,069
XXXV.	"	28° 32' 6"	84° 7' 32"	21,718
XXXVI.	Morshiadi	28° 31' 58"	84° 1' 57"	24,780
XXXVII.	"	28° 29' 37"	83° 59' 22"	22,961
XXXVIII.	"	28° 29' 48"	83° 59' 20"	22,986
XXXIX.	"	28° 35' 38"	83° 51' 16"	26,522
XL.	"	28° 39' 59"	83° 50' 55"	23,611
XLI.	"	28° 39' 11"	83° 46' 22"	22,171
XLII.	Dhaulagiri	28° 41' 43"	83° 32' 9"	26,826
XLIII.	Karāyani	28° 45' 39"	83° 25' 52"	25,156
XLIV.	"	28° 15' 8"	83° 25' 12"	25,299
XLV.	"	28° 43' 56"	83° 21' 18"	21,912
XLVI.	"	28° 41' 2"	83° 21' 20"	25,095
XLVII.	"	28° 40' 21"	83° 19' 6"	23,565
XLVIII.	"	28° 43' 49"	83° 15' 9"	21,141
XLIX.	"	28° 44' 47"	83° 11' 18"	23,779
L.	"	28° 44' 30"	83° 9' 29"	21,727
LI.	"	28° 45' 53"	83° 8' 27"	21,472
LII.	Chambisi	28° 49' 33"	82° 39' 33"	19,415
LIII.	Api	29° 59' 7"	80° 56' 22"	...
LIV.	Parlachachuli	30° 12' 51"	80° 28' 9"	22,673
LV.	"	30° 15' 12"	80° 25' 6"	21,171

Points on the Himálaya mountains arranged in order of longitude by Mr. Trevelney Saunders from the records of the G. T. S.—
(concluded.)


Number.	Name of peak.	Latitude.	Longitude.	Altitude.
LVI.	Nandakot ...	30° 16' 51"	80° 6' 39"	22,538
LVII.	Nanda Devi ...	30 21 58	80 9 31	24,417
LVIII.	Nanda Devi ...	30 22 31	80 0 50	25,661
LIX.	Do. ...	30 22 35	80 0 46	25,587
LX.	East Trisul ...	30 16 14	79 54 51	22,342
LXI.	Do. ...	30 30 56	79 54 31	23,092
LXII.	West Trisul ...	30 18 43	79 49 7	23,382
LXIII.	Nandakua ...	30 22 9	79 45 40	19,916
LXIV.	Do. ...	30 20 56	79 45 36	20,722
LXV.	Do. ...	30 20 57	79 45 35	20,773
LXVI.	Do. ...	30 41 6	79 44 53	22,093
LXVII.	Kamet (Ibn Gamin) ...	30 55 13	79 38 4	25,373
LXVIII.	Nilakanta ...	30 43 52	79 26 56	21,661
LXIX.	Badrinath ...	30 44 16	79 19 20	23,210
LXX.	Do. ...	30 43 22	79 17 52	22,511
LXXI.	Do. ...	30 46 44	79 16 58	22,347
LXXII.	Kedarnath ...	30 47 53	79 6 34	22,790
LXXIII.	Tharlasagar ...	30 51 40	79 2 14	22,582
LXXIV.	Do. ...	30 51 41	79 2 13	22,628
LXXV.	Jauli ...	30 51 18	78 53 53	21,672
LXXVI.	Bas or Sri Kanta ...	30 57 25	78 50 50	20,149
LXXVII.	Bandarpouchh ...	31 0 12	78 35 45	20,758
LXXVIII.	Jannotri ...	31 0 25	78 34 6	20,038
LXXIX.	Swargarufi ...	31 0 8	78 32 32	20,405

After crossing the line of water-parting which, as we have seen, lies at the valley heads to the north of the Tibetan plateau, line of snowy peaks, we come upon the great Tibetan plateau which has a mean elevation of from 13,000 to 17,000 feet above the level of the sea. Puling on the plateau by the Nilang route is 13,800 feet, Chiurkung on the Mana route to Totling is 15,700 feet, Chitung-dhar on the Unta Dhura route to the Satlaj is 15,810 feet, and Buljuing near the head-waters of the Karnali is 15,850 feet. Rabgyaling is 14,000; Gartokh, 14,240 feet; Tirthapuri, 14,820 feet; and lake Rakas Tal over 15,000 feet above the level of the sea. Captain H. Strachey has described the Tibetan tableland lying between the Indian and Turkish water-partings "as the flat top of a great embankment exhibited in all its thickness in the scarp of the Indian Himálaya: the summit, though deeply corrugated with valleys and mountains in detail, being in its general relief laid out horizontally at a

¹ On the physical geography of Western Tibet: London, 1854.

height little inferior to that of its southern scarp." Although the highest summits yet known and measured lie along the Indian slope, very lofty peaks have been seen in all parts of the interior, "and the passes which must be crossed to get from one Tibetan valley to another, even in the very central axis of drainage, generally equal those by which Tibet is reached from India:" so that, on the whole, Captain Strachey was of opinion that the medial depression is but faintly marked in the beds of the great rivers without much affecting the mean elevation of the mass. His description of the mountain system of that portion of Tibet lying to the north and west of Kuumon is the best that we possess, and will usefully conclude our brief review of the features of this portion of the Himálaya-Tibetan mass. "The mountains that compose the bulk of West Nári are not easily understood or defined. On ascending the highest passes we can seldom see anything but a contracted view of mountain tops on all sides, looking very like chaos: no general view of ranges under our feet is ever obtainable as the passes naturally select the ravine-heads and lowest points of the ridge which are not only flanked but often almost surrounded by the higher summits; and the valleys are commonly so steep and narrow, especially in the Rong country, that the view can hardly ever penetrate to an alluvial bottom and the sight of any inhabited place from a pass top is most unusual. When travelling along the bottoms of the valleys, we generally see nothing but a narrow tortuous passage between steep rocky walls, shutting out all extended view, and rather concealing than exhibiting the mountain ranges of which they form but the lowest outworks; consequently it is only by an extended series of observations and inferences, joined and assisted by maps, that any regular arrangement of these mountains can be distinctly established, and my account of them is liable to error in proportion to the defects of my own map. The general plan of the mountain system appears to me to consist of a series of parallel ranges running right across the breadth of the tableland in a direction so extremely oblique to the general extension of the whole as often to confound the one with the other, or to convert the transverse direction to a longitudinal one. The annexed figure may help to explain this. Short transverse necks connecting the main ranges in some parts, and cross fissures cutting through

them in others, together with projecting spurs of a secondary order, will suffice to convert the supposed primary arrangements into all the existing varieties of valley and drainage.



Such connecting necks, when above 18,000 feet, become more or less confounded with the main ranges, and, if not above 17,000 feet, often appear as low watersheds, just dividing the heads of two valleys lying in one line, but draining opposite ways. Secondary spurs also may be so high and so obliquely joined to the primary ranges as to make it difficult to distinguish between the two ; and the cross fissures may sometimes admit a main river to pass through a main mountain mass, in which case the continuity of the range is often evidenced by the extreme narrowness of the rocky gorge or height steepness, and geological correspondence of its sides. Much of the Indian watershed seems to be formed in this way, the great snowy peaks lying mostly on the terminal butt-ends of the primary ranges, sometimes widened by lateral spurs, and the Tibetan passes crossing the low connecting links, whose alignment forms the main watershed, but not the main mountain crest."

It seems strange that so late as the year 1817 the occurrence of glaciers in the Himalaya was considered a matter of doubt by the learned in Europe. There is now no fact more widely attested

and more thoroughly established than the
 Glaciers. existence of glaciers at the head of almost

every valley that descends from the ranges covered with perpetual snow. In size and importance they also fitly compare with those stupendous peaks around them that have placed the Himalaya in the foremost rank of all the mountains of the earth. Colonel Gordon gives us¹ illustrations and descriptions of the great glaciers

¹ Roof of the World, 17.

met by the Yárkand Mission on the journey between Leh and the Kárákoram pass. He mentions the lower Kumdan glacier that comes from the high peaks to the north-west, and continues down the right bank of the stream for over two miles, "forming a perfect wall of ice, rising from the water about 120 feet, and showing a surface covered with countless pinnacles and points." The Remu glacier, also seen by Colonel Gordon, rises amongst peaks and ridges from 19,000 to 24,000 feet high. "It is about 21 miles in length and from one to one and three-quarters mile broad, terminating at an elevation of 15,800 feet above the level of the sea, with a width of about three miles of gigantic cliffs of ice fully 250 feet high." He adds:—"the glaciers of the western Himálaya are twice as extensive as those of the Alps, and are probably the largest in the world, or at all events larger than any others out of the polar regions. One in the Muztágh range is believed to be 34 miles long with fifteen distinct moraines; while in its immediate vicinity is another, 31 miles in length, which may be said to join with it in making 65 miles of continuous ice." Other glaciers have been described by travellers in the ranges between Ladák and Garhwál. In the tract with which we are more immediately concerned we have glaciers at the head of the Jádh Ganga, the Bhágirathi, Vishnu Ganga, Dhauli, Kailganga, Sundardhúnga, Pindar, Kuphini, eastern Rám-ganga, Gori, and Dárma Yáukti. Lieutenant Weller¹ in his visit to the source of the Gori near Milam, describes that river as coming "out in a small but impetuous stream: at the foot of apparently a mass of dirt and gravel some 300 feet high, shaped like a halfmoon. This is in reality a mass of dark-coloured (bottle-green) ice, extending westward to a great distance, and covered with stones and fragments of rock which in fact form a succession of small hills." Here and there were circular and irregularly shaped craters (as it were) from 50 to 500 feet in diameter at top, and some of them 150 feet deep, and higher up these gave place to narrow fissures. This glacier is between seven and eight miles long, and terminates at an elevation of 11,600 feet above the sea. The glacier of the Pindar in Kumaon is the one, however, regarding which we have the most complete information, and we shall confine ourselves, therefore, to its description in detail.


¹ J., A. S., Ben, XI., 1166.

The Pindar river¹ is an affluent of the Alaknanda, and has its rise in a glacier to the west of the Nandakot peak and its valley, where the glacier ends, is about a mile across between the precipitous mountains that bound it. From the foot of the rocks on either side the bottom of the valley slopes inwards with a moderate inclination, leaving in the middle a hollow about 300 yards wide and 250 feet deep, with very steep banks, at the bottom of which flows the river. This comparatively level space between the precipices on either side and the river bed is observable for a mile or more below the end of the glacier, though much cut up by watercourses. The glacier occupies about two-thirds of the whole breadth of the head of this valley, leaving between itself and the cliffs on the east an open grassy slope, which extends along the foot of the moraine for upwards of a mile and a half above the source of the river, and which seems to be a continuation of the level space before mentioned. The first appearance of the glacier is thus described :—"It seems to be a vast rounded mass of rocks and ground utterly devoid of any sign of vegetation, standing up out of a grassy valley. From the foot of its nearer extremity the river, even here unfordable, rushes in a turbid torrent out of a sort of cave ; the top of which is but a few feet above the surface of the water (May). The end, immediately over the source of the river, is very steep and of a dull black colour. It is considerably fissured, the rents appearing to arise from the lower parts, tearing themselves from the upper by their own weight. On a closer examination this abrupt end proves to be a surface of ice covered with sand and gravel and curiously striped by the channel made by the water that runs down it as it melts. Behind this, the glacier rises less steeply, like a bare gravel hill, to its full height, which is probably about 500 feet above the water of the river when it leaves the cave. In some places, however, are seen great fissures both vertical and horizontal, the latter evidently made by the separation of regularly stratified layers."

The glacier is formed by the meeting of two ice streams from
 Feeders of the glacier. gorges, one coming from the north-west and
 the other nearly from the east, and which

¹ From "A description of the glaciers of the Pindar and Kupluni rivers in the Kumaon Himalaya," by Lieutenant R. Strachey, Ben. Eng., J., A. S. B., XVI. 794, and "Note on the motion of the glacier of the Pindar in Kumaon" by the same. *Ibid.*, XVII., ii., 203, and given as nearly as possible in the writer's words as the only scientific examination of these glaciers that we possess.

GLACIER OF THE PINDAR



meet about two miles above the source of the river, as shown in the accompanying sketch.

The feeder from the north-west is larger than that from the east, and its surface is at a considerably higher level for some hundred yards below their first junction. It descends with a great inclination, entirely filling the gorge, down which it comes in a cascade of ice. It assumes the general appearance of a confused mass of irregular steps, which are again broken up transversely into peaks of every shape. The west side of this cascade continues nearly in its original direction after having passed the point below which the glacier bends sharply to the south-west, and in this way completely crosses the glacier. The steps in which it falls, however, also gradually change their direction so as to remain nearly perpendicular to the general current of ice. The transition to the regular level ice is very sudden, and begins much higher up on the west than on the east side. Near the foot of this ice-fall the steps were observed to have their tops considerably overhanging. A small tributary, also descending in cliffs of ice, joins the main glacier from a ravine on the east, not far above where it takes the sudden bend. The feeder from the east is formed by the union of two smaller glaciers, one from the north-east and the other from the south-east, which is the larger of the two. The north-eastern tributary appeared to have no very steep inclination, but was considerably broken up at its junction with the other. Another small glacier joins the main one from the north-west, a short distance below the point where it bends southwards. Its inclination is very great, but it perfectly maintains its continuity of structure to the bottom.

The lateral moraine of the west side of the northern branch of the glacier is first seen near the bend, where
Moraines. it shows itself as a black band along the edge of the ice which in other parts of the fall is quite white. The moraine is small between the bend and the tributary glacier below it, but very rapidly increases, and in its lower parts is a chaos of desolation. This great addition to the size of the moraine is owing to the quantity of *débris* brought down by the small glacier. The ice below the junction is much broken up by crevasses, and rocks and gravel from the moraines on both sides of the tributary glacier are scattered over the space between them, and the moraines at

first sight seem to lose their distinct form. Although there is no clear ice between the moraine that originates on the east of the tributary and the west side of the glacier, the identity of that moraine is sufficiently marked by its colour and by the regular rise above the general surface of the glacier of its top, which remains tolerably even for some way down, being beyond the limit of the disturbance caused by the crevasses along the edge of the glacier. About half way down to the lower end of the glacier, however, the full action of these crevasses reaches the whole of the moraine, and it is scattered or lost sight of in the general confusion of surface. An epoch of peculiar destructiveness to the mountains is marked in one part of this moraine by the accumulation of huge masses of rock from 20 to 30 feet square and as much as 15 feet high, and the stone found on it are generally larger than those on any of the other moraines. The true west lateral moraine, below the tributary glacier, is not very large, nor is its top much elevated above the bottom of the valley, excepting quite at its end.

The lateral moraine of the south-eastern side of the glacier is very large. Its top rises, on an average, probably 250 feet above the bottom of the valley. Along its foot runs a stream, gradually increasing in size, that collects the open drainage of the outer slopes of the moraine. The lower part of this slope is a mass of loose stones and earthy gravel which rolls down from above, as the face of the ice which is visible in the upper fifty or sixty feet of the slope melts and recedes, a process that is constantly going on. On the inner side the slope of the moraine is thirty or forty feet above the level of the clear ice of the glacier. The upper part of the moraine comes down nearly straight from the point where it meets the foot of the north moraine of the east glacier. The north branch glacier being considerably higher than the eastern, the moraine slopes down from the bed of the former to that of the latter, forming a deep angular depression where they meet, that gradually diminishes in depth up to the top of this glacier, which is here entirely covered with *débris*, the moraines of its two sides being scattered all over it for some distance above its union with the north or main branch. The resulting appearance is that the northern branch runs over the eastern, or that the latter runs into the former and is absorbed by it. The eastern tributary brings



From a photograph by Secker

View of the Big Island of Hawaii

down moraines that spread over the whole of its breadth at its extremity. Besides these lateral moraines there is a medial one that is first seen as a dirty stripe along the ice cliffs of the fall at the head of the north glacier. As it comes down the level ice it gradually begins to assume the decided appearance of a moraine, and increasing by degrees at last becomes very large. It continues in a well-defined form for some short distance beyond where the western moraine is dispersed, but there it also is scattered over the ice, and the two become blended together, and ultimately extend to meet the *débris*, which is similarly dispersed by the eastern moraine from the opposite site of the glacier. The whole of the moraines, in the middle of the length of the glacier where it is most regular, are very considerably raised above the general surface of the ice, which in some parts may be as much as one hundred feet below the tops of the western and medial moraines. It would appear that this great elevation is not so much due to the accumulation of *débris* as to the protection afforded by the superincumbent rubbish to the ice below which prevents its being melted. The clear ice beyond the moraine is constantly depressed where exposed, and on the very tops of the moraines pure ice was often seen hardly covered by stones. The protection afforded by the lateral moraines raises the sides of the glacier so much that a very considerable hollow is caused in its middle, which is a striking feature in the first appearance of its lower extremity.

The ice of which the glacier is composed is perfectly pure and clear, but where seen in considerable masses, *Crevasses ; glacier tables.* stripes of a darker and lighter bluish green are distinctly visible. It is composed of bands of ice containing small air bubbles, alternating with others quite free from them. In many places the surface presents a striated appearance, arising from the different degrees of compactness of these differently coloured bands and their consequently different rates of melting. The direction of these coloured views as seen in *crevasses* showed a dip inwards or towards the longitudinal axis and a dip upwards or towards the origin of the glacier in every part, the stratification being more perpendicular towards the head and more nearly horizontal in the lower parts. The direction of the strata in place was also very clearly marked in many parts of the ice, and was placed in

curves, having their branches nearly parallel to the sides of the glacier and their apices directed downwards, the curvature in the centre not being at all sudden. No dirt bands were observed. The crevasses were neither very numerous nor very formidable. They are developed across the direction of the glacier's length on both of its sides, commencing from the small tributary on the west side and from the union of the eastern glacier on the other, and continuing almost to the end, those on the west side being, perhaps, the larger. They are generally wider towards the edges of the glacier, closing up as they approach the centre. They are nearly vertical, and are directed from the sides upwards or towards the head of the glacier, those on the west bearing nearly east and west, and those on the east bearing nearly north and south, thus forming angles of about 45° with the axis of the glacier. Many pools of water (*baignoirs*) were seen on the surface of the ice; some of the largest were said by the guides, who are in the habit of visiting the glacier, to be found in the same place every year. The clear surface of the ice everywhere assumes a more or less undulating appearance from the action of the water that drains from it as it melts and the small streams, into which the drainage collects, end by falling into some of the crevasses. The remains of the last winter's snow was hardly perceptible on any part of the glacier. The occurrence of stones standing up on bases of ice (glacier tables) above the general surface of the glacier is common, but those seen were small. The rocks below the bend in the north-western glacier were covered with grooves or scratches, sloping in about the same direction as the surface of the ice at the spot. These grooves extend to twenty or thirty feet above the present level of the glacier. Almost in every place a space was left between the rock and ice, the latter appearing to shrink from contact with the former, due doubtless to the heat of the rock melting the ice.

The Kuphni river, that rises on the side of the Nanda-kot peak, opposite to the Pindar river, has also its source in a glacier. Both rivers unite at Dwāli, about eight miles from the end of the Pindar glacier and about six miles from the end of that of the Kuphni. General Strachey examined the Kuphni glacier also, and describes the valley for a mile or two below the end of the glacier as having

very much the same general character as that of the Pindar, but somewhat more rugged and desolate in appearance. The glacier commences about two miles above the source of the river and fills the whole breadth of the valley, which is about three-quarters of a mile broad in its upper part. The glacier begins in a precipitous fall of ice some sixty or seventy feet high, which, however, still exhibits the ribbon-like structure. From the foot of the fall the surface was very even, though the slope was still considerable. The main glacier of the Kuphini is joined by two small tributaries on the east and by one on the west, all of which are highly inclined and bring down considerable quantities of *débris*. The moraines are confined to the sides of the glacier, though many small stones are scattered over every part of the ice. As was observed on the Pindar, the protection given by the lateral moraines to the underlying ice leads to the promotion of a medial depression in the glacier at its end. The crevasses here, too, are most strongly marked near the sides and are inclined at an angle of about 45° from the longitudinal axis downwards. The structure of the ice was in all respects similar to that found on the Pindar. On the interesting question of the extension of glaciers at a remote period the inquiries of General Strachey give no precise information. He, however, considers that "some very decided change in the state of things is certainly indicated by the long plateaus before mentioned running for a mile or two below the present terminations of both glaciers nearly parallel to the rivers, but several hundred feet above them." He considers it "impossible that these level banks above the rivers have been caused by deposits from the ravines in the sides of the valleys, for such deposits would have had very irregular surfaces, and indeed their present effect in destroying the regularity of the plateaus is everywhere visible. Had the same appearance been noticed in any other part of the river's course, it would at once have been attributed to the action of water at some former period, and it would have been supposed that the bed had afterwards been excavated to its present depth. If this was the case, the glaciers which the plateau was forming must either have terminated considerably higher up the valleys or have stood altogether at a much higher level. In either of these ways the water could have been delivered at a level sufficiently high to

form the plateau. But it may admit of doubt whether the quantity of water in the rivers, as they are at present, is sufficient to account for such an extent of level deposit or for such a depth of erosion of their beds; for at this great elevation they are not subject to those violent floods that occur lower down, and for nearly half the year too they are inert. The only other way of accounting for the appearance is that it has been occasioned by an extension of the glacier, and that the level top of the plateau shows the limit to which the tops of the moraines reached, as the glacier gradually receded." We have referred on a previous page¹ to the existence of evidence of glacial action far below the present limits of glaciers, and to those who wish to pursue the subject further we commend the records of the Geological Survey and the summary² in the recently published 'Manual of the Geology of India.'

General Strachey has rendered us another important service in his observations on the motion of the Pin-dari glacier recorded in May, 1848. His procedure is thus stated. About 200 yards below the small tributary that enters the main glacier from the north-west a moraine was found heaped up against an almost perpendicular wall of rock, and sufficiently high to command a view of the greater part of the surface of the glacier along the line on which observations were to be made. This line, which is nearly perpendicular to the general direction of the glacier, was marked by two crosses painted white, one on the rock in contact with the old moraine and one on a cliff on the opposite side of the valley. A stake was driven into the moraine at its highest point, close to the rock, on the line between the two crosses, and a theodolite was set up over it. Five other marks were also made on the glacier at intervals along the same line by fixing stakes in holes driven in the ice with a jumper. These marks, which were all carefully placed on the exact line between the crosses by means of the theodolite, were completed at about 0h. 30m. P. M. on the 21st May. On the following day the theodolite was again set up on the same place as before, and being properly adjusted, the cross-wires of the telescope were directed to the cross on the cliff on the opposite side of the glacier. A stick was then set up near the first of the five marks that had been made the

¹ P.² Pp. lxx., 372, 373, 536, 639.

previous day, and was, by means of signals, moved up or down the glacier, till it appeared to coincide exactly with the cross-wires of the telescope, and consequently to be exactly on the line between the two crosses painted on the cliffs. The distance between the centre of the stick and that of the fixed mark was then measured, which evidently showed the downward progress of the ice at that point of the glacier, since the marks were made the day before. The same proceduro was followed at each of the marks. The results were as follows :—

Time of observation.	<i>Distances of fixed marks from standard line.</i>				
	On the west moraine.	On the medial moraine.	At east foot of medial moraine.	Near the middle of the clear ice.	On the eastern moraine.
	ft. in.	ft. in.	ft. in.	ft. in.	ft. in.
21st May h. m.					
21st May 0 30 p. m.	0 ...	1 ...	1 ...	1 ...	0 ...
22nd " 1 15 p. m.	0 5½	1 0½	1 0	1 0½	0 6½
25th " 8 45 a. m.	1 9½	2 9½	2 11½	3 1	1 5½

The motion in 24 hours of the several marks will also be found to be—

Date.	<i>Mean motion (in inches) of ice in 24 hours.</i>					
	On the west moraine.	On the medial moraine.	At east foot of medial moraine.	Near the middle of the clear ice.	On the east moraine.	Approximate mean temperature.
21st to 22nd May ...	5.8	11.9	11.6	11.9	6.5	49°F.
22nd „ 25th „ ...	5.7	7.6	8.4	8.8	3.8	38°F.
General mean ...	5.5	9.7	10.0	10.8	5.1	...

The progress of the lower extremity of the glacier was likewise approximately measured by observing the apparent angular motion of a pole fixed on the top of the eastern moraine, and of a conspicuous

rock lying not far from the middle of the glacier. The results of these observations were—

Date.	Mean motion (in inches) of ice in 24 hours.	
	On the moraine.	Near middle of the glacier.
19th to 20th May	3.0	...
20th „ 23rd „	6.2	8.1
23rd „ 25th „	5.3	10.8
General mean	4.8	9.4

A comparison of the motion of the upper and lower parts of the glacier gave on the lateral moraines 4.8 inches as the mean motion of the ice in 24 hours in the lower part of the glacier, and 5.3 inches in the upper part, and in the middle of the glacier 9.4 inches as the mean motion for the lower part, and 10 inches for the upper part of the glacier. The elevation of the foot of the glacier at the source of the Pindar is 11,929 feet, and of the theodolite station, where these observations were made, was 12,946 feet above the sea.

We have made mention of the snow-line or limit of perpetual snow,¹ which has, at times, given rise to considerable discussion. The height at which precipitations of vapour fall to the earth's surface as snow varies with the temperature of each particular place, and with the annual or even diurnal oscillations of the temperature. In Bhután and Sikkim the ordinary winter limit of snow is about 6,000 feet, and it is rare, says Dr. Hooker, for even three inches to remain on the ground as many days at 7,000 feet.² According to General Strachey, the height at which snow is certain to fall in Kumaon is about 6,500 feet, and at an elevation of 5,000 feet it will not fail more than one year out of ten. The lowest level to which sporadic falls of snow are here known to descend is about 2,500 feet, of which there are two authentic instances on record since the British occupation, the first in 1817 and the second in 1847. In the valley of Kashmir, at an elevation of 5,500 feet, the snow falls every year,³

¹ These observations on the snow-line are based on the article in J., A. S., Ben., XVIII., i. 267, entitled, "On the snow-line in the Himalaya," by Lieutenant R. Strachey, Engineers, on notes placed at my disposal by the same writer, and on the works of recent travellers.

² Hooker, Quar. J., Hort. Soc., VII., 144;

³ Moorcroft, II., 107.

Griffith Post. Papers, I., 236.

and further west as low as 4,000 feet, whilst at least one fall of snow is recorded at Peshawar, which has an elevation of only 1,250 feet above the level of the sea.¹ Campaigning experiences during the late war show that the winter snows descend to a very low elevation in the valley of the Kábul river and at Kandahár. If we follow the lower boundary of the winter snow on a mountain as it melts with the advance of summer, we at length ascend to a height at which the summer influence is insufficient to entirely melt the snow, and from which, as the season advances towards winter, we have gradually to descend in order to follow the line of snow. The line to which the snow recedes, and from which it again advances in one complete revolution of the seasons, is called the snow-line. The snow above that line is called perpetual snow, not as observed by Professor Forbes, that the continuance of snow at any spot implies that it never melts there, but only that some always remains unmelted.

According to our best authorities, the height of the snow-line on the most southern exposures of the eastern Himálaya. Observations in the Himálayan slope that carry perpetual snow ranges from 15,000 to 16,000 feet all along that part of the chain that lies between Sikkim and the Indus, whilst to the north towards Tibet it has a considerably higher elevation. Before proceeding further we may observe, with General Strachey, that "all estimates of the snow-line are, in the very nature of things, subject to no little uncertainty; for, independently of the variations of the seasons from year to year there are naturally considerable differences in the level at which the snow lies on steep or slight slopes and on north or south exposures, between the latter of which a difference of as much as a thousand feet may at times be observed. Besides this, too, there is some practical difficulty in the actual observation of the snow-line, for the process of judging by the eye whether the snow upon one's path and still more on contiguous mountain sides begins to exceed the bare spaces is neither easy nor susceptible of much precision. Hence the errors and uncertainty to be looked for in all our conclusions must be considerable, amounting no doubt to several hundred² feet." Dr. Hooker estimated the height of the snow-line on the most southern spurs of the snowy mountains in Sikkim to be at about 15,500

¹ Bom. Geog. Jour., X., 39.

² Captain H. Strachey, J. R. G. S., XXIII. 69.

feet.¹ Of the peaks covered by perpetual snow, the elevation of which is noted in Dr. Hooker's map, Chola, on the boundary between Sikkim and Bhután, is the lowest, (17,300 feet), and at the same time the most southern and the Chola Pass immediately to the south of the peak, and rising to 14,900 feet, he found to be free from snow at the beginning of November. Somewhat further to the north, near Younghong, the lower limit of perpetual snow was directly measured in September and found to be nearly 16,000 feet. To the west of Kanchanjunga in eastern Nepál, the south-eastern descent from the Kambacho pass was found to be free from snow, a little from the summit, at the beginning of December, and on the northern approach the snow was supposed to become perpetual at about 15,000 feet, though the fresh falls of the previous October forced Dr. Hooker to be in some measure guided by the people of the country in this estimate. On the ascent to the Wallanchun or Wallungsum pass the snow-line was again estimated, though under similar circumstances of doubt, to be at 15,000 feet. The Pandit crossing the same pass, his Tiptala, on the 16th August, 1871, found it covered with snow,² and Dr. Hooker on December³ 31st, 1848, crossed "with snow on both sides up to the shoulder."

The following are the results of trigonometrical measurements of the elevation of the inferior edge of the snow observed on spurs of the Trisul and Nanda Devi groups of peaks, made by General R. Strachey before the winter snow had commenced in 1841 :—

Point observed.	<i>Height of snow-line.</i>		Mean.	Remarks.
	On south exposure.	On west exposure.		
No 1 ...	16,706	15,892	16,298	These heights were calculated from observations made with the theodolite at Almora and Binsar. The distance of these two places, which served as a base, was obtained by measurement from a map of points fixed by the G. T. Survey. The elevations of the two places were taken from Captain Webb's trigonometrical survey.
2 ...	17,007	
3 ...	17,206	14,898	16,051	
4 ...	15,347	
Mean ...	15,566	15,395	{ 16,174 15,980	

¹ Himalayan Journals.
derived from the Wallungsumgoia to the south: the pass itself has an elevation of 15,616 feet.

² Reports, 1871, p 1: the name appears to be derived from the Wallungsumgoia to the south: the pass itself has an elevation of 15,616 feet.
³ J., A. S., Ben., XVIII., i, 524.

The points 1, 2 and 3 are on ridges that run prominently out in a south-westerly direction, from the great peaks of Trisul G. (G. T. S.) same as Webb's No. XI. and H. (G. T. S.), same as Webb's No. XII. The dip of the strata being to the north-east, the faces exposed to view from the south are for the most part very abrupt and snow never accumulates on them to any great extent. This will explain the difference between the heights at which snow was observed on the southern and western exposures, the ground having been less steep on the latter and better able to retain the snow; but in these places it was in very small quantities, and had probably fallen lately, so that its height may probably indicate the elevation below which the light autumnal falls were incapable of lying rather than the inferior edge of the perpetual snow. It is further to be understood that below this level of 15,000 feet the mountains were absolutely free from snow, excepting those isolated patches that are to be seen in ravines or at the head of glaciers, which do not affect such calculations as these. The point No. 4 was selected as being in a much more retired position than the others, and is situated not far from the head of the Pindar river, between the peaks of Nanda Devi, I. (G. T. S.), same as No. XIV. of Webb, and of Trisul H. (G. T. S.) It was quite free from snow at 15,300 feet. On the whole, therefore, General Strachey is inclined to consider that 16,000 feet may be given as a close approximation to the maximum height to which the snow recedes every year on the most southern and external ranges in Kumaon.

This result appears to accord well with what has been observed by Dr. Gerard in a visit made by him to the Shatúl pass in the Bisahr range expressly for the purpose of determining the height of the snow-line.¹ He reached the pass, the elevation of which is 15,000 feet, on the 9th of August, 1822, and remained there till the 15th of that month. The southern slope of the range was generally free from snow, and he says that it is sometimes left without any whatever. On the top of the pass there was no snow, but on the northern slope of the mountain it lay as far down as 14,000 feet. On his arrival rain was falling, and out of the four days he was there, it rained and snowed for the greater part of three. The fresh snow

¹ Tours I., 289-347.

that fell did not lie below 16,000 feet, and some of the more precipitous rocks remained clear even up to 17,000 feet. Dr. Gerard concludes that 15,000 feet is about the height of the snow-line on this range, but it will be seen that Dr. Gerard was there rather early in the year, and General Strachey, from what he heard from others who crossed the range later in the season, thinks that 15,500 feet will probably be a better estimate, even if it should not be carried still higher. At the beginning of the month of July, Captain Gerard found heavy snow on the northern face of this Bisahr range at about 15,000 feet, and the Kunlia pass, the elevation of which he states to be 17,000 feet appears never to be free from snow¹ Dr. Thomson² agrees that the estimate formerly made by General Strachey of 15,500 feet, which his subsequent researches led him to believe was a little too low for Kumaon, is as nearly as possible correct for the Bisahr range. He adds—“Captain Herbert, in his geological report, had fixed upon 15,000 feet, which is a little too low even in the district of Bisahr, to which his estimate, I believe, refers. In the trans-Satlaj Himálaya, from the diminished amount of summer cloudy weather, the snow-level is probably a little higher.” The Chanba range and the Pir Panjal, south of Kashnir, both of which rise immediately from the low external sandstone hills, just enter, he tells us, the region of perpetual snow. The highest peaks of the former are about 16,700 feet, and its mean height about 15,000 feet above the sea, and its snow-line will consequently be not far from 16,000 feet. Major A. Cunningham also places the snow-line on the most southern ranges of the Himálaya to the west of the Ganges at about 16,000 feet.³

When, however, we advance into the interior of the chain, across the snowy range, after having once passed over any range of sufficient height to come within the limits of perpetual snow, we invariably find that there is less snow on all such ridges of similar altitudes so that when we arrive at the Indian watershed, the snow-line has risen to about 18,500 feet, and on the summit of the tableland it reaches to an elevation of 20,000 feet. Dr. Hooker observed this phenomenon in Sikkim, and bears testimony to the gradual rise of the snow line as we enter among

¹ *Ibid.*, II., 71.² *Travels*, p. 487.³ *Ladák*, pp. 73-77.

the peaks covered with perpetual snow. Near Zemu, twenty miles north-east from Kanchanjunga, he found little snow on south exposures at the beginning of July. A little further in, above Phalung, in the middle of the same month, the snow-line was supposed to be about 16,500 feet, and at the end of the month many plants were obtained at 17,000 feet. Another ascent in the same vicinity about the same time did not carry our traveller to perpetual snow at 16,800 feet. On the flanks of the Kanchanjha broad summits were seen quite bare of snow at 18,000 feet. Dr. Campbell¹ who accompanied Dr. Hooker on his return journey in September, notes that vegetation ceased at the foot of the Dankia pass at 18,000 feet, and there was no trace of it within 500 feet of the summit on either side. There was no snow on the road as he ascended the north face nor as he descended the south face, but it lay in patches amongst the rocks all the way on both sides. On the mountain to the west of the pass snow lay deep in hollow places, but these may have had glacial ice in them though the surface of the snow was then smooth. The line of snow would here be 19,800 feet, and further north in Tibet it rose to 20,000 feet. Bhurutso, on the 18th October, had not a particle of snow on it at 18 or 18,500 feet, whilst in the Lachung valley in Sikkim to the south snow was lying at about 15,000 feet. Dr. Campbell adds:—"South of the Himálaya, the quantity of snow that falls is very much greater than in Tibet, and from the greater moisture of the air and cloudiness of the sky, it is not carried off with the rapidity of evaporation which obtains in Tibet, where you do not find even a rill of water from the melting snow. Besides in Tibet the snow falls in light, feathery skiffs, and not in flakes. I believe that the lowest snow-line we saw on the mountains to the north of us in Tibet must have been upwards of 22,000 feet. On the Kambajang range, which, comparing them with Bhurutso, must be 20,000 feet at least, there was not a particle of snow."

We have the results of General Strachey's experience for
 Kumaon. Garhwál. Towards the end
 of August, 1848, he crossed the Barjikang
 pass leading from Rálam to Juhár on a subordinate ridge between
 the Nandakot and Panchachuli peaks. Although this pass has an

¹ J., A. S. Res., XXV., 565.

elevation of about 15,400 feet, not a vestige of snow was met with on the ascent from the south-east, and only a very small patch remained on the north-western face, and indeed, in no considerable quantity, up to 17,000 feet. The vegetation on the very summit of the pass was far from scanty, though it had already begun to break up into tufts, and had lost the character of continuity it had maintained to within 500 or 600 feet of the top; but many species of flowering plants, all evidently flourishing in a congenial climate, showed that the limits of vegetation and regions of perpetual snow were still far distant. This place is within ten to fifteen miles of the most southern border of perpetual snow. The Unta-dhúra pass has an elevation of about 17,300 feet, and lies to the north of the great peaks nearly at the crest of the watershed. There was no snow along the southern ascent to this pass, at the top of which General Strachey arrived in September, 1848, in a little drizzle of rain that at last turned into snow. The ground was quite free from snow, being worked up into a deep black mud by the feet of the cattle that had crossed it. There was, however, on the north side of the pass an accumulation of snow some little way down, extending perhaps 200 feet, apparently the effect of the drift through the gap in which the pass lies. No snow was seen on the hills on either side within some few hundred feet, and the snow-line was certainly above 18,000 feet. Vegetation reached to within 300 or 400 feet of the summit.

The Chor-hoti pass (18,000 feet) and the Marshak pass, (18,400 feet), both to the north of Níti, have a position
 Garhwál. relative to the great snowy masses nearly similar to Unta-dhúra. The Marshak pass was crossed in July by General Strachey, a time rather too early to judge fairly of the snow-line, which is also obscured by the presence of a glacier that fills up the valley by which the pass is approached.¹ On the Chor-hoti pass in September there was not a vestige of snow on any part of the southern face of the ridge that the route crosses, but on the north face was a patch that was plainly perpetual, descending some hundred feet to a glacier which was connected with that just mentioned to have been crossed at Marshak or Balchak. The snow-line was, therefore, here no doubt near 18,500 feet. The Kyungar and

This expedition is apparently referred to in J., A. S., Ben., XIX., 79.

Balchha passes, each about 17,500 feet in elevation, and both lying to the north of the Unta-dhúra pass, were equally free from snow on their southern faces in September, small quantities only being found on the northern aspects. The highest points on the ridge, over which the latter of these passes leads, only just exceed 18,000 feet in elevation, and in fact it does not come within the limits of perpetual snow, nor does it appear snowy when viewed from the Tibetan plain to the north of it. The vegetation on all these passes reaches to about 17,500 feet. Lieutenant Weller visited the Balchha pass¹ on the 1st June, 1842, and found "towards the top of the ascent a tolerable quantity of snow, but in detached portions." The Lákhar pass also, to the north of Unta-dhúra, was crossed by General Strachey in September. It has an elevation of about 18,200 feet, and was found free from snow on both sides, as well as the Jainti ridge some 200 feet higher. This latter is, however, a somewhat detached spur, and the snow-line was manifestly near, for unbroken snow could be seen in more sheltered places considerably below this elevation. General Strachey thinks that, on the whole, 18,500 feet may be considered a fair average height of the snow-line in this locality. Lieutenant Weller crossed the Unta-dhúra pass at the end of May, and found more soil than snow visible, whilst snow was scattered thinly on either side, but the northern slope presented one unbroken sheet of steep snow.² In September (28th) Captain Manson found the last ascent to this pass quite free from snow. A detached peak, Lanjar, a little to the north of the Niti pass, and having an elevation of 18,400 feet, was found by General Strachey nearly quite free of snow having only a patch lying in a ravine on the north side of the hill. Two other peaks near the Balchha pass, seen from Lanjar, having an altitude of 18,100 and 18,200 feet respectively, were also quite free from snow, so far as could be ascertained at the distance. Mr. J. H. Batten, who visited the Niti Pass in 1837, found it free from snow, of which the first heavy fall did not occur till the 11th of October.³

In the more western part of the mountains the authority of the Gerards, of Dr. Thomson and of Major A. Cunningham coincide in fixing the snow-line at much about the same level as that just assigned to Kumaon. Captain J. D. Cunningham also accepts

¹ J., A. S., Ben., XII., 97.

² *Ibid.*, XII., 87.

³ *Ibid.*, VII., 816.

the results of General Strachey's observations. In upper Kunaor Captain Gerard¹ found a little snow on either side of the Kynbrang pass (18,300 feet) in July, but none on its summit, and the summits in the neighbourhood, though attaining a height of 18,000 or 19,000 feet, were only just tipped with snow.¹ The Gangtang pass, also 18,300 feet and lying a little farther to the west, was snowy for the last few hundred feet at the end of the same month. The Kynbrang pass is on the Indian waterparting, and the Gangtang pass a little within it, but both the observations were made before the snow-line had attained its maximum elevation. The gradual descent of the snow-line as we advance southwards is shown by the fact of the Charang pass having an altitude of only 17,400 feet, and lying between the passes above named and the Bisahr range, being said to be never free from snow, though early in July it had already melted up to 16,300 feet. North of the Satlaj, under the peak Leo Porgyul, the surface was found to be free from snow in October up to 19,000 feet or even higher, while west of that river on the Manirang or Rupak pass, having an elevation of 18,600 feet, the summit was covered with newly fallen snow at the end of August, showing that the level of perpetual snow was nearly at its maximum. Snow was, however, met with on the road to the pass, but this was due to avalanches and drifts and to the fact of the road lying in a deep glen. Dr. Thomson, who visited the Kárákorum pass in August, 1848, estimated the snow-line on the journey back to Sáser at 17,500 to 18,000 feet but to the northward and eastward it was much higher, probably not less than 20,000 feet.² Trotter³ also notes that the Kárákorum pass (18,550 feet) is always free from snow in summer, whilst the Sáser further south is seldom, if ever, free from snow.

Regarding the height of perpetual snow on the tableland of western Tibet Captain H. Strachey is still the best authority. He writes:—"from a series of minute observations on the snow-level, made during two years, in the course of which I crossed twenty-five passes elevated from 15,000 to 19,000 feet at various seasons between the end of

¹ *Tours in the Himalaya*, II., 73, 117, 133, 242: *Account of Koonawur*, 157-9.
² *Travels*, p. 437. ³ *Report*, p. 11.

April and the beginning of November, I have arrived at the following conclusions. The snow-line in the central and northern parts of west Nári attains an extreme height of nearly 20,000 feet. It lowers on approaching the Indian Himálaya, and on the southernmost parts of the Indian watershed descends perhaps as low as 18,000 feet." The mountains under 20,000 feet in height in the northern and more open parts of the tableland will, he adds, be almost entirely denuded of snow during the latter part of the summer. General Strachey, during his visit to the tableland north of Garhwál during September, 1848, found snow only in patches in sheltered ravines, but the highest summits in the district through which he passed were only 18,400 feet. Perpetual snow was not found on any of the hills between the Indian water-parting and the Satlaj. The height of the snow-line on the south face of the peak of Kailás was observed in the month of September by means of a theodolite, and found to have an elevation of nearly 20,500 feet, and the altitude of a peak on the verge between the Satlaj and the Indus which was only tipped with snow in August was in like manner determined to be 20,500 feet above the sea: so that, making a fair allowance for the difference between the northern and southern exposures, the mean snow-line was in both cases about the same. The limit of snow on the Pámír is reported to be between 16,000 and 17,000 feet and on the Alái Pámír about 14,000 feet.

Throughout Kumaon and Garhwál there are several lakes, but the chief in size and beauty occur in par-
 Lakes. gana Chhakháta, the Westmoreland of Kumaon. These are known as Naini, Bhím, Sát, Naukuchiya, Malwa, Khurpa, Sukha, Sariya, Khuriya, &c., with the affix 'tál' or 'lake' attached. The following table gives some information regarding the principal lakes:—

Name.	Height above sea-level.	Greatest length.	Greatest breadth.	Greatest depth.	Approximate area.
	Feet.	Feet.	Feet.	Feet.	Square feet.
Naini Tál ...	6,407	4,733	1,518	93	5,149,000
Bhím Tál ...	4,500 ¹	5,580	1,490	87	4,900,000
Naukuchiya Tál ...	4,000 ¹	3,120	2,270	132	4,849,500
Malwa Tál ...	3,400 ¹	4,480	1,833	127	4,909,600

¹ Approximate measurements.

It has been suggested that these lakes were formed by glacial action, but Mr. Ball in a recent paper¹ combats this view, and assigns their existence to landslips which closed up the valleys in which they occur. Be this as it may, they form one of the most remarkable and beautiful features of the Lower Himálaya in Kumaon. The lake of Naini lies in a valley which runs about north-west and south-east, and is surrounded on all sides except the east by lofty ridges, Sher-ka-danda, Chína (8,568 feet), Deopátha (7,989 feet), and Ayárpátha (7,721 feet). Bhím Tál lies in a comparatively open valley with a hill to the south of the lake rising some 1,800 feet above its level. Further east in the same valley is Naukúchiya Tál, occupying a hollow in the slope, and without any remarkable hills around it. The Sít Tál or seven lakes, lie within a circle of hills between Bhím Tál and the valley of the Naini Tál river, and Malwa Tál lies to the north of Bhím Tál in a deep valley, the sides of which rise up abruptly from the level of the lake. The only lakes of importance in Garhwál are the Gudiyar Tál in patti Dasauli Malli and Diuri Tál in patti Káliphat Malla, neither of which can compare with the Kumaon lakes in size or beauty.

¹ *Rec., Geo. Sur.*, XI, 2, p. 174.

CHAPTER V.

METEOROLOGY.¹

CONTENTS.

Preliminary sketch of climate. Contrast between the eastern and western parts of the plain and the Himálaya. Radiation, solar and nocturnal. Temperature :—Diurnal and annual ranges,—Vertical decrement,—Height of snow-line. Pressure and winds :—Barometric tides,—Mountain winds,—Annual variation of pressure,—Monsoons. Humidity :—Vertical distribution of vapour,—Relative humidity,—Cloud. Rainfall :—Distribution on plains and on the Himálayan slope,—Annual variation,—The winter rains,—The monsoon rains.

THE climatological conditions of these hill districts are a most important element in their physical geography, and will therefore require to be treated at considerable length. An exhaustive discussion of the meteorology cannot yet be attempted, but sufficient data have already been collected to serve as a basis for a general description of the climate, and at the same time to throw some light on several of the more interesting problems of meteorology. In this latter respect the Himálaya, on account of its less distance from the equator and its greater elevation, presents many points of advantage as compared with the Alps and other European mountain systems ; and already some important general conclusions regarding the physics of the atmosphere have been drawn from the observations that have been made in it. The mere statement of the fact that nearly all the snowy peaks and most of the passes over the Indian watershed stand above the lower half of the atmosphere, and thus completely cut off all communication between India and Central Asia, except in the upper strata, indicates how much regarding the general movements of the atmosphere may be learnt from observations taken in India and the Himálaya.

Of late years, a good deal has been done in the way of collecting trustworthy meteorological data for the mountain zone by the establishment of Government observatories at certain points. The places where observations are made at the public expense must

¹ Written by Mr. S. A. Hill, Meteorological Reporter to the Government of the North-Western Provinces and Oudh, for this volume.

always, however, be few, and it is desirable that more should be done in the way of enlisting the services of volunteer observers. Temperature and rainfall observations are now made at many tea-gardens in Kumaon ; but, as a rule, so little attention is paid to the hours of reading, the exposure of the instruments, and the continuity of the registers, that the results are of no value for scientific discussion and comparison. By far the most important of the observations taken in the north-west Himalaya, prior to the establishment of regular observatories, were those collected by Lieutenant (now General) R. Strachey, of the Bengal Engineers, in 1848 and 1849. Some of General Strachey's deductions from them have been given to the world in the *Proceedings of the Royal Society* and the *Journal of the Asiatic Society of Bengal* ; but others have not yet been published, though they were long ago embodied in a work on the "Physical Geography of the Himalaya" that has been placed at the disposal of the compiler of this volume. Considering the scanty nature of the materials General Strachey had to work with, the conclusions arrived at were wonderfully accurate ; and though some of them were opposed to the generally received opinions of the European meteorologists of the day, they have been confirmed in almost every respect by the more extensive data subsequently obtained from the Himalayan observatories.

In the following pages a somewhat detailed discussion of all the data available for meteorological inquiry will be given after a brief general sketch of the climate. The several elements of meteorological observation will be taken in the natural order of cause and effect, commencing with solar radiation and afterwards passing on to temperature, barometric pressure and winds, and the distribution of vapour and rain.

The order of the three seasons on the plains of Upper India—the cold, the hot, and the rainy—is now well known even in Europe. After the close of the rains at the end of September or beginning of October the sky is serene and the atmosphere transparent. Owing to the absence of cloud and the rapidly diminishing proportion of water vapour, the air is also very diathermanous ; that is, it permits the free passage of heat from the sun to the earth in the daytime, and in the calm nights that prevail at this season the radiation of

Sketch of climate.

heat into space goes on so rapidly that the earth's surface and the air resting on it become very cold before morning. The months of October and November are thus characterized, not only by clear skies and calms, but by a great temperature range and heavy dews at night. These conditions prevail through the greater part of December, and towards the end of that month and in the beginning of January, the exposed thermometer sometimes falls 10 degrees below freezing at places as far down the plain as Allahabad and Benares. In the Panjáb it is much colder, and there the shaded thermometer sometimes reaches the freezing point.

About the end of December and in January and February, however, clouds often interfere with the free radiation of heat at night, and the daily range of temperature for these months is less, on the average, than that of November. Some rain usually falls at this time of the year, especially in the Panjáb and the higher districts of the North-Western Provinces. In March and April the temperature rises rapidly, especially at a distance from the mountains, and the air becomes extremely dry. Hot winds from the west or north-west blow down the valley of the Ganges and rapidly change the appearance of the whole country from that of a highly cultivated plain to a parched and sandy desert, almost the only green things left being the groves of mango trees. In April, the daily range of temperature over the plains is at a maximum, exceeding 30 degrees in most parts of the North-Western Provinces and the Panjáb. The nights are still tolerably cool, though in the daytime the thermometer ranges as high as 110° F. or even higher sometimes.

During May and the first half of June the temperature continues to increase, though much less rapidly than in March and April, until by the 15th or 20th of June, if the periodical rains have not commenced, the temperature is probably higher in North-Western India than anywhere else in the world. In the North-Western Provinces the shaded thermometer has only been known to rise once or twice above 120° F., but in the Panjáb temperatures as high as 123° or 124° have been recorded. The days in June are thus only a few degrees hotter than those of April; but, as the rainy season approaches, the range of temperature diminishes and the nights become insufferably hot and close.

Rain seldom falls in the hot weather, the falls that do occur generally taking place during thunderstorms. About the middle of May, however, the quantity of water vapour in the air begins to increase rapidly, betokening the approach of the rainy season. This vapour is probably brought by the prevailing south-west upper current of the atmosphere which seems to descend gradually until it merges with the surface sea winds of the Bay of Bengal and forms "the south-west monsoon" or prevailing wind of the rainy season. In Northern India the lowest strata of the sea winds are deflected from their normal course by the mountains and directed towards the seat of highest temperature in the Panjáb, thus appearing as east or south-east and not as south-west winds. Along the foot of the hills these easterly winds are felt occasionally by the middle of May, when the quantity of vapour in the air first begins to show signs of a rapid increase.

During the latter half of June the sea winds increase in strength and gradually advance along the foot of the Himálaya, until, by the beginning of July, the rains have usually set in all over Northern India. In ordinary years rain continues to fall, not steadily but with frequent intermissions or "breaks," until about the end of September, when the easterly winds cease, except close to the hills, where they last a month longer, and are succeeded by calms or feeble currents from the west. In the Panjáb the rains begin later, are lighter and more intermittent, and end sooner than in the North-Western Provinces, and the length and intensity of the rainy season increase regularly as we approach the sea in Bengal. During the rains the temperature averages about 85° over the greater part of Northern India. The daily range at this time varies from 8 to 12 degrees, being greatest in the driest districts.

The extremes of heat and cold are much greater in the Panjáb and the upper part of the North-Western Provinces than in Bengal, for two reasons—the greater distance from the sea and the higher latitude. On account of its proximity to the sea and its heavy rainfall, Bengal is moist and cloudy at all seasons compared to the Panjáb. This condition of the atmosphere, by retarding the radiation of heat, renders the climate of Bengal more equable than that of the Panjáb, just as an insular climate is more equable than a continental one. Again, the latitude of

the Panjáb, which is 7 or 8 degrees higher than that of Bengal, causes its winters to be much colder and its summers much hotter. At first sight it seems anomalous that a place should be the hotter the more distant it is from the equator, at any season of the year; but when it is borne in mind that the quantity of heat received from the sun is directly dependent upon the length of the day as well as on the elevation of the sun above the horizon, the anomaly disappears. Various mathematical physicists from Halley downwards, including Poisson¹ and more recently Meech,² have calculated the total heating effect of the sun in different latitudes during a day or other given period of time. The latest investigation of this kind has been made by Wiener³ of Carlsruhe, who finds that while the maximum of heat for the whole year falls on the equator, the maximum for the 21st of June is at the north pole, where the sun remains above the horizon for twenty-four hours, and has an altitude of nearly $23\frac{1}{2}$ degrees for the whole of that time. In the summer half year, from equinox to equinox, most heat falls on a zone about $25\frac{1}{2}^{\circ}$ north of the equator, and during the three months nearest to the summer solstice—that is, from the 7th of May to the 6th or 7th of August, the zone of greatest heat lies about 41° N. The total heat received during these three months by an area in latitude 40° or 41° N. is more than a fifth greater than that which falls on an equal area at the equator. The actual increase of temperature produced is much more than this, for the mean temperature is determined by the balance between the gain of heat during the day and the loss at night. When the gain of heat from the sun at any place is greater than at the equator, on account of the length of the day, the loss at night must be correspondingly less.

This excess of solar heat in summer, together with the dryness of the air and the absence of cloud, seems to account for the excessively high temperature of June and July in the extreme north of the Panjáb and on the plains of Yárkand and Káshgar still farther north. In the moister zone of the mountains, the direct action of the sun is less observable; but beyond the Indian watershed it is by far the most important factor in determining the character of the seasons.

¹ *Théorie de la Chaleur*, 1835 edition, page 478.

Contributions to Knowledge, Vol. IX.

² *Zeitschrift der Oesterreichischen Gesellschaft für Meteorologie*, Band XIV, page 133.

³ *Smithsonian Contributions to Knowledge*, Vol. IX.

Regarding the succession of the seasons in the mountain zone, General Strachey says :—

“The same general sequence of the seasons takes place in the mountains as in the plains. Here, however, every altitude has its own special temperature, from the lower valleys where the heat is still overpoweringly great, to the regions of eternal frost ; but at all elevations in summer the force of the sun’s rays is excessive. The summer rains, too, gradually diminish in strength as we move along the chain from east to west, being at their maximum in Sik-kim, but still being felt slightly on the ranges north of Pesháwar. The heaviest falls invariably take place on those portions of the chain most exposed to the south ; increasing in amount up to a certain height [not very exactly determined, but probably about 4,000 feet]; at the same time every high and continuous ridge most sensibly diminishes the supply of rain that falls on the country to the north of it, and we find, as we approach the Indian watershed, that the quantity is very small, and that the monsoon only just drops a few partial showers on the southern border of Tibet. The winter, as may be supposed, is extremely rigorous on the summit of the table-land ; and at this season, or in spring, the only important precipitations of moisture take place in the form of snow, but they are exceedingly small in quantity.”

The reason why every altitude has its own special temperature is that the air is warmed chiefly by contact with the hot ground on which it rests, and but little by direct absorption of the solar rays. The air in contact with the ground, expanding and becoming less dense, rises up, but in doing so its heat is rapidly converted into the work of expansion ; the result being that the temperature of the upper strata can never rise so high, on the average, as that of the air near the ground. Dry air, if heated only at the bottom, would lose 1 degree Fahrenheit for every 183 or 184 feet of ascent. In moist air that is precipitating rain, and thus being warmed by the latent heat of the condensed vapour, the rate of decrease is much less. The rate actually observed, both in balloon ascents and on mountain sides, is less than that calculated theoretically ; because even dry air is to some extent warmed directly by the sun’s rays, while air saturated with moisture has a very considerable absorbing

power. On mountain slopes also the temperature falls less rapidly than in the free air over the plains—at all events for the first nine or ten thousand feet of ascent, the reason being that the air is heated by contact with the mountain sides.

No data at present exist from which the average intensity of solar radiation, and its variations from time to time, can be estimated with any approach to exactness. Any deductions made by passing from radiation to other meteorological phenomena must therefore to a great extent be based on theoretical considerations.

The instrument hitherto used to measure the intensity of the sun's heat has been a maximum thermometer with a blackened bulb surrounded by a thin glass case more or less completely exhausted of air. If the exhaustion were perfect, the temperature of the instrument would be determined by radiation to and from surrounding objects; including under these the glass case of the instrument which is in contact with the air, the sun, the ground, the clouds, and the open sky. Were solar thermometers all made exactly alike and exposed under absolutely identical conditions, the excess temperature of the instrument above the contemporaneous temperature of the air would be a measure of the excess of radiant heat falling on it from objects above the horizon over that which passes away from it in an upward direction. The following table gives the average value of this difference for each month at six stations. Corrections have been applied as far as possible for differences of instrument and exposure, except at Dehra, for which the corrections are not known :—

I.—Monthly mean excess temperature of the solar thermometer above the maximum in shade.

Stations.	Elevation in feet.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
Leh ...	11,538	49.9	53.1	62.3	63.2	58.7	58.1	54.1	62.8	63.8	68.9	54.1	50.9	58.7
Chakrata,	7,652	71.1	67.1	69.4	69.5	69.1	66.3	63.0	61.5	70.3	69.8	69.4	68.5	67.9
Ranikhet,	6,069	61.1	60.6	62.6	60.4	60.3	57.1	50.8	52.1	59.8	57.3	52.7	55.7	57.5
Dehra ...	2,232	51.0	52.3	57.6	57.1	56.8	54.3	53.8	53.1	57.7	58.5	55.6	51.9	54.9
Roorkee...	887	52.2	51.5	55.9	54.6	53.7	51.3	51.9	51.8	54.5	54.5	55.1	52.5	53.3
Bareilly ...	868	50.0	52.6	54.1	54.4	55.3	54.2	50.7	50.1	49.3	52.0	52.4	46.9	52.0

If the air were absolutely diathermanous, the altitude of the sun above the horizon and the vertical thickness of the atmosphere above the place of observation should have no effect upon the temperature differences in the table, which should therefore be the same for all the stations and for every month of the year. But the air having some absorbing power, the differences should be greatest when there is least air for the sun's rays to pass through ; that is, at the highest stations and in the summer months. Up to Chakráta the excess temperature of the solar thermometer does increase with a fair degree of regularity ; but it appears to be less at Leh than at Chakráta, contrary to all theory. There is also no regular increase apparent in the heating power of the sun as the season changes from winter to summer. The truth is that the indications of the black-bulb thermometer are affected by so many disturbing causes, that after all possible corrections they are of little or no value for inter-comparison ; though with the same thermometer, at the same place, and under absolutely constant conditions of exposure, the figures for one year may be to some extent comparable with those for another.

The results of observations with the nocturnal radiation thermometer are even more unsatisfactory, owing to differences in the height of the instrument above the ground and in the nature of the ground surface itself, whether grassy or bare.

II.—Monthly mean depression of the grass thermometer below the minimum in shade.

Stations.	January.	February.	March.	April.	May.	June.	July.	August.	Septem-ber.	October.	Novem-ber.	Decem-ber.	Year.
Chakráta,	9.5	6.5	7.2	7.5	6.3	6.0	8.1	4.3	5.2	8.8	9.2	11.0	7.2
Ránikhet,	14.5	13.6	13.9	13.3	12.0	9.0	5.3	6.0	8.9	13.5	17.2	19.1	12.2
Dehra ...	4.4	4.4	4.3	4.7	4.8	3.6	3.3	1.5	3.3	3.5	4.8	4.6	3.8
Roorkee...	6.3	5.8	6.4	6.3	5.0	3.5	2.4	2.9	3.9	6.3	7.3	6.3	5.2
Bareilly...	7.0	7.5	8.0	10.0	10.5	8.8	4.3	3.3	3.4	6.4	5.2	8.0	7.1

The figures in table II. serve to show that the depression of the nocturnal radiation thermometer below the minimum in shade is less in the rainy season than in the dry, and that both at the hill stations and on the open plain the refrigeration of the earth's surface during

the night is probably greater than at Dehra, where the observatory is situated in a well-wooded park. They do not throw any light on the question whether the ground surface cools more rapidly at night on mountain tops or on the plains; though it is probable that in the clear, calm nights of the cold weather the difference, if any, is in favour of the plains; since there the air cooled by contact with the ground remains in contact with it, whereas on the mountains the cooled air constantly drains away, and is replaced by warmer air from the surrounding free atmosphere.

It has been already stated that, in the western Himálaya, every elevation has its characteristic mean annual temperature. Each elevation has probably also a distinctive form of variation of temperature during the year, and the daily variation is different at different altitudes, in range if not in general form.

For a proper discussion of the distribution of temperature in a hilly country a very large number of observations would in most cases be required; and these should be made at places chosen so as to be at nearly equal distances from each other vertically, and at the same time fairly distributed in latitude and longitude. On the southern slope of the Himálaya it fortunately happens that differences of latitude and longitude make but little difference in the mean annual temperature. The sea-level values of the mean temperature at the Sub-Himálayan stations from Lower Asám to Ambála all lie between 76 and 78 degrees Fahrenheit, and the temperatures of places at about 7,000 feet elevation along the whole range from Darjiling to Marri in the north of the Panjáb do not differ more than 2 or 3 degrees.

Both along the plains and at the level of the hill sanatoria the highest mean temperatures are found to characterize the regions lying between the Káli and Satlaj rivers. The chief reason for the great uniformity of mean temperature at the same elevation that prevails over the whole Himálayan region—that is to say, through more than 7 degrees of latitude and 17 of longitude—is the greatly increased heat of summer in the north-west as compared to the east. In Bengal and Sikkim the sun's rays when most intense are to a great extent cut off by cloud, whereas

in the Panjáb and the north-west Himalaya the winter is almost if not quite as cloudy as the summer. For these reasons Darjiling has very nearly the same temperature in January as Simla, Chakráta, or Mussooree, while in May and June it is seven or eight degrees cooler. The comparatively low temperature of the summer at Darjiling renders the mean for the year nearly two degrees lower than that of Marri in the extreme north-west, though the effect of latitude is apparent in the low temperature of Marri in January.

On account of this uniformity of temperature a small number of observations, at places chosen specially with reference to height above the sea, will enable us to ascertain the most important features of temperature distribution in the Himalayan districts of the North-Western Provinces. The following table gives the mean monthly temperatures of twenty-one places, including the two stations on the plains that were given in the previous tables. All the other places except Dharmśála lie in one or other of the three districts of Kumaon, Garhwál, and Dehra Dún, or in parts of Kunáwar, Lahul, and Ladák to the north of Dehra Dún. The monthly means from a Government observatory at Dharmśála in the Panjáb have been inserted, though this station lies nearly two degrees west of Dehra Dún, because it was considered desirable to have some trustworthy figures for places about 4,000 feet above the sea ; and the only other station near that altitude is Háwalbágh in Kumaon, for which we have but one year's observations.

III.—Mean monthly and annual temperatures of places in Kumaon, Garhwal, Dehra Dun, and adjacent districts.

Place.	Latitude N.	Longitude E.	Elevation in feet.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.	Number of years.
RANIKHET	29° 11'	79° 27'	568	57.5	63.0	73.0	83.5	89.1	90.5	84.3	84.0	89.0	76.4	66.5	58.6	75.8	11
ROOKER	29 53	77 56	887	58.7	61.1	70.5	81.5	87.9	90.8	85.2	84.1	89.7	75.5	65.0	57.8	74.9	17
Nainagar	30 13	78 50	1,950	71.0	80.0	84.0	79.0	84.0	79.0	1
Kānī	30 30	77 50	2,000?	58.3	61.0	63.6	77.7	81.2	86.0	83.7	80.2	77.2	70.8	63.2	59.7	71.8	1-3
DEHRA	30 30	78 8	2,233	54.8	58.0	65.8	76.1	82.1	84.9	80.2	79.0	77.3	70.4	63.5	56.2	70.6	12
Hawalbagh	28 38	79 37	4,114	47.0	53.0	61.0	66.0	73.0	76.0	78.0	79.0	75.0	69.0	60.0	53.0	65.8	1
DHARMADIA	33 20	76 33	4,495	44.6	49.4	57.3	64.3	72.3	75.8	70.7	69.5	69.0	64.6	57.7	50.5	62.8	4-5
Panti	30 10	78 53	3,310	60.0	69.0	73.0	73.0	69.0	67.0	61.0	1
Almora	29 53	79 38	3,346	46.3	52.3	57.4	64.7	70.3	73.0	73.2	73.5	73.5	65.4	57.9	51.2	63.2	6-7
Lahughat	29 34	80 4	3,449	44.3	43.8	52.3	60.9	66.0	71.0	71.1	70.7	68.7	63.0	51.9	46.4	59.4	3-5
Musoorie (1)	30 24	78 10	5,850	43.4	47.0	57.2	65.3	67.0	70.8	64.4	63.8	63.2	63.0	56.6	47.6	59.6	1-3
RANIKHET	29 53	79 29	6,059	46.1	49.2	56.3	65.2	67.8	71.8	67.9	67.2	66.4	61.0	55.7	50.2	60.4	9
Kānī Tal	29 50	79 31	6,463	43.8	43.3	53.3	60.1	66.0	69.1	68.4	67.3	65.4	59.8	52.7	46.4	58.2	10
Musoorie (2)	30 28	76 7	6,937	41.5	43.4	53.0	59.6	63.5	68.0	64.7	64.1	63.0	57.5	52.2	46.4	58.7	2-14
CHAKOTA	30 40	77 33	7,032	41.6	43.7	50.4	59.6	64.5	68.0	64.5	64.1	63.0	57.5	52.2	46.4	58.7	11
Laudhar	30 27	78 11	7,311	37.6	44.3	49.4	56.6	61.8	68.0	63.8	63.9	64.3	56.4	49.4	44.1	55.2	4-8
Kānī (Kandwar)	32 0	79 30	9,396	28.8	34.6	37.2	46.8	56.7	63.7	58.6	63.0	61.8	54.1	43.6	36.1	49.4	3
Kānī (Lahol)	33 34	77 1	10,343	24.0	36.0	44.0	47.0	49.0	64.0	63.0	60.0	53.0	46.0	37.0	37.0	44.9	1
Niti	30 48	79 24	11,464	58.0	58.0	55.0	1
Lam (Ladakh)	34 10	77 43	11,388	17.6	27.1	30.3	40.1	46.6	53.6	60.5	57.5	49.3	37.9	30.7	24.4	36.3	1
Spiti Valley	33 10	78 0	12,000	17.5	18.7	22.3	37.3	45.9	56.9	61.7	58.3	55.7	38.3	33.6	17.2	37.2	2-7

Authorities for the above table.

- BARKILL :** Registers of the Government Meteorological Observatory, 1869 to 1879
- BOORKE :** Ditto, ditto, 1863 „ 1879
- DHARMILA :** Ditto, ditto, 1871 „ 1875
- BANIKHET :** Ditto, ditto, 1871 „ 1879
- CHAKRATA :** Ditto, ditto, 1869 „ 1879
- LEH :** Ditto, ditto, 1871 „ 1877
- DEHRA :** Registers of the Observatory at the G. T. Survey Office, 1868 „ 1879
- NAINI TAL :** Registers of Government Observatory, 1863 to 1869 ; and observations by Dr. Payne in 1851-54, given at page 496 of the third volume of the *Results of a Scientific Mission to India and High Asia*, by the brothers Schlagintweit.
- MUSBOORKE (1) :** Records of an observatory at Saint Fidelis's School for 1877-80 ; somewhat fragmentary.
- (2) :** November to April, Sir A. Waugh's and Mr. Mackinnon's observations in 1853-54, also one year's observations from Dove's tables ; May to October, observations at the Survey Office for 1866-79, furnished by Mr. Hennessy, F.R.S.
- LANDAUR :** Registers kept at the Convalescent Hospital in 1852-54, and 1866-67. Observations for the three years 1877-79 have been communicated by the medical officer in charge, but the means deduced from them appear to be 4 or 5 degrees too high.
- KALSI :** Figot—*Calcutta Journal of Natural History*, Vol. VI., 1837-38.
- ALNORA :** Observations at the Regimental Hospital. 1852-54 and 1866-69.
- HAWALBAGH :** Schlagintweit, page 494, on authority of Mr. Batten ; year unknown.
- LONCHIT :** Clelland's *Geology of Kuluon*, page 179 : parts of 1830, 1831, 1834, and 1835.
- KARDONG :** Schlagintweit, page 513 ; 1853-56.
- KANAM :** Cunningham's *Laddak*, page 181 ;¹ 1827-28.
- NPITI VALLEY :** Ditto, page 183 ;² 1846.
- BRIXAGAR :** { Manuscript observations by General Strachey and his brothers,
PAWET : { 1847-49.
XITI : }

The figures for the regular meteorological observatories (printed in small capitals in table III.) and those for the observatory at

¹Recomputed from the maxima and minima and corrected by means of the observations of Chakrata and Leh. ²Recomputed and corrected by the Leh observations.

the Survey Office, Mussoorie, are either directly calculated from observations at 4 A. M., 10 A. M., 4 P. M., and 10 P. M., or have been corrected to represent the means of observations taken at these hours. They probably differ very little from true daily means. The mean temperatures for the other places have been calculated in various ways, and many of them are open to considerable doubt.

The mean annual temperature diminishes pretty regularly as height increases. In the table there are only two exceptions to the rule that the higher a place is the cooler it is. It will be seen also on comparing places of nearly equal altitude and not very far apart that the highest temperatures belong to those which, lying behind the outermost high ridge, are subject to a much smaller rainfall than stations situated on the ridge or in valleys opening out to the south and exposed to the full force of the rainy winds. Thus Ránikhet and Almora are too hot in comparison with Naini Tál. The difference in temperature as well as in humidity between places situated at equal heights on the outer and inner ranges of Kumaon is sufficiently great to be easily recognizable without the aid of instruments, and is well known to all who have ever resided in the hills. The variation of temperature between the hottest and coldest months and the daily range of the thermometer are also greater, as a rule, in the interior than on the outer hills, owing to the larger proportion of cloudy sky and greater humidity of the air in the latter region.

Both the diurnal and the annual range of temperature decrease on ascending from the plains up to a height of 6,000 or 7,000 feet, and beyond that they again increase, their greatest values being attained at the highest stations where observations have been made. These places, however, lie to the north of the Indian watershed, where the humidity is doubtless less than on the southern side, and the observed ranges of temperature are probably higher than they would be at equal altitudes on this side of the snowy range. Moreover, the annual range in Tibet and Ladák is greater than on the Indian side of the chain on account of the difference of latitude, as has already been pointed out. In table IV. the daily and yearly ranges of temperature at twelve places are compared, and from it these relations will be readily seen.

IV.—Mean diurnal and annual ranges of temperature at places in the Hindlaya.

Place.	DIURNAL RANGE.													Annual range of monthly means.
	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.	
Bareilly	26.3	25.5	28.0	31.0	28.3	22.1	14.2	14.2	15.0	24.3	30.4	28.0	23.9	33.0
Roorkee	26.5	24.6	28.3	31.4	28.9	22.2	14.5	14.6	17.2	27.8	32.9	29.3	24.8	34.1
Dehra	23.4	23.1	24.0	26.3	24.4	18.6	12.8	12.5	15.0	23.4	25.9	24.3	21.2	30.1
Dharmasala	15.8	18.6	20.4	21.3	21.6	18.2	12.6	12.6	15.4	19.6	20.2	18.5	17.9	31.5
Mussooree, (1),	14.0	13.2	14.0	16.0	19.3	15.4	9.4	8.0	10.3	13.6	16.3	12.3	13.2	25.2
Ranikhet	16.1	17.0	18.2	19.6	18.1	15.0	11.2	10.9	13.1	15.3	17.2	16.1	15.5	25.2
Mussooree, (2),	19.0	16.5	10.7	9.9	12.0	14.2	16.8	27.0
Chakrata	18.3	18.4	18.1	19.7	19.4	16.3	12.5	11.3	12.0	17.3	18.0	18.3	16.6	26.4
Landaur	15.3	13.7	18.2	19.7	20.0	21.2	13.1	9.8	12.2	16.5	17.0	18.2	16.6	28.8
Kanam	15.2	17.2	22.0	19.0	18.5	17.3	16.3	15.5	88.9	20.5	19.0	18.3	18.1	37.8
Leh	28.5	30.0	28.0	31.0	30.1	31.8	11.2	31.3	31.4	30.5	26.7	25.2	29.6	43.9
Spiti	19.3	27.7	29.2	18.6	22.0	29.0	32.1	31.3	28.4	28.0	18.0	32.2	28.7	44.5

The table shows clearly that the minimum range for both day and year is reached at Ránikhet and the lower Mussooree station—that is, about 6,000 feet above the sea. The dependence of the diurnal range upon the humidity of the air and the proportion of cloud at each station is distinctly brought out in the variations from month to month. At all the stations but Leh and the Spiti valley, which lie beyond the snowy mountains, the months of the year which are driest in India—April and May—have the largest daily thermometric range, and the most humid months—July and August—have the smallest. There is a secondary minimum of temperature range coinciding with a maximum of humidity in January, and a secondary maximum in the dry and cloudless month of November. At Leh, where the most frequent precipitation of moisture during the year takes place in winter, the range is somewhat greater in the summer than in the winter months.

Owing to the greater annual range of temperature on the plains than on the hills, the diminution of temperature in the first 6,000 feet of ascent is most rapid in the hottest months and least so in the cold season. Between Roorkee and Chakráta the difference is less than 11 degrees in December and more than 23 in May. In the clear still nights of the cold weather, especially in November and December, before the winter rains and snows set in, the nocturnal

loss of heat goes on almost as freely on the plains as on mountain peaks. It is thus not unusual to find the temperature of the exposed thermometer at Roorkee nearly as low as at Chakráta, and it very frequently falls lower at Roorkee than at Dehra, where the observatory is surrounded by trees. In December the mean temperature diminishes between Roorkee and Dehra at the rate of only one degree in 1,034 feet, while in May and June it falls one degree in 230 feet.

The low temperature of the plains in the winter season, especially in the morning, is doubtless due in part to the draining down of cold air from the mountain slopes through the river gorges. This, however, cannot appreciably affect the temperature of places at a long distance from the mountains, though it may have a very considerable effect on that of Roorkee, close to the foot of the Siwálíks. The minimum temperature of the day is thus one or two degrees lower on the average at Roorkee than at Dehra in the months of November and December, and in January the minimum temperatures of the two places are equal. In the mountain country itself it often happens for the same reason that the temperature of the air at the bottom of a valley is distinctly lower than on the adjacent ridges. A similar inversion of the normal variation of temperature with height has been noticed in Europe during calm weather in winter.

From March to June the absorption of heat in melting and evaporating the snow on the outer hills, and in evaporating the rain that falls there in frequent showers when no rain falls over the plains, keeps down the temperature; so that in May and the first half of June, when the plains are at their hottest, the decrease of temperature on ascending through 6,000 or 7,000 feet is more than twice as great as in December.

In the Panjáb, where the latitude is higher and the humidity of the air over the plains is never great, the annual range of temperature at places on the plains is higher than in the North-Western Provinces, while in the hills there is much less difference. The annual variation in the decrease of temperature with height is accordingly much more distinctly marked in the extreme west of the Himálaya than it is in Dehra Dún. The difference of

temperature between Ráwal Pindi and Marri is 19·5 degrees in July and only one-third as great in December. On the other hand, in the eastern Himálaya, where the air at all elevations up to 9,000 or 10,000 feet is nearly equally moist, and where the range of temperature, especially over the plain, is much less than in the north-west, the decrease of temperature with height is most rapid in February and March and least so in June and July. The slow rate of decrement in the rainy season is doubtless due to the liberation of latent heat in the condensation of vapour; this heat rendering the atmospheric strata in which condensation occurs warmer than they otherwise would be, while the constant precipitation of rain prevents the lower strata from becoming very much hotter than the rain drops which pass through them. The effect of the rainy season in retarding the fall of temperature as we ascend is distinctly seen between Bareilly and Naini Tál or Ránikhet, but is not seen between Roorkee and Chakráta.

The great annual range of temperature at more elevated stations, especially such as lie behind the first snowy range and receive little or no precipitation, causes even greater differences in the rate of decrease of temperature with height, but in the opposite direction. In January, the difference of temperature between Chakráta and Leh is 24 degrees, but in August it is only 4 degrees. The greater length of the day in summer at Leh, and the absence of cloud to obstruct solar radiation on the surrounding mountain sides, render the summer months at that station, 11,500 feet above the sea, as hot as they would be on the southern side of the snowy mountains at an elevation of 8,500 or 9,000 feet. If General Cunningham's figures for the temperature of the Spiti valley are to be trusted, the heat of summer at an elevation of 13,000 feet is still more excessive. The relation of this to the greater height of the snow-line on the northern than on the southern side of the Himáláyan system is obvious.

In the following table the mean rate of decrease of temperature in the first 6,000 or 7,000 feet of ascent has been worked out for each month. In Dehra Dún the lower station is Roorkee and the upper one Chakráta; in Kumaon, Bareilly has been taken for the lower station, and instead of choosing either Naini Tál or Ránikhet for the upper one, the monthly mean temperatures of both

places have been taken and assigned to the mean elevation of the two. This was considered preferable to taking the figures for either hill station alone, because Ránikhet appears to be a little hotter than the average of places at the same elevation, and Naini Tál is probably somewhat cooler than the average.

V.—Decrement of temperature with height in the Himálaya.

Month.	Dehra Dún, difference of elevation, 6,165 feet.		Kumaon, difference of elevation, 5,008 feet.		Mean rate of decrease.	
	Difference of temperature.	Height for 1°.	Difference of temperature.	Height for 1°.	Height for 1°.	For 1,000 feet.
	°	Ft.	°	Ft.	Ft.	°
January ...	15·1	408	12·9	442	425	2·85
February ...	17·4	354	15·7	363	358	2·79
March ...	20·1	307	17·1	333	320	3·13
April ...	22·9	270	20·6	278	274	3·65
May ...	23·3	265	22·2	257	261	3·83
June ..	22·8	270	20·1	283	277	3·61
July ...	20·7	298	16·7	341	319	3·14
August ..	20·0	308	10·7	341	325	3·08
September ...	19·7	313	17·1	333	323	3·09
October ...	18·0	343	16·3	350	356	2·81
November ...	12·8	482	12·3	463	473	2·11
December ...	10·9	567	10·2	559	563	1·78
Year ...	18·6	331	16·5	345	336	2·96

In Dehra Dún there is a regular annual periodic variation in the rate of decrease of temperature with height, but in Kumaon the regular variation is interrupted in July and August, when the rate of decrease is slightly retarded by the fall of rain, as has been explained above. When the mean for both districts is taken, the regular variation from month to month is only slightly broken in August.

The figures given in the table include not only the decrease of temperature due to increase of elevation above the sea, but also a certain diminution caused by an increase of latitude equal to about a degree in 6,000 feet of elevation. The change of temperature with latitude in the Himálaya is small and to some extent masked by the contrast between the sunny valleys of the interior and the cool and cloudy outer ranges, but nevertheless it exists. On the plains of the North-West Panjáb the temperature falls as the latitude increases at a mean rate of about 1·1° F. for each degree of latitude, when corrections are made for differences of height above the sea. Probably much the same rate obtains in the western Himálaya at moderate elevations. The mean temperature of Yárkand, in latitude

39° N. and 4,200 feet above the sea, appears from the observations of Captain Trotter, R.E., of Drs. Bellew, Henderson and Souilly, to be about 54°F. At the same elevation in Kumaon the mean temperature is between 65° and 66°, and since the difference of latitude is ten degrees, the temperature appears to diminish about 1·1° or 1·2° for a degree of latitude. There is also a certain variation of the mean temperature with the longitude, places situated towards the east of the chain being cooler than those towards the west on account of the cloudiness of the summer months. In order to determine the true variation of temperature with height it is necessary to make allowance for these variations in latitude and longitude. The mean annual temperature of any place in the western Himalaya may thus be looked upon as a function of four quantities—(1) the sea-level temperature at a point taken as the zero of latitude and longitude, (2) a constant multiplied into the height of the place above the sea, (3) a constant multiplied into the latitude, and (4) a constant multiplied by the longitude. From the Schlagintweits' work and the reports of the Indian Meteorological Department fairly trustworthy mean temperatures for about thirty places in the Himalaya between the Nepál frontier and the Indus can be obtained, more than half of these being given in Table III. When treated by the method of least squares the observations admit of being thrown into the form $T = 78\cdot5^\circ - \cdot00277h - 1\cdot20 (\lambda - 29^\circ) - \cdot105 (L - 73^\circ)$, a formula which represents the observations with a mean error of about three-quarters of a degree. It ignores, of course, the differences of temperature between such places as Almora or Ránikhet and others of equal elevation on the outermost range. The decrement of temperature with height when latitude and longitude remain unchanged appears therefore to be 2·77 degrees in 1,000 feet or 1° in 361 feet. In the eastern Himalaya the decrease is more rapid, the observations taken at Darjiling and Gwálpára giving a mean rate of 1° in 320 feet. At great elevations on the table-land, too, it is probable that the temperature diminishes more rapidly than on the southern slope of the mountains. General Cunningham's observations in Spiti and Rápuhu during the month of September, 1847, give a mean increase of about 280 feet in elevation for one degree of fall in temperature; and from the observations taken by Dr. Souilly on the return journey from Yárkand over the Karakoram

pass and through Ladák in September, 1875, Mr. Blanford has deduced a mean fall of temperature equal to one degree in 227 feet.¹

The isotherms for equal altitudes in the western Himálaya and Ladák run from about W. 6° N. to E. 6° S., being thus three points less inclined to the parallels of latitude than the general direction of the mountain axis; but in eastern Tibet, Nepál, and Sikkim they probably bend to the southward. The mean temperatures of Káthmandu and Darjiling, at elevations of 4,354 and 6,912 feet above the sea, appear from the observations of ten years to be 61·7 and 53·9 degrees respectively, but according to the preceding formula they should be 66·7° and 60·1°. The mean temperatures of the four stations, Darjiling, Káthmandu, Gwálpára, and Sibságar, in the eastern Himálaya and Asám, may be represented very accurately by the formula $T = 77·7° - ·00312h - 1·53 (\lambda - 26°) - ·23 (L - 85°)$. In this part of the chain the temperature decreases at a mean rate of one degree in 321 feet of ascent, and the isotherms run from W. 10° N. to E. 10° S., the direction of the mountain axis being nearly due east and west. The isotherms are thus slightly curved, with the concavity towards the south-west, whilst the general line of the mountains is considerably curved in the opposite direction. This is merely another form of the statement in page 209, that the highest temperatures characterize that part of the chain between the Nepál frontier and the Satlaj.

Supposing the uniform rate of decrease worked out in Table V. to hold good for the whole southern slope of the North-West Himálaya, since the difference of latitude is nearly propotional to that of height, a mean temperature of 49°F., equal to that of London, would be attained at a height of 9,600 feet, and the annual range of temperature would probably differ little from that observed in England. The hill sanitaría, lying between 6,000 and 7,000 feet, possess climates comparable, as regards temperature, to those of Nice, Mentone, and other health resorts in the Riviera; all the towns along the coast of France and Italy from Marseilles to Florence having mean temperatures between 58° and 60°F. The annual range of the monthly means at the Himálayan stations does not exceed 25 or 26 degrees, whereas on the Mediterranean coast it varies from 28 or 29 degrees at Nice to 35 or 36 at Florence and Rome.

¹ *Indian Meteorological Memoirs*, Vol. I., page 233.

Nico and Ránikhet have exactly the same mean temperature, but Nico is seven or eight degrees hotter than Ránikhet in July and August, and several degrees colder all through the winter and spring months, except in January, when the temperatures of the two places are nearly equal.

At the superior limit of natural forest in the Himálaya, about 12,000 feet above the sea, the mean temperature is probably ten degrees above freezing. In the Alps a species of pine, *P. cembra*, forms natural forests on the borders of perpetual snow, where the mean temperature is several degrees under the freezing point. This difference of habit between the Himálayan and Alpine pines is very curious, and it is difficult to suggest any reason for it, since the natives of the Himálaya and Tibet find little difficulty in growing poplars and fruit trees in sheltered situations up to 13,000 feet or higher.¹ A mean temperature of 32° would be attained at a height of 15,400 feet, which is 2,000 feet above the upper limit of cultivated trees in Tibet.

The zone which has a mean temperature of 32° in any month will probably be near the lower edge of the snows in that month, especially in summer, when the light falls of snow on the outer hills have been all melted away. If the height of this zone be calculated for every month, the highest value obtained will be near the perpetual snow-line. With the uniform rate of decrease given in the last column of Table V., the result for July and August, when the snow line is highest, is about 17,630 feet. This result is very much higher than that given in Humboldt's *Asie Centrale*, on the authority of some of the earliest European travellers who penetrated into the country. General Strachey has, however, shown² that some of these mistook the lower limit of glaciers for the line of perpetual snow. The elevations of the snow line on the Trisúl and Nanda Devi groups of peaks, determined by trigonometrical observation from Almora in the latter part of 1848, before the winter snows set in, varied, according to Strachey, from about 17,000 feet on the eastern face of each group to 15,500 on the west; this latter was, however, in part probably newly fallen autumnal snow. The conclusion from these observations was that the height of the snow-line on the "more prominent points of the southern edge of the belt (of snowy mountains) may fairly be reckoned at 16,000 feet at the very least." The Schlagint-

¹ Schlagintweit's *Meteorology of India*, page 565.

² *Journal of the Asiatic Society of Bengal*, April, 1849.

weits found that the average height of the snow-line on the southern face of the Himálaya from Bhután to Kashmir was 16,200 feet. At page 72 of General Cunningham's book on Ladák the mean height of the snow-line on the peaks north of Simla, determined trigonometrically, is given at 16,665 feet, the highest point observed being about 1,000 feet higher; and in Mr. Drew's volume on the Kashmir territories, published in 1875, the elevation of the snow-line on the outermost zone is stated to be 16,500 feet. The height obtained by calculation as above suggested is thus probably a little too great, though it comes surprisingly close to some of Strachey's observations on Nanda Devi and those of Cunningham on the peaks of Kulu. Some observations have recently been made by the Trigonometrical Survey officers at Mussooree to determine the height of the snow-line on the peaks above Jamnotri, but the results have not yet been published.

For the ranges north of the Indian watershed we have not sufficient data to calculate the approximate height of the snow-line from temperature observations; but the high temperature of Leh and the Spiti valley in July and August show most distinctly that it must be much higher than on the southern side. The limit of perpetual snow on the ridges bordering on Tibet, especially those which lie beyond the Satlaj, is given by Strachey as 19,000 feet at least. Dr. Gerard many years ago, and more recently Mr. Drew, assigned an elevation of 20,000 feet to the limit of the snow in Rúpshu. One reason why this limit is so high—the intensity of solar radiation in summer—has already been mentioned; another is the insignificant quantity of snow that falls each winter in these regions that are almost completely cut off from the great southern vapour currents.

The lower limit of the snow in winter is usually about 6,500 feet in Kumaon and somewhat lower in Dehra Dún and the hills north of the Panjáb. While it lies the temperature does not rise much above 32°, but it seldom falls in sufficient quantity to lie more than a few days at a time. About one year in ten the snow comes down as low as 5,000 feet. The lowest level attained in the first half of the present century was about 3,000 feet in 1817 and 1847. In 1877 and 1878, though the snowfall was heavier than it had been for many years, it did not come down much below 5,000 feet. A slight fall of snow is said to have been observed at Lahore on the 12th of January, 1874,¹

¹ *Pioneer*, 17th January, 1874.

but no notice of this unprecedented occurrence was taken in the meteorological report for that month.

One of the most important effects of solar heat upon the atmosphere is the disturbance of its statical pressure relations, which in turn gives rise to those movements of adjustment towards equilibrium that are recognised as winds. The diurnal heating and cooling of the air causes certain oscillatory variations of pressure called the barometric tides, and gives rise to wind movements, such as the land and sea breezes and certain mountain winds that will be described below. The great annual change of temperature between summer and winter, in like manner, causes an annual variation in the height of the barometer and sets up those great currents of the lower atmosphere that are called the monsoons. Since the temperature is constantly changing no such thing as a simple statical distribution of pressure ever exists, though the averages of long series of observations may approximate more or less nearly to what a statical distribution would be. The air is constantly in motion either horizontally or vertically, and these movements cause variations of pressure, just as variations of pressure produce movements. Cause and effect, as regards pressure variations and winds, are thus so inextricably mixed up that it is next to impossible to disentangle them and show their relations clearly.

There can be little doubt that both the daily and the yearly inequality of pressure grow less as we ascend; and the annual variation at least becomes quite altered in character at a moderate elevation; but since the barometric variations depend upon the range of temperature which is possibly very great at great altitudes, while at a height of 6,000 or 7,000 feet it is less than on the plains, the decrease of the pressure variations with height is not strictly proportional to that of the total pressure. Table VI. gives the mean monthly pressures at a number of stations, and Table VII. the average daily range between 9-30 or 10 A.M. and 3-30 or 4 P.M. The figures for Bareilly, Roorkee, Dehra, Mussooree (both stations), Ránikhet, Chakráta, and Leh, have been corrected for the index errors of the barometers and are thus comparable with each other, except in so far as the different lengths of the periods of observation may cause the averages to differ; the others involve an unknown barometer error, which does not, however, affect the range of pressure, either diurnal or annual.

VI.—Monthly and annual means of Pressure at places in the Himalaya.

Place.	Elevation in feet.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.	Num-ber of years.
Bareilly	563	29-417	29-366	29-363	29-148	29-003	28-924	28-831	29-002	29-027	29-263	29-393	29-489	29-187	11-12
Boothoo	887	29-123	29-037	29-967	29-856	28-744	28-615	28-639	28-694	28-796	28-969	29-090	29-182	29-868	12
Debar	2,332	27-734	27-639	27-633	27-533	27-463	27-346	27-346	27-403	27-496	27-638	27-740	27-763	27-667	12
Pharmania	4,485	26-478	26-449	26-442	26-429	26-361	26-318	26-296	26-333	26-465	26-545	26-624	26-620	26-446	9
Almora†	5,546	24-704	24-633	24-668	24-588	24-543	24-423	24-434	24-463	24-678	24-710	24-771	24-731	24-603	1
Mussoorie (1).	5,835	24-267	24-264	24-304	24-260	24-211	24-133	24-104	24-143	24-317	24-390	24-331	24-323	24-236	1-3
Ranikhet	6,069	24-096	24-076	24-070	24-086	24-010	23-933	23-926	23-959	24-023	24-106	24-157	24-134	24-046	5-7
Naini Tal	6,462	23-965	23-834	23-858	23-833	23-760	23-687	23-698	23-716	23-753	23-879	23-905	23-894	23-807	5
Mussoorie (2).	6,831	23-834	23-267	23-249	23-294	23-366	23-439	23-492	23-438	...	1-14
Chakrata	7,052	23-259	23-220	23-243	23-236	23-194	23-163	23-113	23-166	23-325	23-268	23-308	23-287	23-223	4-5
Leh	11,508	19-804	19-578	19-640	19-649	19-668	19-660	19-616	19-640	19-692	19-723	19-742	19-701	19-639	4-6

† Upon one year's observations by General Strachey.

The double barometric tide that occurs regularly every day, especially in tropical countries, is still one of the least understood of atmospheric phenomena. It has been observed at all elevations in the Himālaya to which barometers have been carried, and with no considerable difference of phase,¹ though the range and general form of the oscillation are not the same at different heights. It is thus probably something of the nature of a wave of expansion and contraction propagated upwards and downwards with a velocity equal to that of sound.

The amplitude of the daily tide bears an obvious relation to the diurnal range of temperature, for it is greater over land areas than over the sea, and in table VII. it is seen to decrease pretty regularly from the plains up to stations situated at an elevation of 6,000 or 7,000 feet, where the range of temperature is least, becoming greater again at Leh, where the temperature range is large. Moreover, on the plains there is a well-marked annual variation of the daily range of the barometer, having its maximum in the hot-weather months, when the temperature range is greatest, and its minimum in the rainy season. But the inequality of temperature is not the only cause on which the observed barometric tides depend; for their amplitude, as well as their general form, varies with the season, the latitude and the position of the place with respect to the sea and to mountain ranges.

¹ Professor Loomis, in the *American Journal of Science* for 1879, finds that at the top of Mount Washington the daily maxima and minima set in three hours later than at the foot, though the difference of elevation between the two stations is only a little over 3,000 feet. This is quite unlike anything that occurs in the Himālaya, and is doubtless due to other causes than those which produce the diurnal tides. Hourly observations made in India show that up to a certain moderate elevation the daily barometric maxima and minima are slightly retarded; but this is due to the mountain winds described below. At Leh, in the upper Indus valley, the diurnal winds cause the morning maximum to occur nearly an hour earlier than on the plains. This will be seen from the following table:—

Place.				Height.	Morning maximum.	Afternoon minimum.
				Feet.	H. M.	H. M.
Calcutta	18	9 35	16 26
Allahabad	307	9 34	16 30
Hazārībāgh	2,010	9 47	16 24
Simla	7,971	10 26	16 51
Leh	11,503	8 48	16 29

The observed diurnal movements of the barometric column are in fact made up of several parts, only one of which is directly caused by the alternate heating and cooling which the air undergoes every day, though the others are all indirect effects of it. The direct effect of heating and cooling upon the pressure of the air has been more or less clearly explained by Espy, Davies, and Kreil, but it has not yet been shown by strict mathematical reasoning that their explanation accounts for the whole of the phenomenon. There is firstly, with a rise of temperature in the morning, an increase in the elastic force of the air, indicated by a rise of the barometer. But the increased elastic force immediately sets up a movement of expansion, either vertically or it may be in some cases laterally, by which the pressure is diminished. The actual movement of the mercury in the barometer is determined by the difference of these two actions ; and consists of a rise at first, up to 9 or 10 A.M., on the average, followed by a fall which goes on until some time after the hottest period of the day. It is easily seen that as long as the temperature continues to rise more and more rapidly, that is up to 9 A.M. or a little after, the first effect must outweigh the second, and the barometer will rise ; but as soon as the rate of increase of temperature begins to grow less, expansion will prevail and the barometer will fall. The expansion will not cease at the instant when the temperature reaches its maximum, but owing to the accumulation of motion it will go on for some time longer. Thus, there ought to be a barometric maximum about the time of most rapidly increasing temperature and a minimum in the afternoon. In much the same way it can be shown that there should be a maximum in the evening, when the temperature is falling most rapidly, and a minimum about the coldest time of the morning.

The coincidence between the barometric minima and the extremes of temperature is usually very far from exact, the barometer in this country standing lowest in the mornings about two hours before the time of minimum temperature, and in the afternoon about an hour and a half or two hours after the hottest time of the day. This may perhaps be explained on the principle of forced oscillations, that in the successive transformations which the energy undergoes, the oscillations approximate more and more nearly to simple harmonic waves with the maxima and minima separated by equal intervals. The

diurnal curve of temperature approaches more nearly to a simple wave form than that of the solar radiation which falls upon the earth at any place, and the double daily oscillation of pressure can be almost exactly represented by two waves superimposed.

At most places in India and the Himālaya the minimum temperature of the day occurs about sunrise, that is at 6 A.M. on the average of the year, and the maximum is attained about 2 P.M. The daily rise of temperature therefore occupies only eight hours of the twenty-four, and the fall the remaining sixteen. On the principle that the height of the barometer varies with the rate of change of temperature, the morning maximum should be much more decided than that of the evening ; and this is found by observation to be the fact, especially in the interior of India and other continental countries. Over the sea in tropical regions the periods of increasing and decreasing temperature are probably more nearly equal, and there the difference between the day and night waves of pressure is less.

At places near the equator the epochs of maximum and minimum pressure hardly vary from month to month ; but in higher latitudes the morning maximum and afternoon minimum approach each other when the days are short, and become more widely separated in the long days of summer.

Besides this primary oscillation of pressure caused by the heating and cooling of the air every day, the barometer indicates other changes due to the transfer of air by winds blowing to and from the place of observation, and perhaps also in some places it indicates other changes again due to the repetition of previous oscillations in the form of free waves. The great regularity and considerable range of the barometric tides over tropical seas where the daily range of temperature is small, may be thus to some extent caused by the repetition of the waves of previous days. In high latitudes, where the days and nights are usually of very unequal length and the variation of temperature is irregular, the tides become feebler, and near the pole disappear altogether, for in forced vibrations of any kind regular periodicity in the cause is an essential condition.

Near the coast the land and sea breezes modify the form of the diurnal pressure curves both at sea and on land. The transfer of air from sea to land during the earlier hours of the night renders the

nocturnal fall of pressure on land less, and at sea greater, than it would otherwise be, while the land breeze which blows in the fore-noons has a similar effect in the opposite direction. Over the water the morning minimum thus comes to be the lower of the two, in opposition to the usual rule.¹ Among the mountains a very similar semi-diurnal transference of air takes place, causing two distinct types of barometric tides—the one characteristic of valleys and the plains near the mountain system, and the other of high ridges and detached peaks. These types can be readily distinguished in the following table, which gives the variations from the daily mean at the hours nearest the turning points of the tides:—

	4 A. M.	10 A. M.	4 P. M.	10 P. M.
Bareilly	... —·021	+·060	—·047	+·008
Roorkee	... —·020	+·057	—·036	—·001
Ránikhet	... —·025	+·041	—·018	+·013
Naini Tál	... —·028	+·043	—·012	—·003
Chakráta	... —·022	+·036	—·015	—·001
Simla ²	... —·047	+·063	—·032	+·022
Leh	... +·011	+·037	—·049	+·001

At Bareilly and Roorkee, on the plains, the variation from the mean at 4 P. M. is twice as great as at 4 A. M. At all the Himálayan stations except Leh, which is in a valley between two ridges, this relation is reversed. Over the plains and on the outer hills, as at Naini Tál and Chakráta, the pressure at 10 P. M. hardly differs from the mean of the day. There is a small positive variation for this hour at Bareilly, and as we recede from the mountains and approach the sea the variation becomes greater; but there can be little error in concluding from the above table that along the southern border of the Himálaya the pressure rises in the evening just sufficiently to touch the mean. At Simla and Ránikhet, some twenty or thirty miles in towards the centre of the mountain system, there is, however, a well-marked evening maximum. This seems to indicate that the air continues to accumulate over the interior of the mountain zone for some time after the current has changed on the outer ranges and the air has commenced to flow back towards the plains.

¹ Blanford, J. A. S. B., Vol. XLVI., Part II., page 45. ² From General Boileau's observations in 1843-45. The daily range given by these figures is nearly twice as great as that of the other stations at the same altitude. The reason is probably some difference in the form of the barometer or in the mode of applying the correction for capillarity.

The transfer of air from the plains to the mountains in the daytime and its retransfer to the plains at night, which, by partly counterbalancing the afternoon fall of the barometer in the mountains and correspondingly increasing it on the plains, cause the peculiarities of the pressure variations seen in the preceding table, are brought about by the expansion and contraction of the air under the influence of heat and cold. In the daytime the air over the plains expands more than that over the hills, because the total thickness of air is greater and the range of temperature is probably higher. The surfaces of equal pressure, which we may assume to be horizontal on the average, are thus raised more above the plains than on the mountains, and the air under the influence of gravity flows down the incline towards the mountains. At night the air contracts and these surfaces sink more above the open plains than in the hills, and there is thus a slope or gradient in the opposite direction.

The following is General Strachey's description of the diurnal variation of the wind in Kumaon :—

“At most seasons of the year we find that on the Himálayan slope winds blow up the valleys during the day, that is from about 9 A. M. to 9 P. M., and down them during the corresponding hours of the night, or from 9 P. M. to 9 A. M. At the debouches of the principal streams into the plains, these night winds blow downwards with great violence, particularly in winter. In the interior of the mountains they are more moderate ; and at great elevations, and in the central parts of Tibet, the nights are almost always nearly calm. The diurnal currents from the south, on the other hand, increase in force as we ascend in height ; and along the Indian watershed and the neighbouring parts of Tibet they are excessively strong ; so that in travelling there, I have often looked forward to the afternoon, when they are at their height, with real dread ; and the natives of the country invariably endeavour to cross the high *passes* of the Indian watershed early in the day, for the purpose of avoiding the fury of the afternoon wind. As we advance further into the tableland, however, their power rapidly ceases.

“These winds, though on the whole blowing from the south-west during the day and from the north-east at night, that is perpendicular to the general line of the mountains, are naturally constrained

to follow the course of the deep valleys up which they pass, so that their direction is subject to endless local variation ; and, excepting on the tops of the hills, little information can be obtained by a register of the direction of the wind on the Himálaya, beyond the fact of there being an up or down current. In the part of Tibet I visited, near the Indian watershed to the north of Kumaon, the day wind seemed to commence in the south-east quarter about 9 A. M., and gradually to shift round with the sun as the day advanced, ending in the south-west quarter about 9 P. M. On several occasions in these localities I also noticed the wind blowing faintly from the north early in the morning. * * * * *

“The calm nights of the table-land and the higher mountains would (according to the theory above stated) be a consequence of their position in the centre of the mountain area, where the down current would originate, and therefore have the least force, though it be still felt in the faint northerly winds that are often observed near the Indian watershed.

“The violent night winds from the gorges by which the principal rivers leave the mountains would not appear to be altogether due to the same cause which produces the ordinary down winds, but to the accumulation of cold air in the deeper valleys to which I have before alluded. The air collected in these aerial lakes, as they may be called, having no means of escape but the openings through which the drainage is carried off, pours from them in a current the velocity of which will be dependent on the depth and area of the mountain basin from which it flows.”

General Cunningham also states that in Ladák and Spiti the southerly or south-westerly day wind usually begins about 9 A. M., the wind blowing faintly from the north about midnight and from the north-east in the early morning.

The day and night winds are probably strongest about 4 P. M. and 4 A. M., and the pressure and temperature observations made at these hours on the plains and at the hill stations indicate clearly enough that the direction of the baric gradient is from the plains towards the mountains in the afternoons, and from the mountains towards the plains in the mornings. When the pressures of Roorkee and Bareilly at 4 A. M. are reduced to the level of Chakráta and

Ránikhet respectively, and are corrected for any residual gradient to or from the mountains discovered by similarly reducing the mean pressures to the level of the hill stations, it is found that there is a pressure difference of $\cdot 077''$ at Chakráta and $\cdot 055''$ at Ránikhet, sending a wind towards the plains. At 4 p. m. the gradient is towards the hills, and is equal to $\cdot 062''$ at an elevation of 7,000 feet between Roorkee and Chakráta ; while between Bareilly and Ránikhet it is equal to $\cdot 045''$ on the average of the year. On the southern border of the mountain zone the gradient causing the down wind at night is therefore rather greater than that which causes the up wind during the day.

When the pressures of Roorkee at 4 A. M. and 4 P. M. are reduced to the level of Leh, a station beyond the Indian watershed, the gradients are found to be $\cdot 033''$ in the morning and $\cdot 182''$ in the afternoon. The pressure difference causing the day wind at great elevations thus appears to be nearly six times as great as that which causes the night wind ; but this relation is much exaggerated, no doubt, by the peculiar form of the pressure variation at Leh, which cannot be taken as a typical mountain station. This peculiar variation is doubtless due to the position of Leh in a narrow valley between two parallel mountain ranges. In the daytime the air of the valley expands and flows towards the mountains, and at night it again accumulates over the valley. In this way the nocturnal barometric tide is completely obliterated, and the afternoon fall of the barometer is rendered much greater than it would be on an open plain at the same altitude.

In April, May, and June the afternoon winds of the mountains blow with greatest violence, because in these months the range of temperature both on the plains and among the mountains is greatest. In these three months we find the afternoon fall of the barometer on the plains at a maximum, while at the hill stations it is less than in the cold weather. The nocturnal inequality of pressure is then at a minimum on the plains and valleys and at a maximum on the hills.

The annual variation of pressure differs from the diurnal in that no part, or an exceedingly minute part, of it is due to direct increase or decrease of elastic force accompanying gain or loss of heat. The

rise of temperature in the first half of the year and the fall in the latter half are accomplished so slowly that the increase or decrease of elastic force cannot accumulate, but is lost in expansion or contraction. The annual variation is thus almost entirely a secondary effect due to the movement of air both vertically and horizontally. When the temperature of the air over India changes the air expands or contracts, and the hypothetical surfaces of equal pressure widen out or come closer together than they were before ; and since the annual variation of temperature over the south of India is very small in comparison with that which occurs over the northern plain and in Central Asia, the vertical range through which these surfaces travel in the course of a year will be greater on the Himálaya than under the equator. In the cold weather, for example, the planes of 30, 29, 28, &c. inches are wider apart vertically over Ceylon than in Northern India, while at the end of the hot and in the rainy season the opposite relation obtains.

If there were no lateral movements of the air the pressure at a station on the plains would be nearly constant all the year round, while at the hill stations it would be least in winter and greatest in summer, because in the latter season a larger fraction of the total atmosphere than usual would be elevated above the place, while in the winter less than usual would lie above it. In winter, however, the planes of equal pressure in the upper regions of the atmosphere over India all slope towards the north, and down this slope winds blow, causing an accumulation of air over Northern India which renders the total pressure observed on the plains at that season greater than in summer. As regards mountain stations, it depends entirely on the height of the place whether the influx of air from the south will be more or less than sufficient to compensate for the contraction and sinking of the atmosphere in winter. At all the hill stations in Table VI. above 5,000 feet elevation there are indications of a winter minimum of pressure, though this is not the lowest minimum except at Leh, the most elevated station of all. There the pressure is least in the beginning of February, whereas at all the other stations, as on the plains, it is least in June and July.

During the cold weather winds are usually blowing out from Northern India towards the south along the surface of the ground at the same time that other currents are blowing northward in the

upper strata ; the apparent direction being modified in either case by the rotation of the earth on its axis and by friction against the ground surface. On the plain of the Ganges the conformation of the surface makes the lower winds have a north-westerly direction.

As the temperature rises the air over India expands and a larger and larger proportion of the total atmosphere is lifted above the level of the hill stations. In consequence of this the barometer at first rises at the higher hill stations ; and it simultaneously sinks over the plains and the lower hills owing to the outward movement of the air. As the season advances more and more air is removed from India by the strong day winds which blow in the hot weather as well as by the winds over the Indian watershed that have been already described, while but little is restored by the feeble night winds that come from the opposite quarters ; the barometer continues to fall over the plains, and the rise observed at the hill stations in spring is soon also changed to a fall, except at Leh, where the barometer continues rising until May. In the upper half of the atmosphere, that is, above the plane of 15 inches pressure, the summer depression of the barometer, which at Leh is feebler than that of winter, probably disappears altogether, and the barometer stands highest in the hottest season as it would do at all elevations if there was no transfer of air from place to place by lateral currents or winds.

When the temperature of Northern India is at its maximum in the latter half of June, the planes of equal pressure are widest apart, and they all slope towards the north in the lower half of the atmosphere. Winds consequently blow in from the sea towards the land in the lower strata, and there are possibly upper currents in the opposite direction, though the existence of such has not yet been established. This relation continues until the autumnal equinox, after which the temperature falls rapidly, and the atmosphere contracts and sinks so as to reproduce the conditions characteristic of the cold weather. The cooling of the air at this season, like the heating of it in spring, produces a differential effect on the height of the barometer at the hill stations, which again have a maximum of pressure in November.

When the effects of the two actions above described—the expansion and contraction of the atmosphere vertically, and the lateral

transference of air by winds—are borne in mind, some curious and at first sight inexplicable peculiarities of the annual variation of pressure become intelligible. For example, on the plains the barometer almost invariably stands higher in December than in January, though January is the colder month of the two. This anomaly at once disappears when we remember that the total pressure of the air on the plains, considered statically, is made up of two parts—that of the air from the plains up to the hill stations, and that of the air lying above the hill stations. The latter part appears from the observations of Leh to be greatest in the first fortnights of May and November, and least in the corresponding parts of February and August; and if the monthly means for any station on the plains or lower hills be subjected to harmonic analysis, the annual variation will be found to be very closely represented by two harmonic waves—one of annual period, reaching its maximum at the time of greatest cold in the beginning of January, and the other of six months' duration nearly coinciding in phase with the pressure variation at Leh. The amplitude of the first of these undulations, which is as much as six tenths of an inch at some places on the plains, rapidly diminishes as we ascend, and passing through a zero value at about 10,000 feet elevation, re-appears at Leh in nearly the opposite phase, the minimum falling in winter. The amplitude of the half-yearly oscillation increases slightly as we ascend, but it appears to vary with distance from the plains in a horizontal direction rather than with height. The observed pressure on the plains, being due to the superposition of the two waves, is highest in December—that is, between the dates when each wave separately attains its maximum.

The truth of this theory of the annual change of pressure may be more clearly seen from Table VIII., where the monthly variations of the barometric weights of three successive strata of the lower atmosphere from their annual mean values are compared with the simultaneous variations of temperature. The last double column gives the variations for the whole thickness of the atmosphere from the plains up to 11,500 feet above sea-level.

VIII.—Annual variation of pressure and temperature in the lower atmospheric strata over the Hindlaya.

Month.	Roorkie to Dehra, 1,345 feet.		Dehra to Chakrata, 4,820 feet.		Chakrata to Leh, 4,451 feet.		Roorkie to Leh, 10,516 feet.	
	Temperature variation. °	Pressure variation. "	Temperature variation. °	Pressure variation. "	Temperature variation. °	Pressure variation. "	Temperature variation. °	Pressure variation. "
January	... -17.0	+0.67	-15.3	+1.31	-18.2	+0.92	-18.2	+2.90
February	... -13.2	+0.46	-12.6	+1.25	-14.9	+0.79	-14.5	+2.30
March	... -4.6	+0.14	-5.3	+0.45	-7.5	+0.40	-6.4	+0.99
April	... +6.0	-0.18	+4.4	-0.29	+2.1	+0.26	+3.6	-0.21
May	... +12.3	-0.40	+9.9	-0.75	+7.3	-0.37	+9.6	-1.56
June	... +14.9	-0.52	+13.0	-1.31	+13.0	-0.80	+14.0	-2.63
July	... +10.0	-0.40	+8.9	-1.19	+14.7	-0.66	+13.3	-2.25
August	... +8.8	-0.30	+8.2	-0.99	+13.0	-0.45	+11.7	-1.74
September	... +7.3	-0.20	+6.7	-0.73	+8.7	-0.32	+8.4	-1.23
October	... +0.2	-0.01	+0.5	0.00	-0.1	+0.08	+0.2	+0.07
November	... -9.0	+0.29	-6.1	+0.92	-6.4	-0.01	-7.6	+1.20
December	... -15.9	+0.48	-12.0	+1.31	-11.3	+0.23	-14.0	+2.02

1 From the average of the three stations—Roorkie, Chakrata, and Leh.

The barometric weight of each stratum and of the whole thickness—that is, the difference between the observed pressures at the top and bottom—varies inversely with the temperature, and the one variation is as nearly as possible proportional to the other. The only exception worth noting is that in the month of November the statical pressure of the stratum between Chakráta and Loh is less than it should be according to the temperature figures. This anomaly, however, would probably disappear from the means of a longer series of observations.

The annual variation of the wind in Northern India is for the most part such as should accompany the pressure variations above described, according to the usually received “convection current” theory ; but there is one important feature of the winds of the plain that has not yet been satisfactorily explained—namely, the prevalence during the hot weather of strong north-westerly winds when the distribution of temperature and pressure should, by the theory of convection currents, give rise to winds with a southerly element. These “winds of elastic expansion,” as they have been called by Mr Blanford, actually blow sometimes from places where the mean pressure is low to others where it is slightly higher. They are the strongest winds of the year on the Indian plain ; they blow only in the daytime, and since there is no compensating current of any appreciable strength at night, they are probably the chief agency in that removal of air from Upper India which causes the great summer depression of the barometer. They are not confined to India, but are equally characteristic of Afghanistan ; and Colonel Prejevalsky encountered winds perfectly similar in everything except temperature in various parts of the Gobi desert and on the Alashán plateau north-east of Tibet. On the southern slope of the Himálaya these winds are sometimes met up to elevations of 6,000 or 7,000 feet, and when they blow the air is unusually dry and full of dust. At greater elevations, however, they are either not felt or become undistinguishable from the ordinary up currents that blow during the day.

IX.—Mean resultant wind directions at places in the Hindkya.

Place.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Number of years.
Bareilly	N. 57° W.	N. 54° W.	N. 56° W.	N. 56° W.	N. 82° W.	N. 85° E.	S. 66° E.	S. 64° E.	N. 81° E.	N. 62° W.	N. 54° W.	N. 65° W.	9-12
Roorkee	N. 71° W.	N. 51° W.	N. 76° W.	N. 78° W.	S. 21° W.	S. 6° E.	S. 40° E.	S. 34° E.	S.	S. 30° E.	S. 18° E.	N. 74° W.	10-13
Delhra	S. 72° W.	S. 67° W.	S. 79° W.	S. 74° W.	S. 58° W.	S. 66° W.	S. 67° W.	S. 56° W.	N. 61° W.	N. 64° W.	N. 78° W.	N. 89° W.	9-10
Dharmasala	S. 71° W.	S. 73° W.	S. 80° W.	S. 69° W.	S. 59° W.	S. 66° W.	S. 71° W.	S. 71° W.	S. 57° W.	S. 66° W.	S. 52° W.	S. 59° W.	4-5
Ratukhet	S. 81° W.	S. 54° W.	S. 57° W.	S. 63° W.	S. 63° W.	S. 75° W.	S. 77° W.	S. 72° W.	S. 61° W.	S. 53° W.	S. 56° W.	S. 60° W.	7-10
Naini Tal	S. 49° E.	S. 39° E.	S.	S. 57° E.	S. 49° E.	S. 40° E.	S. 41° E.	S. 44° E.	S. 44° E.	S. 41° E.	S. 38° E.	S. 41° E.	7
Chakrata	S. 69° W.	S. 19° E.	S.	S. 59° W.	S. 35° W.	S. 74° W.	S. 51° W.	S. 55° W.	S. 50° W.	S. 29° W.	S. 29° W.	S. 18° W.	6-10
Leh	S. 8° W.	S. 11° W.	S. 28° W.	S. 22° W.	S. 40° W.	S. 32° W.	S. 57° W.	S. 55° W.	S. 29° W.	S. 37° W.	S. 27° W.	S. 17° W.	4

From the preceding table it is seen that though the change of the prevailing wind from north-west to east or south-east at the commencement of the rains is very distinctly marked on the plains, no such change takes place at the hill stations. Even at the lowest of these, Dehra, the resultant wind varies only from about three points north to the same distance south of west. At all the higher stations the prevailing direction in every month is southerly or south-westerly, with modifications depending on the form of the ground,—at Naini Tál, for instance, the winds are generally south-easterly. The only notable variation of the wind direction is a deflection towards the east at Chakráta (also at Simla, Marri, and other stations on the north-western Himálaya) at the time when the winter snows and rains are heaviest. The cause of this has not yet been ascertained.

The wind direction at the hill stations changes so little from month to month because the winter monsoon is of no great vertical thickness, while that of the summer months extends to a much greater elevation than the highest station at which observations have been made. When northerly or north-westerly winds are blowing on the plains, the return current from the south-west is felt on the mountains at all elevations above the first few thousand feet; and when southerly winds blow over the plains, the return current, if it exists at all, lies at a very great altitude. The existence of this return current from the north during the summer monsoon may possibly be proved by cloud observations. Dr. Scully's observations on the way back from Yarkand in August, 1875, tell neither for nor against it, the resultant of all the wind directions observed at elevations above 14,000 feet being due west.

In the next table the vertical thickness of each monsoon current on the Himálayan slope has been computed approximately from observations made at pairs of hill stations in the north and south of India. The northern stations are Rookee and Chakráta, and the southern ones, Colombo and Newara Eliya in Ceylon. To render the figures directly comparable, the observed pressures at the hill stations have been reduced to the common elevation of 7,000 feet, and those of the lower stations to sea-level, as was done by Mr. Blanford in drawing up a similar table in the *Indian Meteorologist's Vade Mecum*, page 175.

X.—Vertical thickness of the Monsoon Currents.

Season.	Month.	Mean pressure at 7,000 feet.			Mean pressure at sea-level.			Neutral plane.	
		Ceylon.	Debra Dün.	Gradient send- ing wind to- wards north.	Ceylon.	Debra Dün.	Gradient send- ing wind to- wards north.	Approximate pressure.	Approximate elevation in feet.
		"	"	"	"	"	"	"	"
Summer. Winter.	November ...	23.388	23.353	+ .035	29.904	30.021	- .117	24.45	5,780
	December ...	23.387	23.337	+ .050	29.911	30.078	- .167	24.89	5,260
	January ...	23.379	23.310	+ .069	29.915	30.070	- .153	25.39	4,690
	February ...	23.389	23.270	+ .119	29.917	29.994	- .077	27.35	2,650
	June ...	23.318	23.179	+ .169	29.856	29.483	+ .373	17.96	14,130
	July ...	23.313	23.160	+ .183	29.871	29.508	+ .366	16.81	15,930
	August ...	23.311	23.205	+ .139	29.879	29.576	+ .303	17.80	14,370
	September ...	23.359	23.270	+ .089	29.897	29.684	+ .213	18.67	13,140

At 7,000 feet elevation the pressure gradient in both seasons is such as to send a current from south to north, while at sea-level it is only in the winter season that the wind blows from north to south. The neutral plane separating the lower wind current from the supposed upper return current is nearly 16,000 feet above the sea in the height of the rainy season ; but in the cold weather, especially in January and February, the neutral plane is below the level of the hill sanitarium.

The heights in the table represent only the approximate mean positions of the neutral plane for the several months. In reality its height is constantly fluctuating, and thus in the winter season it often sinks so low as to strike the Indian plain below the base of the hills. A moist easterly or south-easterly current then blows for several days at a time in Upper India, bringing the winter rains, while in Southern India the wind may be northerly. The prevailing direction of the wind on the plains is, however, always northerly in the cold weather, in accordance with the mean position of the neutral plane.

The pressure gradients both at sea-level and at 7,000 feet are much greater in summer than in winter. In the latter half of October and the beginning of November there is hardly any gradient either way, and at that time feeble winds and calms prevail. The velocity of the wind being directly proportional to the baric gradient (except perhaps in the case of anomalous currents like the "winds of elastic expansion" which blow down the valley of the Ganges in the hot weather), this velocity should be greater in the rainy season than in winter.

XI.—Monthly mean velocity of wind in miles per diem.

Station.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
Roorkee,	55.8	65.9	67.3	74.6	91.9	101.0	18.3	65.5	58.8	35.3	31.3	35.7	63.5
Bareilly,	70.6	95.4	95.1	110.4	117.2	134.5	105.5	75.9	71.6	41.0	42.3	51.6	84.3
Dehra ...	40.1	49.2	57.6	65.2	64.7	54.0	34.9	29.6	34.1	42.6	50.9	47.4	47.3
Ránikhet,	71.9	77.0	87.1	131.3	98.0	120.0	129.1	150.3	136.4	116.6	99.5	95.0	109.4
Chakráta,	114.5	114.6	129.4	137.0	135.8	128.1	110.1	88.5	100.3	116.1	105.9	103.3	117.7

Table XI. shows that on the plains and at Ránikhet (for which station more observations are required to get a good average) the wind velocity is least in winter, but that at Chakráta it is least in the rains. At the hill stations the winds are chiefly of the diurnal kind, and are feeblest when the temperature range is least—that is, in the month of August.

The quantity of water vapour present in the air at any time is a most important meteorological condition. It depends on the temperature, the distance from the sea or other evaporating surface, and the direction of the wind. These relations are very distinctly marked on the North Indian plain, where the high temperature range in the yearly period and the semi-annual change of the winds combine to render the proportion of vapour in the air during July and August nearly three times as great as in December and January. A high temperature cannot of course increase the quantity of vapour in the air, unless it be in a region where vapour is being generated. Accordingly we find in Table XII. that the vapour pressure at Roorkee, on the drier part of the plain, hardly varies from December to April, though as soon as the sea winds set in, which they sometimes do in the middle of May, the proportion of vapour rapidly increases. At Bareilly, where the surrounding country is moister and better wooded than at Roorkee, there is a slight increase of vapour in the hot-weather months.

XII.—Monthly mean tension of aqueous vapour at places in the Hindkoja.¹

Place.	Elevation in feet.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.	Number of years.
Bareilly ...	568	·354	·353	·413	·456	·570	·751	·953	·939	·866	·610	·422	·351	·585	11-12
Roorkee ...	887	·310	·336	·362	·360	·460	·677	·902	·903	·820	·543	·368	·324	·531	12-13
Dehra ...	2,232	·294	·325	·372	·418	·511	·721	·881	·878	·794	·321	·356	·301	·531	12-13
Dharmasla ...	4,455	·271	·281	·352	·439	·459	·539	·643	·627	·601	·444	·294	·274	·436	4-5
Mussooree ...	5,850	·202	·206	·249	·280	·322	·471	·643	·620	·621	·402	·281	·240	·378	1-3
Ranikhet ...	6,069	·190	·203	·235	·264	·341	·472	·557	·598	·536	·354	·253	·199	·353	6-7
Chakrata ...	7,052	·177	·191	·222	·265	·332	·455	·583	·582	·519	·844	·226	·187	·340	8-11
Kumaon and Garhwal ² .	8,000 to 9,000	...	·140	·220	·400	·280	?	...
Ditto ...	9,000 to 10,000	·420	·250	?	...
Ditto ...	10,000 to 11,000	·310	·350	·330	?	...
Leh ...	11,538	·103	·120	·134	·142	·167	·217	·339	·313	·246	·144	·115	·093	·177	4
Kumaon and Garhwal ² .	12,000 to 13,000	·210	·250	·320	?	...
Ditto ...	13,000 to 14,000	·200	?	...
Ditto ...	14,000 to 15,000	·140	·210	·200	·120	?	...
Ditto ...	15,000 to 16,000	·140	·130	·110	?	...
Ditto ...	16,000 to 17,000	·120	·180	?	...
Ditto ...	17,000 to 18,000	·130	?	...
Ditto ...	18,000 to 19,000	·120	·150	?	...

¹ These are the means of observations at 10 A. M. and 4 P. M. There is a certain diurnal variation of vapour tension, but its range is very small and it varies a good deal at different places. For most of the stations in the table the correction for this variation is unknown, and therefore all the figures have been given uncorrected.

² From miscellaneous observations by General Strachey.

The tensions in the foregoing table have been computed from observations of dry and wet bulb thermometers. The formula used for the most part has been Apjohn's ; but all the figures for Leh and Mussooree, and those for the last four years at the other stations, have been computed by means of certain tables based on August's formula.

On the southern slope of the mountains the annual variation of vapour tension is similar to that which obtains on the plains, though because of the considerable evaporation from the forest-covered slopes, and the occasional showers of rain which fall, the increase of vapour during the hot weather goes on much more uniformly than on the plains. At Leh, where hardly any precipitation occurs at any time of the year, but in the neighbourhood of which there is some cultivated land irrigated from the hill streams, the annual variation of vapour tension is determined almost entirely by the temperature. In the valley of Yárkand the quantity of vapour in the air is similarly determined by the temperature and the extent of irrigation.

In the mountains the mean vapour tension decreases very rapidly with the height, on account of the rapid decrease of temperature as we ascend. If Dalton's law, that in a mixture of gases or vapours the pressure of each is the same as if it filled the whole space alone, were applicable to the atmosphere, as is sometimes supposed even yet, then the pressure or tension of vapour observed on the plains ought to be reduced one-half on ascending through 29,000 feet; but it is found by observation that a vapour pressure equal to half that observed on the plains is attained at an elevation of 7,000 or 8,000 feet. This was pointed out by General Strachey in the *Proceedings of the Royal Society* for March, 1861, where he has shown that the observations of Mr. Welsh in balloon ascents, those of Dr. Hooker in Sikkim, and his own observations in Kumaon (most of which are included in Table XII.), make it perfectly certain that the proportion of water vapour which exists at any given elevation is determined, not by Dalton's law, but simply by the temperature. The vapour raised from the earth's surface is constantly diffusing upwards, and would go on doing so until it attained the state of equilibrium represented by Dalton's law ; but the temperature falls so rapidly as the height increases that saturation point is reached and the vapour is partially condensed into cloud or rain long before the barometric equilibrium is attained.

In the third column of Table XIII. the figures given in the previous table have been compared in a manner suggested by Strachey. The tension of vapour at sea-level under Kumaon and Garhwál has been computed for each month, by multiplying the mean of the observed tensions at Roorkee and Bareilly into the ratio between the tension of saturated vapour at the sea-level temperature and that of saturated vapour at the temperature of the plain; that is to say, the temperature is supposed to be corrected for elevation above the sea while the degree of saturation remains constant. The figures in Table XII. have then been divided by the corresponding tensions at sea-level, and the average of the fractions for all the months has been calculated for each elevation. Finally, from these results the ratio of the tensions at each even thousand feet above the sea has been found by interpolation. The second column of the table gives the results of Sir Joseph Hooker's observations in Sikkim compared with those taken at the meteorological observatory of Gwálpára near the foot of the hills, and the fourth column has been computed from the observations in General Cunningham's *Ladák* and those taken by Dr. Scully on the way back from Yárkand in 1875. The latter have been published in the *Indian Meteorological Memoirs*, No. VIII. The base station for the Kashmir group is Ráwal Pindi. The figures opposite 7,000 feet in the second and fourth columns are derived from the monthly means of the Darjiling and Marri observatories.

XIII.—Proportions of vapour tension at various elevations in the Hindúlaya.

Height.	Sikkim.	Kumaon.	Kashmir and Ladák.	MEAN FOR HIMALAYA.		
				Observed.	Computed by Strachey's method.	Computed by Hann's method.
Sea-level ...	1·00	1·00	1·00	1·00	1·00	1·00
1,000 feet ..	·89	·92	·92	·91	·91	·91
2,000 " ...	·83	·89	·85	·86	·83	·82
3,000 " ...	·74	·87	·69	·77	·75	·74
4,000 " ...	·67	·74	·52	·61	·68	·67
5,000 " ...	·64	·70	·51	·62	·62	·61
6,000 " ...	·59	·60	·57	·59	·56	·55
7,000 " ...	·53	·56	·60	·56	·51	·50
8,000 " ...	·50	·53	·41	·48	·46	·45
9,000 " ...	·44	·49	·28	·40	·42	·41
10,000 " ...	·40	·47	·27	·38	·38	·37
11,000 " ...	·37	·35	·28	·33	·34	·33
12,000 " ...	·33	·28	·21	·28	·31	·30
13,000 " ...	·30	·25	·23	·26	·28	·27
14,000 " ...	·27	·20	·19	·22	·25	·24
15,000 " ...	·27	·17	·15	·20	·22	·22
16,000 " ...	·22	·15	·16	·18	·20	·20
17,000 " ...	·20	·15	·14	·16	·18	·18
18,000 " ...	·19	·14	·15	·16	·16	·16
19,000 " ...	·18	·13	·14	·15	·14	·15

The mean of all three sets of observations probably represents very closely the actual diminution of vapour pressure on ascending in the Himálaya. It decreases with more regularity than either the Kumaon or Kashmir series of observations alone, and it agrees very closely with the series for Sikkim, where the relative humidity, or percentage of saturation, varies much less than in the western Himálaya. The last column of the table but one gives a series of ratios calculated on the assumption that the degree of humidity is the same at all elevations, and that the temperature of the southern slope of the Himálaya decreases at the mean rate above found—one degree in 361 feet. This series agrees very closely with the average of the results given by observation ; though from 2,000 to 10,000 feet the calculated ratios are all considerably less than those observed in Kumaon. During the hot-weather months the degree of saturation on the plains below Kumaon falls exceedingly low, while on the hills, as has already been stated, the air remains much moister ; at Dehra, for example, the vapour pressure in March, April, May, and June is greater than at Roorkoe owing to local evaporation. Thus on the average of the year the relative humidity of the air in the Kumaon hills is considerably greater than over the plain. On the other hand, the observed ratios from 14,000 feet upwards are less than those given by calculation, because most of the observations at these altitudes were made at places lying behind the snowy range.

In the last column are given the ratios calculated by the logarithmic formula, $\log p = \log P - \frac{h}{29088}$, where h is expressed in feet. Dr. Julius Hann, in an article in the Austrian Meteorological Society's Journal for 1874, page 193, has deduced from all the available observations on mountains and in balloons a similar formula in which the numerical constant is 6,517 metres, or 21,382 feet. On the assumption that this formula holds good to some distance beyond the limits of observation, we find that an elevation of 23,000 feet, or about the average height of the snowy peaks, the quantity of vapour in the air is only one-tenth of the quantity at sea-level. The extreme dryness of Tibet and Ladák is thus easily accounted for.

The logarithmic formula has the advantage of enabling us to calculate approximately the total quantity of vapour in the air at any time, by an application of the integral calculus. Using the generally received values for the density of water vapour and its

co-efficient of expansion with heat, and extending the integration to an infinite height above the ground, it is found that the depth of water that would be formed by the complete condensation of the vapour over a given area is almost exactly three times the height of the mercurial column which measures the pressure of the vapour at the bottom. In the rainy season for example, when the pressure of vapour over the Indian plain is equal to about an inch of mercury, the complete precipitation of the vapour would yield only three inches of rain, that is, less than the quantity which sometimes falls in two or three hours. A continuous downpour amounting to fifteen or twenty inches, such as frequently occurs in India, must be fed by a powerful indraught of moist air.

The relative humidity of the air is probably greater at all elevations on the Himálayan slope than either on the plains or on the Tibetan plateau beyond the Indian watershed; and it is doubtless greater on forest-clad slopes and valleys than on steep and bare mountain sides. On a high ridge, too, which intercepts and deflects upwards the prevailing south-west winds, thereby cooling them and partially condensing their vapour, the degree of saturation is greater than on the lower ridges or valleys behind it; for the air in sinking after crossing the high ridge is warmed and rendered capable of absorbing more moisture than it has been able to retain in crossing the ridge. Thus Naini Tál, independently of the influence of the lake, is always much moister than Ránikhet or the notoriously dry and bare station of Almora. The registers of the meteorological observatories do not, however, illustrate this very well; for at several of them observations have only been taken in the daytime, when the relative humidity is below the mean; and the humidities recorded at the old observatory of Naini Tál are quite untrustworthy and in many cases impossible. At Bareilly, Roorkee, Ránikhet, and Chakráta observations were taken both night and day for some years, at the hours of ten and four. If the means of the four observations at these hours be adopted as daily means, Chakráta appears to be the most humid of the four stations, and Roorkee and Ránikhet the driest, though the difference between Chakráta and Roorkee or Ránikhet is less than might be anticipated. The humidities of the other places in Table XIV. have been calculated approximately from the monthly means of temperature

and vapour tension. The figures for Leh in the winter months are doubtful; the psychrometer generally giving unreliable results when the temperature falls much below freezing.

XIV.—Approximate mean humidities of places in the Hindálaya.

Place.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
Bareilly ...	69	63	51	39	43	59	82	83	81	70	65	69	65
Roorkee ...	69	65	53	36	37	54	78	80	77	65	62	68	63
Dehra ...	69	69	58	47	45	63	86	90	86	69	62	67	70
Dharmśála ...	83	73	67	57	51	55	79	80	77	69	56	67	68
Mussooree ...	66	64	53	45	49	62	99	99	98	70	61	72	70
Ránikhet ...	60	58	52	39	49	61	85	86	79	60	53	54	62
Chakráta ...	68	64	56	49	57	68	94	94	89	64	49	58	67
Leh ...	100	100	79	57	52	50	62	66	68	63	67	71	70

The relative humidity of the air at all elevations up to 11,500 feet is subject to a double annual variation, one maximum occurring at the time of greatest cold, and the other in the middle of the rainy season. At Leh the summer maximum is very faintly marked, nine-tenths of the vapour brought by the south-west monsoon being cut off before reaching the station, and at Dharmśála, north of the Panjáb, the air appears to be slightly more humid in winter than in summer. At all the other stations the maximum degree of humidity is reached in August.

In April and November the air is dry, especially in the former month, when, during hot winds from the north-west, the percentage of saturation over the plains often falls as low as 5 or 6. In the hills, at Almora and Ránikhet, the humidity of the air frequently sinks to 25 per cent., but is seldom less than 20 per cent. The month of November and the beginning of December appear to be quite as dry as April at the higher hill stations, where these months are rainless, while showers sometimes fall in April; but on the plains, because of the low temperature of November and the moisture left in the ground by the summer rains, the air is still comparatively moist. In the cold weather the Sub-Himálayan stations are more humid than the hill stations on the average of the twenty-four hours, probably because the air, which is cooled and has its relative humidity increased by radiation during the night, drains away from the hills and collects over the plain. At this season the air at the hill stations appears to be drier in the mornings than in the evenings.

The humidity of the upper regions of the atmosphere, as indicated by clouds, is always greater in the daytime than at night. On the outer slope of the Himálayan chain the variation is doubtless quite as distinctly seen as anywhere else in the world, but it has not been recorded in the observatory registers, where only the amounts of cloud seen at 10 A.M. and 4 P.M. have been entered. The variation is, however, probably very similar to that which occurs over the plains, where the sky is most serene about 10 P.M. and most cloudy at the hottest time of the day, when the upward convection currents are strongest. In Table XV. the figures represent the means of the 10 A. M. and 4 P. M. observations, and they are therefore a little above the true mean for the day.

XV.—Average proportion of cloudy sky, in tenths of the expanse.

Station.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.
Bareilly ...	2.87	3.12	2.97	1.96	1.46	3.98	7.11	6.18	4.50	0.73	0.72	1.65	3.12
Roorkee ...	2.96	3.47	3.12	2.25	1.71	2.90	4.97	4.50	4.58	0.99	0.82	2.18	2.87
Dehra ...	3.73	4.07	3.55	2.50	2.75	4.02	7.50	7.00	4.83	1.08	1.33	2.74	3.76
Dharmasála ...	4.99	3.91	3.42	2.17	2.60	3.26	7.60	6.76	4.09	0.74	0.95	2.02	3.54
Ránikhet ...	3.66	4.42	3.78	3.34	3.58	5.41	8.58	8.42	5.64	1.69	1.31	2.99	4.40
Chakráta ...	4.23	4.75	4.44	3.42	3.75	5.14	8.89	8.57	6.79	2.28	1.74	3.62	4.80
Leh ...	6.32	6.38	6.27	5.78	5.77	5.50	4.89	4.68	4.44	4.19	5.12	5.91	5.44

The annual variation of cloud is similar to that of the relative humidity of the air near the ground. It has two maxima, in the cold weather and the rainy season, and two minima, in April or May and in November. April is cloudier than November, probably because the upward movement of the air during the day then prevails over the downward movement at night, while in November the prevalent movement is downward. In this way the air in the upper strata is dynamically cooled in the hot-weather months and dynamically heated in November. The variation of humidity at the hill stations is intermediate in character between that observed on the plain and the variation in the cloud-bearing strata of the atmosphere.

No direct observations of the heights of clouds above the ground have been made in the Himálaya. The ordinary clouds of the rainy season that look like broken cumulus from below are often not more than 5,000 or 6,000 feet above sea-level, hill stations like Naini Tál and Mussooree being frequently enveloped in them

for days. They sometimes even extend down to the level of the plains, the whole mass of the mountains up to the snows being then shrouded in fog. From the vapour tensions given in Table XIII., and the temperature decrements in Table V., it is possible to calculate approximately the average height at which a mass of air rising up from the plain would reach the dew-point and begin to form cloud. In January this height is a little over 4,000 feet above the plain, or about 5,000 feet above sea-level. In April and May the height above the plain is 8,000 feet, and at this time of the year it is rare to find clouds resting on the outer ridges of the *Himālaya*, though great banks of them are formed every day along the southern face of the snowy range. In the rainy season, that is, between the middle of June and the end of September, the average height at which clouds would commence to be formed in a rising column of air is 3,100 feet above the plain or about 3,900 above sea-level. This probably coincides very nearly with the zone of greatest rainfall on the mountains. In August, when the air is most humid and rainfall most frequent, the average lower limit of cloud is probably about 3,200 feet above the sea.

Regarding the upper limit of cloud nothing is known. The light feathery ice-cloud called *cirrus*, seen above the plains of Tibet and the passes over the Indian watershed, appears quite as high as when viewed from the Indian plain. It is probably formed at all elevations to which water vapour extends, though what the upper limit of vapour is we do not know. If we assume the *cirrus* clouds over the Tibetan plateau to be twice as high as the plateau itself, say 30,000 feet above sea-level, the quantity of vapour in the air would be only one-twentieth of that observed on the plains of India, but it would probably be quite sufficient to form light clouds.

The distribution of rain both on the plains and on the mountains has already been described in a general way.

Rain and snow.

The plains of Northern India, between the mountains and the *Jumna* river, or a line drawn north-westward from *Delhi*, beyond the river, may be divided into roughly parallel zones of equal rainfall, that which receives the greatest amount of precipitation lying nearest to the *Himālaya*. The breadth of each of these zones gradually diminishes towards the north-west and widens out in the direction of *Bengal*, because, the prevailing wind of the rainy season

being easterly over the plain, the supply of vapour gradually diminishes and the rains become lighter as we pass from east to west. In Table XVI, the average monthly rainfalls of 15 places on the plains near the base of the hills of Kumaon, Garhwál, and Dehra Dún, are given. The first group of stations is at an average distance of 20 to 30 miles from the base of the hills, and the distance of the other group is under 20 miles. In both groups the stations are arranged in order from east to west. The table might be extended indefinitely in both these directions as well as southwards, but little would be gained by doing so, since the distribution of rain above described is seen clearly enough from the figures as they stand.

The average rainfall of the line of stations at a distance exceeding twenty miles from the Himálaya is 40·1 inches, and that of the stations at a distance less than twenty miles is 46·6 inches. In each group the total rainfall gradually diminishes in passing from the extreme east of Rohilkhand to the neighbourhood of the Ganges, where it increases suddenly and again gradually shades off to the westward. The mean wind directions for the rainy season at Roorkee, Meerut, and Delhi indicate that there is frequently a sort of eddy formed at that season near the upper course of the Ganges, probably by the meeting of the south-east winds of the plain with south-west winds from the Arabian Sea that have been deflected northward by the Aravali hills in Rájputána, and this may be the cause of the increased rainfall that is observed.

On the mountains the rainfall varies rapidly with height, and its quantity is to a very great extent dependent on the situation of the place to the windward or leeward of high ridges and peaks. At fairly exposed stations of nearly equal altitudes there is a gradual diminution of the annual rainfall on passing from west to east, and between the Ganges and Jumna there is a slight increase perfectly comparable to that which occurs on the plain in the districts of Bijnor and Saháranpur. Thus the annual rainfall of Darjiling is 120 inches, that of Naini Tal 91 inches, that of Mussooree 95 inches, and that of Chakráta, Simla, and Marri 62, 68, and 58 inches respectively.

The next table gives the average monthly and annual rainfall of twenty places on the Himálayan slope, classified into three groups according to their positions near the foot of the slope, on the outer high ranges or on the inner ranges and valleys.

XVI.—Rainfall of stations on the plains near the base of the Hindlaya.

Station.	Elevation in feet.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.	Number of years.	
Nawdganj	...	640	1.28	1.15	0.55	0.27	1.29	5.82	16.94	11.79	8.02	1.51	...	0.49	49.11	16-17
Raheri	...	690	1.12	1.40	0.63	0.28	1.25	5.35	17.62	11.69	6.55	1.16	0.02	0.43	47.40	16-17
Tafardwana	...	720	1.35	1.63	0.73	0.36	1.06	6.95	14.02	11.71	5.91	0.79	...	0.63	44.34	16-17
Bijnor	...	800	1.14	1.22	0.86	0.34	0.70	4.81	10.47	9.82	5.41	0.53	0.05	0.40	35.45	31-33
Dooband	...	870	1.31	1.60	1.20	0.31	0.88	2.86	10.31	8.16	3.49	0.28	0.03	0.61	31.04	16-17
Sahampur	...	950	1.42	1.51	1.09	0.42	0.82	4.46	12.32	9.77	3.68	0.50	0.16	0.61	36.76	31-33
Ambda	...	820	1.41	1.38	1.01	0.63	1.03	4.18	11.53	8.75	4.57	0.84	0.14	0.71	36.18	25-26
Purampur	...	620	1.32	1.90	0.53	0.50	0.62	5.39	20.26	15.00	8.80	2.25	...	0.82	57.41	6-7
Pilibhit	...	650	0.99	1.41	0.45	0.37	1.15	6.25	16.17	12.06	9.72	1.31	...	0.48	50.36	16-17
Badarpur	...	720	1.18	1.42	0.80	0.32	1.16	5.74	15.11	11.30	5.39	1.02	...	0.47	48.91	20-21
Kashipur	...	750	1.48	1.53	0.79	0.49	1.43	6.05	12.53	11.70	5.29	0.83	...	0.69	43.81	16-17
Dhampur	...	780	0.96	1.84	0.74	0.58	0.95	4.22	13.12	11.87	5.29	0.50	...	0.68	40.70	16-17
Nagina	...	850	1.18	1.79	0.81	0.54	0.95	5.03	14.24	13.02	6.24	0.54	...	0.72	45.06	16-17
Najibabad	...	870	1.49	1.92	0.92	0.78	1.02	5.64	15.50	14.11	7.23	0.88	...	0.71	59.20	16-17
Bourtee	...	890	1.95	1.63	0.95	0.36	1.04	4.83	12.62	12.05	4.75	0.72	0.19	0.42	41.51	26-27

XVII.—Rainfall on the Hindlayam slope.

Station.	Elevation (in feet).	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Year.	Number of years.
INNER RANGES															
Champurāt	5,550	2-12	3-59	2-23	1-49	3-54	7-46	14-93	11-36	6-72	2-14	...	0-97	56-55	16-17
Pithoragarh	5,500	1-72	2-83	1-84	1-72	3-28	7-68	13-14	9-02	4-59	1-36	...	0-82	48-92	16-17
Almora	5,490	1-88	2-08	1-54	1-09	2-16	5-90	9-84	8-11	4-57	0-93	0-19	0-65	38-94	35-37
Ranikhet	6,069	1-64	2-61	1-60	1-62	3-11	5-81	13-69	11-11	5-05	0-78	0-08	1-18	48-28	9-10
Pauri	5,350	2-61	2-98	2-29	1-29	2-63	5-42	13-71	12-65	5-03	0-64	0-20	0-77	50-22	30-31
Shrinagar	1,750	2-20	3-44	2-34	1-05	2-25	4-35	8-80	8-05	2-71	0-66	0-02	0-72	36-63	16-17
Leh	11,358	0-10	0-01	0-01	...	0-05	0-03	5-20	0-71	0-21	1-52	3
OUTER RANGES															
Naini Tāl	6,600	3-04	3-24	2-34	1-78	2-97	15-42	24-60	23-77	10-24	1-80	0-15	1-59	90-84	30-31
Musoorree (St. Fidelia's School).	5,850	2-48	3-67	1-47	3-27	4-28	13-99	37-16	64-90	19-20	0-20	...	2-23	142-25	1-3
Musoorree (Hospital)	6,550	2-00	3-29	2-98	1-90	2-71	9-90	29-85	31-27	9-64	1-09	0-09	1-01	95-23	16-28
Chakrata	7,052	1-41	3-05	3-41	1-88	3-13	7-97	17-23	16-31	5-94	0-61	0-18	1-16	62-23	11-12
Landaur	7,511	2-47	1-44	1-68	2-49	4-60	7-24	26-41	30-86	7-59	0-62	0-68	0-93	87-13	5
BRIDGES, SWALES, AND DERHA DOK.															
Kilpari	800	1-32	1-72	0-84	0-56	1-35	10-61	19-86	15-63	8-79	1-45	0-01	0-50	60-12	16-17
Haidwāni	1,430	1-40	3-18	1-78	0-79	1-32	10-61	21-74	20-67	10-51	1-45	...	0-61	74-06	16-17
Hardwar	1,100	2-98	2-80	1-04	0-08	0-45	4-38	10-33	16-59	5-56	1-03	0-06	0-39	45-69	5-6
Mohan	1,330	3-65	2-91	1-82	0-83	0-79	9-23	19-86	17-48	6-70	1-72	0-58	0-50	65-07	4
Rhogpur	2,450	1-47	3-27	1-23	1-27	4-87	8-63	5-83	58-30	19-92	0-80	...	0-10	1-57-69	2-3
Dehra	2,230	2-06	2-10	1-41	0-73	1-39	8-32	24-61	23-66	9-14	0-75	0-07	0-52	74-86	31-33
Ambari	1,800	1-67	2-38	0-85	1-13	2-67	11-65	24-58	23-73	8-20	0-42	...	0-55	77-83	2-3
Kāsi	2,000?	2-21	3-42	2-22	0-96	1-54	8-02	25-09	26-91	10-04	0-89	0-04	0-99	82-53	16-17

1 Interpolated from upper station in some years.

The influence of an elevated ridge in diminishing the rainfall of the valley behind it is seen on comparing the rainfall of Almora with that of Naini Tál, or even by comparing Srinagar with Pauri, though both of these lie far in the interior of the mountain system. A much greater contrast is observable between Bhogpur, at the foot of the mountains overhanging the gorge of the Ganges above Hardwár, and Dehra, in the Dún, behind the central and highest part of the Siwálik chain. The rainfall of Bhogpur, given by the observations of two and a half years is, however, probably too high.

The variation of rainfall with height can only be roughly determined, because every high ridge and peak thus cuts off the supply of vapour from the lower ground to the north of it. In Table XVIII. an attempt has been made to determine it approximately from the rainfall figures in the first two sections of Table XVII., together with those of two or three places in the hills north of the Panjáb and the observations made by General Strachey at Niti in 1849. The ratio between the mean rainfall of each hill station and that of the nearest station or stations on the plains, for the same years, is given in the last column.

XVIII.—Rainfall of the outer slope of the Himálaya compared to that of the neighbouring plain.

HILLS.			PLAINS.			Difference of elevation.	Rainfall ratio.
Station.	Height in feet.	Rainfall.	Nearest station or stations.	Height in feet.	Rainfall.		
Kilpurí ...	800	80.1	Rudarpur ...	720	43.9	80	1.37
Hardwár ...	1,100	45.7	Roorkee ...	890	40.2	210	1.14
Pathánkot ...	1,160	50.2	Gurdáspur ...	900	30.1	260	1.67
Mohan ...	1,330	65.1	Roorkee ...	890	39.8	440	1.64
Ilaldwáni ...	1,430	74.1	Rudarpur ...	720	43.9	710	1.69
Ambári ...	1,800	77.8	Saháranpur and Ambála,	890	30.0	910	2.59
Káisi ...	2,000	82.3	Ditto ...	890	34.0	1,110	2.42
Nárpur ...	2,050	79.3	Gurdáspur ...	900	30.1	1,150	2.60
Dehra ...	2,230	74.9	Roorkee ...	890	41.5	1,340	1.80
Bhogpur ...	2,450	157.7	Ditto ...	890	30.0	1,560	5.26
Pálapur ...	4,000	118.0	Gurdáspur ...	900	28.7	3,100	4.11
Dharnisála ...	4,490	123.2	Ditto ...	900	33.1	3,590	8.72
Mussooree (1) ...	5,850	142.2	Roorkee ...	890	32.0	4,960	4.44
Mussooree (2) ...	6,550	95.2	Ditto ...	890	41.5	5,660	2.29
Naini Tál ...	6,600	90.9	Rudarpur ...	720	45.2	5,880	2.07
Simla ...	6,950	68.6	Ambála ...	820	36.2	6,130	1.90
Chakráta ...	7,050	62.2	Saháranpur and Ambála,	890	38.4	6,360	1.71
Landaur ...	7,510	87.1	Roorkee ...	890	42.8	6,620	2.05
Niti ...	11,460	5.5	Roorkee and Rudarpur,	800	45.0	10,660	0.12

¹ Rainfall of July, August and September, 1849; the observations at Niti taken by General Strachey.

By grouping together the ratios for the places lying between the even thousands of feet, many of the irregularities that appear in Table XVIII. are cleared away and the results may be accepted with more confidence. The excessive rainfall of Bhogpur, for example, will to some extent counterbalance the defect at Dehra caused by the position of the latter station behind the ridge of the Siwāliks. The following figures are thus obtained :—

Height above plain.			Mean height.	RAINFALL RATIO.	
				Observed.	Calculated.
Feet.			Feet.		
0 to	...	1,000	435	1.68	1.63
1,000	„	2,000	1,290	3.04	3.01
3,000	„	4,000	3,350	3.91	3.77
4,000	„	5,000	4,740 ¹	3.48	3.15
5,000	„	6,000	5,770	2.18	2.39
6,000	„	7,000	6,370	1.89	1.89
10,000	„	11,000	10,660	0.12	0

The ratios in the last column are calculated by means of a formula, $R = 1 + 2.12 h - 0.47 h^2 + 0.025 h^3$, given in the official *Report on the Rainfall of the North-Western Provinces and Oudh*, published in 1879. It was originally computed from somewhat different data, but it represents the observed ratios in the above table as closely as can possibly be expected, considering the nature of the observations. At elevations greater than 9,585 feet above the plain this formula gives increasing values for the rainfall, and is therefore inapplicable; but from 7,000 feet above the plain upwards the rainfall ratio may be approximately represented by a logarithmic formula, $\log R = 2.151 - 0.287 h$. In both formulæ h is to be expressed in thousands of feet.

The mean rainfall along Rohilkhand and the Doáb, at a distance of twenty miles from the hills, is about 43 inches, and the mean elevation of this line above sea-level is 800 feet. Applying the formulæ in the preceding paragraph to these data, we find that

¹ The observations for the lower Mussoree station being for a very short period only, the figures for the stations immediately above and below it in Table XVIII. have been included in striking the average.

the average rainfall on the southern slope of exposed mountain ridges in Kumaon and Garhwál would probably be the following :—

At	800 feet above the sea	48 inches.
"	1,000 "	60
"	2,000 "	125
"	3,000 "	159
"	4,000 "	164
"	5,000 "	149
"	6,000 "	122
"	7,000 "	88
"	8,000 "	52
"	9,000 "	27
"	10,000 "	14
"	11,000 "	7
"	12,000 "	4

From the table it appears that the maximum rainfall occurs about 4,000 feet above the sea. The exact height of the maximum zone determined by the formula is 2,948 feet above the plain, or about 3,750 above sea-level. This agrees very closely with the mean altitude at which a rising column of air reaches its dew-point in the rainy season.

By far the most important if not the only cause of rain in the Himálaya is the cooling of the air by expansion as it ascends the mountain slope. It has been already seen that in the rainy season, the direction of the wind at all elevations in the Himálaya up to 15,000 or 16,000 feet, if not higher still, is from some southerly quarter. Near the foot of the hills the prevailing direction is south-easterly, but at most of the stations from the level of Dehra Dún upwards the wind blows from some point to the west of south ; that is, more or less nearly at right angles to the axis of the mountain zone. The air in rising to surmount the barrier has its heat rapidly converted into the work of expansion, and it commences to precipitate rain when the temperature falls to the dew-point. When once condensation begins, the rate of decrease of vapour with height will be a measure of the quantity condensed or the rainfall. This rate is greatest at the lowest elevations ; and thus rain should be heaviest at places on the outer slope of the mountains where a rising stream of airs usually begins to precipitate moisture, as the observations prove to be the fact.

The slight rainfall of places like Almora and Srinagar, to the leeward of a higher mountain mass is caused by the partial exhaustion of the vapour in crossing the mountains and by the dynamical heating of the air as it streams down towards the valley, both causes diminishing the tendency to condensation. The rapid decrease of rainfall on ascending beyond 6,000 or 7,000 feet is due simply to the exhaustion of the vapour, but at all elevations the influence of high ranges in cutting off the supply of vapour is easily seen. Regarding the rainfall of 1849 on the Tibetan table-land, General Strachey says:—"In the country beyond Níti no register was kept; but during a week of rainy weather in the middle of August 1·5 inches fell at Níti, while at Sanjar, beyond the watershed, where I was then encamped, at 16,500 feet, the rain never exceeded a very faint drizzle, and could hardly have been susceptible of measurement." At the Leh observatory all through the summer the rainfall hardly ever exceeds a few drops, and the greatest fall in a month during several years was an inch and a half. Sir Joseph Hooker's experience in Sikkim supplies us with facts quite parallel to these. In August, 1849, he says 26·8 inches fell at Darjiling, while in the interior, at the same elevation, but in the rear of the first masses of snowy mountains only 12·5 inches were measured. Between the 8th of September and the end of the month only 1·7 inches fell at Mome Samdong, about 15,500 feet above the sea, while at Darjiling 10 inches fell, and other instances of a similar nature might be cited.

The variation of rainfall with season is very distinctly marked in India. At all the stations in Tables XVI. and XVII., and at almost every station on the North Indian plain, the driest month of the year is November. In the great majority of years no precipitation whatever occurs in this month or in the first half of December, except perhaps on the higher mountains towards the north-west where the winter snows usually begin before the end of November. About Christmas a few showers of snow usually fall on the outer hills, and at the same time there is a slight precipitation of rain over the plains of the Panjáb and the North-Western Provinces. These winter snows and rains increase in quantity and in frequency on the hills and in the north-west Panjáb until February or March,

but on the plains of the North-Western Provinces and Behar the maximum occurs in January.

The cause of the winter rains and snows has already been pointed out in describing the annual changes of the winds. In October and about the beginning of November the air over Northern India is as near as it ever attains to a condition of statical equilibrium. It is subject to the diurnal oscillations called the barometric tides and to the accompanying mountain winds, but there is little permanent movement of the air in any direction. During the cold weather, however, the neutral plane of pressure gradually sinks and the south-west upper currents of the atmosphere are then forced to ascend the slope of the mountains where they precipitate more or less of the vapour they contain. On the lower hills the temperature increases so rapidly in March and April that the tendency to precipitation, and consequently the rainfall, becomes less than in January and February, though the upper currents continue to blow from nearly the same direction as in winter. Along the southern declivity of the great snowy range, however, thunderstorms are of daily occurrence at this time of the year, and above the snow line considerable quantities of snow are frequently precipitated. In Ladák the heaviest falls of snow observed by Captain H. Strachey in 1848-49 occurred in April; but during the three years, 1876-78, there was no precipitation at Leh in that month.

April and May are the months in which hail is most frequently noticed in the Himálaya. No regular registers of this phenomenon have been kept, but nearly every year several hailstorms occur in the outer hills, and the stones are often of large size. On the 11th of May, 1855, a hailstorm occurred at Naini Tal in which many stones of 6, 8, 10 and even 24 ounces were observed to fall, the circumference of these varying from 9 to 13 inches. In 1878 there was a storm in which large hailstones fell, some of them so heavy that they punched holes through the zinc roofs of the houses, while the quantity was so great that it lay in shady places, where covered with leaves, for nearly a month.

About the middle of June usually, and sometimes before the end of May near the foot of the hills, the hot north-west winds of the plains give way to sea winds from the Bay of Bengal. The

whole of the lower atmosphere over India is then moving towards the Himálaya; and the upward deflection of the air currents by the mountains causes frequent precipitations of rain in the manner already described. The rainy season almost always commences sooner on the mountains than on the plains, for saturation is reached first at high elevations and then propagated downwards by the cooling effect of the falling rain drops and the cutting off of the sun's heat by clouds. At most places in India, and in the inner parts of the Himálaya, July is the rainiest month, but on the outer slope of the mountains August is equally or sometimes even more rainy, especially towards the north-west of the chain.

While the rainy season lasts, the parts of Ladák about Leh, and the Tibetan plateau generally, receive perhaps, on the whole, less precipitation than in winter, because the temperature is then so much higher than on the Indian side of the chain, this high temperature greatly decreasing the relative humidity of any air that may reach the plateau from the south. It is probable also that the high snowy peaks, lying above the limit of the monsoon current proper, receive less precipitation in summer than in winter. At the turn of the seasons, however, about the end of September, falls of snow amounting to several feet in thickness sometimes occur on the passes over the Indian watershed.

Shortly after the autumnal equinox, about the end of September or beginning of October, the rains cease all over Northern India. The sudden cessation of the rainy season seems to be in some measure determined by the rapid diminution of solar heat, as the sun retreats to the south of the equator. It is possibly to this that we must attribute the somewhat remarkable regularity of the recurrence of two or three days' incessant rain frequently experienced in Kumaon about the 20th of September. On the plains, also, it is well known to the natives of the country that if rain falls in the *nakshatra* (lunar mansion) of *Hathiya*—that is, in the last week of September or first week of October, it is likely to be heavy. Excessively heavy rain, like that of the 17th and 18th September, 1880, when 30 inches fell in little more than two days at Naini Tál and produced a disastrous landslip, cannot, however be thus produced by a simple loss of heat, but requires a powerful indraught of moist air to keep up the supply of vapour. The

heavy rain observed on the plains at "the break-up of the monsoon," and probably also that which falls at the same time on the outer hills of Kumaon, must be due chiefly to the minor storms of a cyclonic character that are frequently formed near the head of the Bay of Bengal at the turn of the season, and pass inland in a north-westerly direction.

CHAPTER VI.

ECONOMIC MINERALOGY.

CONTENTS.

Mining history. Experiments in Garhwál. The Kumaon Iron Works Company. Causes of present poor returns. Mode of mining: excavation: removing the ore: crushing the ore: cleaning the ore: drainage. Smelting operations: refining the ore. Division of profits. Improvement needed in appliances: Chili furnace: English system. Outturn. Financial results of the settlement of the revenue derived from minerals. Gold: Tibetan gold-fields: Thok Jalung mines. Silver. Copper mines: Rái mine and its ores: Sira mines: Gaul and Sor Garang. Bournonite. Copper mines of Garhwál: Dhanpur: Dhobri: Pokhri: Chaumattiya mines: Rája's mine: Nola mine: Thála mine: Danda mine: Tálapungla mine: Kharna mine: Agar Sera mine. Iron mines: Kumaon iron mines: Gíwár mines: Garhwál mines: Nágpur: Dasauli: Iriyakot. Lead. Arsenic. Lignite. Graphite. Sulphur. Borax. Gypsum. Soap-stone. Asbestos. Silajít. Limestone. Building stone. Roofing slate. Alum.

THE mineral resources of the Kumaon division early obtained the attention which their traditional value assumed to be due to them, and it was one of the directions to the first Commissioner to procure specimens of the ores to be found in Kumaon and transmit them to the mint for assay. Specimens of copper ore from the mines in Sira and Gangoli were accordingly forwarded to Calcutta in 1815, but the report was not favourable; for, if the specimens sent were fair samples of the ores in general, it was doubted whether the mines could be worked to advantage.¹ The Government were, however, not satisfied with this report, and in 1817 deputed Mr. A. Laidlaw as mineralogical surveyor to accompany Lieutenant Webb's party through Kumaon. His orders² were to consider the examination of the mineral resources of the country his primary duty, though at the same time he

¹ To Government, dated 26th June, 1815: from Government, dated 13th January, 1816. For further information on the mineral resources of Kumaon see *Moorecroft's Travels*, I., 7; *As. Res.*, XVIII., 236; *Rec. Geol. Sur.*, 1871, 19: II., 66: (*Lawder*) III., 43; *J. A. S. Ben.*, VI., 653; *Glean. in Sc.*, I., 230.

Graphite, *Glean. in Sc.*, III., 230; *J. A. S. Ben.* XXIV., 208; *N.-W. P. Rec.*, III., N. S., 371; *G. India Rec.*, XVII., 58.

Iron ores and works, *N.-W. P. Rec.*, XVIII., 1; *Rec. G. India, Sep.*, VIII., 37: XVII., 1: XXVI.: Bhábar iron works, *Agra*, 1856.

Copper ores and works, *J. A. S. Ben.*, VII., 934 (*Drummond*): VIII., 471 (*Glasford*): XII., 453, 769 (*Lushington*): XIV., 471 (*Reckendorf*); *As. Res.*, XVIII., 239; *N.-W. P. Rec.*, III., N. S., 23; *Glean. in Sc.*, I., 328.

Lignite, *As. Res.*, XVI., 387, 397.

² From Government, dated 11th July, 1817. Mr. Laidlaw died at Pithoragarh in 1836, and I have not been able to procure any of his original reports

should not feel himself debarred from bestowing attention on any other matters deserving of scientific research, so far as such investigation did not interfere with his more immediate duties. These instructions further go on to say:—"You should ascertain the existence or otherwise of mineral productions applicable to purposes of public use, or available as a source of revenue; and report on the practicability of bringing them to account. The existence of iron and copper ores in considerable quantity has already been ascertained; but as the working of these metals might injuriously affect important articles of British import, it is not designed that your attention should be occupied in detailing any practical arrangements for that purpose; you will not, however, consider yourself debarred from prosecuting enquiries into any circumstances regarding them which may be of sufficient interest to science to merit particular notice. In every part, indeed, of your researches it is the wish of Government that, in the first instance at least, you should contemplate rather the general capabilities of the country which you are to visit than the special means necessary for bringing them into action in any particular district; though, of course, the facility or difficulty with which metallic ores or other useful minerals could be raised and brought to market must be a leading point in your observations on them. The minuter details of machinery, mode of working, &c., are what it is meant to postpone, as these will be the subject of future determination, when the whole result of your survey shall be before Government."

In 1826 Captain J. D. Herbert submitted his report on the mineralogical survey of the province.¹ This was followed by a report² in 1838 by Captain H. Drummond, of the 3rd Light Cavalry, on the copper mines at Rái in Gangoli and at Sira in Bárabisi. Captain Drummond had brought with him from England an experienced Cornish miner, Mr. Wilkin, to examine the mines already worked, and proposed that, with a view of ascertaining their actual value, a certain sum should be advanced by "Government for an experimental opening of such mines as might appear best suited to the object in view." This proposal was accepted³ by Government, and a sum of Rs. 2,415 (subsequently increased by Rs. 1,000) was

¹ To Commissioner, dated 10th January, 1826.
934; and Stat. Kum., p. 268.

² J. A. S. Res., VII.,

³ To Commissioner, dated 26th November, 1838.

allotted to carry out the designs furnished by Captain Drummond. A report¹ on the experiments then undertaken was rendered by Mr. G. T. Lushington, Commissioner of Kumaon, in 1842.

The place selected for the important purpose of determining the advantages or otherwise of working the mines under European superintendence was the Pokhri copper mine in pargana Nāgpur in Garhwāl. The works were carried on from 1838 to 1841 with a net loss to Government of Rs. 7,384. The Commissioner considered this complete failure to be due to the poorness and scarcity of the ores found, and not in the least to any want of skill or industry on the part of Mr. Wilkin, or any injudicious selection of the places for experiment. Mr. Reckendorf, a mining engineer, visited these mines again in 1845, and in commenting on the deductions to be drawn from Mr. Wilkin's experiment, gave it as his opinion that the operations then undertaken should not be considered as conclusive against the value of the mines.² The experiment should have been confined to driving a shaft some thirty or forty fathoms below the old mines, and not to collecting ores which might have been a good addition in smelting other ores, but the smelting of which alone could never be profitable.

Again, there were no proper appliances for smelting, the loss from which by the native method adopted was very great, and the ores used, from their nearness to the surface, had already lost much of their value by the slow metamorphosis of pyrites into sulphate of copper. On the whole, Mr. Reckendorf's opinion was favourable to more extended and expensive operations in the hands of a private company. Nothing of importance, however, resulted from this the first attempt to obtain some accurate information regarding the mineral resources of the Himālaya. Captain Drummond also combated³ the conclusions arrived at by Mr. Lushington as to the mines not affording a fair field for investment of capital, and in support of his views quoted the testimony of Captain Glasford (Executive Engineer), Captain J. D. Herbert (Superintendent of the Mineralogical Survey), and the experts Wilkin and Reckendorf, who had actually visited and partially worked the mines. He urged that the sums

¹ J. A. S., Ben., XII., 434, and Stat. Kum., p. 267.

471; and Stat. Kum., p. 265.

² J. A. S., Ben., XIV., 471; and Stat. Kum., p. 265. ³ In a pamphlet published in London in 1845, and Stat. Kum., p. 262.

hitherto expended were barely sufficient to pay for the expenses of discovering and laying open the lode, and were utterly insufficient to carry on the experiment in the only way in which it could be made to pay. Nothing was undertaken until 1852, when the mines were again opened on the same footing, but the result was failure as complete as before. No attempt has since been made towards placing the copper mines of Garhwāl under European superintendence. In 1872, however, a European leased the mines and continued the extraction of the ore according to the native method, but was obliged to abandon the process, as he found the cost of the metal when manufactured more nearly approached the current rate for silver than that obtainable in the market for the best foreign copper. The copper mines in Kumaon have never been worked under European superintendence, and any remarks that I have to make on their value or fiscal history will be found under the notice of the mines themselves.

Connected with the history of mining enterprise in Kumaon special prominence must be given to the Kumaon Iron Works Company. Kumaon Iron Works Company still in existence, and whose origin is no doubt due to the continued belief, in spite of successive failures, in the possibility of turning the utilisation of the mineral resources of the province into a profitable investment. These had again in the regular cycle of inquiries become the subject of much speculation. From a review of the information before Government and the results of certain experiments made in 1856 the Directors of the East India Company sent out a Mr. Sowerby and a large staff of mining assistants in 1857 to carry on the smelting of iron on account of Government in the interior.¹ The fact was soon established that iron of an excellent quality could be manufactured at rates below the cost of iron imported from England, and a number of private individuals under the style of Davies and Co. were permitted to undertake operations for the same purpose in other parts of the lower hills.

¹ Report on the Government iron works in Kumaon, with plans, specifications, and estimates for establishing iron works in Kumaon, and remarks on the iron deposits of the Himalayas by W. Sowerby, C.E., printed as No. XXVI. of the Sel. Rec., Government of India (Public Works Department), Calcutta, 1859. These papers give a review of the English, Ulverston, Continental, Belgian and French, Rhineland, Black Forest, Bohemian and Styrian Iron Works, and estimates and plans for adapting the approved processes of those iron countries to the Kumaon mines.

The avowed object of the Government enterprise was to induce private companies to work by demonstrating the financial and physical possibility of carrying on iron works as a remunerative industry in this province. Messrs. Davies proposed to take over the tract between the Dhabka and the Bhakra, and their proposals were accepted, with an assurance that they might proceed in confidence to make their arrangements, as Government would grant the lease sought. They therefore took over the Khúrpa Tál works in the rains of 1858, and paid their cost price in 1863. This company also erected buildings at Káládhungi at a cost of Rs. 1,25,000. On the failure of the Government works at Dehchauri Colonel Drummond offered to take them over at a valuation. These works were given over to Drummond and Co., who paid the capital under agreement into the treasury in 1861. The forest rules were relaxed in favour of both companies, so as to allow them entire control over the fuel supplies, and eventually in November, 1862, both companies were amalgamated under the title of the North of India Kumaon Iron Works Company (Limited). Soon after the formation of the company instructions were received from the Secretary of State to construct a trainway to Khurja on the East Indian Railway, chiefly, it would appear, to afford an outlet for the iron manufactures of Kumaon. The trainway was to be laid with cast-iron rails manufactured at Dehchauri, and the company lost no time in making several thousand maunds of pig iron. Before the rails were made, however, the Government announced its determination not to undertake this line itself, but to hand its construction over to the Oudh and Rohilkhand Railway. It was essential to the success of the enterprise that some such outlet should exist, and the company accordingly determined to close its works for a time and await the opening of the line.

A license was granted by Government, but, unfortunately for the company, not executed till it was too late, and the deed of agreement contained briefly the following clauses:—1.—That a capital of 3½ lakhs of rupees should be paid before execution of the deed. 2.—That the company should pay Rs. 83,585 as the liquidated value of the works at Dehchauri, Ramgarh, and Khúrpa Tál, made over to them by Government, in four instalments, on or before the 1st September, 1862, 1863, 1864, and 1865, respectively; and in default of one

payment the whole to become immediately due. 3.—To pay yearly, for the first three years from 6th June, 1861, a rent of Rs. 1,500, and thereafter a yearly rent of Rs. 2,500 and a royalty of one rupee per ton of cast or wrought iron produced, and eight annas per ton of iron ore raised and sold without being smelted; the said royalties not to be paid unless they exceed, and only so far as they exceed, the rents of Rs. 1,500 and Rs. 2,500 before named; payments to be made on the 1st May yearly. 4.—To erect during the third, fourth, and fifth years suitable furnaces with requisite appliances for the smelting or blasting of iron (no number mentioned), and during these third, fourth, and fifth years to manufacture at least 750 tons of iron per annum on an average, and thereafter till the end of the term of 50 years manufacture 2,500 tons of iron per annum on an average of three years, to be struck in May each year. 5.—That the area of the forest at the close of every ten years should be covered with at least nine-tenths of such timber like trees as stood upon it at the commencement of the term, and when less than nine-tenths the company should plant to the necessary extent, failing which they should pay for each default Rs. 20,000. 6.—Not to transfer their lease without the consent of Government. 7.—To keep all roads (not being public ways made by Government) used by them in repair, also their works. 8.—At the close of their term remove their buildings, &c., first giving Government the option of purchasing them at a fair valuation. 9.—On the failure to pay or manufacture as stipulated, Government to enter upon and possess the works. 10.—But if such failure is not due to the neglect of the company, they shall be free from such penalty.

The purchase-money of the Khūrpa Tāl works was alone paid up, but the forfeiture clause was suspended by the local Government in 1868. The forest tract was never given over to the company in the meaning of the agreement, but its revenue has been separately collected and credited by Government. The license was not sent up from Calcutta till June, 1864, by which time the company had begun to see that the speculation would not turn out a profitable one. In fact the license deed was not prepared for signature until after the company had suspended operations, and then it was ruled by the Solicitor to Government that it ought not to be signed. The

map of the tract to be given over to the company was not completed till 1869. So much for the relations of the company to Government, and the delay in dealing with its affairs which has been shown throughout. In June, 1861, permission was obtained for the company's manager to draw against their capital. This was expended with little result, and in 1864, as above mentioned, the company was wound up. This result was in a great measure, no doubt, due to the company being unable to raise capital in the market, owing to the defect, or rather the want, of title, which appears to be ascribable to no fault of their own. Since 1865 correspondence has been carried on in reference to the affairs of the company and plans have been proposed for its resuscitation. In 1872 the works were visited by Mr. Jones of the Roorkee Workshops and valued; he made them then worth Rs. 1,26,733, with a debt to Government of about Rs. 80,000. There can be no doubt that the works can hardly be said to have had a fair trial, and the valuable opinion of the Commissioner of Kumaon may be quoted to the effect that there is every reason to believe that, if carefully supervised and fed with capital, the works should at least turn out as favourable under any circumstances as the East Indian Railway. There is no doubt that, in the distance, the fuel difficulty exists, but at Dehchauri and Káládhungi for many years this can scarcely be felt, and under penalties to replant, the company may fairly be allowed to have an unlimited supply from the neighbouring forest. At Ramgarh it is doubtful whether iron manufacture will pay, as the ore, though of the finest quality, lies at a considerable distance from any forests of any considerable magnitude, so that until it has been definitively settled whether coal does or does not exist in Kumaon the eventual absolute success of these Kumaon mines must remain problematical.¹ The increase of railways in Northern India and the development of the resources of this province must sooner or later press these difficulties into notice, and they will then obtain a final solution. "Too much has been written and too little done" hitherto in this direction.

In reviewing the causes of the poor returns from the different
 Causes of present poor returns. mines, one that presses itself into notice on
 the most cursory inquiry is the comparative

¹ For the materials for this note I am indebted to the office of the Commissioner of Kumaon and a note drawn up by Sir Henry Ramsay, than whom the Company and Kumaon has no more warm well-wisher.

inaccessibility of the principal mines. The copper mines of Sira and Gangoli, equally with Pokhri and Dhanpur, are situated on high cliffs in the interior. The talcoose and calcareous formations in which the ores are found occupy the high precipitous mountains which build up the outlying spurs of the principal range, and some lie within it. This chain itself is metalliferous, as the lead mines at Ghirti between Milam and Níti, the copper indications at Tola and elsewhere in the Juhár country, and the copper and iron mines at Polar near Rudrnáth combine to show. The absence of coal and the increasing cost of wood fuel, with the distance it has to be carried when the forests near the mines have been exhausted, materially enhances the cost of production, while the difficulties of carriage in the tracts where the mines lie are often such as almost to preclude the transport of ore for smelting, and the forests in the neighbourhood of most mines only suffice for the most moderate requirements. Another difficulty is the want of labourers. The present workmen only come to work in the mines from the latter end of October to the beginning of April, and many of the less productive mines have been abandoned owing to the miner class turning to agriculture and to supplying the labour market at Ránikhet and Naini Tál. This want, however, could be supplied from Nepál were regular wages and constant employment once established. Sea-borne copper, though inferior to native copper, is from its cheapness preferred, and until capital is invested in opening up the larger mines and conducting the whole operations on a sufficiently large scale to warrant the permanent investment of capital in machinery and proper furnaces, and other appliances for the more economical working of the ore, mining enterprise must remain as it is—a practical failure in this province. It may be said that these extended experiments have already been tried in the case of the Kumaon iron works, but this remark will scarcely apply, as that is another of those unsatisfactory operations which stopped just at the point where further progress would have decided the question for or against the possibility of making mining speculation a remunerative one in Kumaon. I shall now briefly describe the mode of working and the financial results of the settlements of the revenue from mines from the official reports and papers before describing each mine.

The mode of working the mines is the same in Garhwál and

Kumaon, and the suggestions for its improvement will serve for all classes of minerals. A

Mode of mining. A gallery or passage is cut in the face of the hill with such slight declivity outwards as is sufficient to carry off the water. These adits have more of the nature of burrows than that of the shafts known in European mining. The section is always small, and in those parts where the hardness of the rock occasions any difficulty in working the passage is scarcely sufficient to admit of a person in a creeping posture. In no place will it allow of an erect position. Where necessary, frames of timber formed of unsawn branches of trees, rudely and even carelessly constructed, are set up to support the roof and sides. Accidents are therefore not uncommon, and the frequently falling in of the mines is one result of these imperfect protections.¹

The ore as well as the rock is excavated by a very different
Mode of excavation. kind of pickaxe, the handle being made of a piece of wood with a knob at one end, into which a piece of hard iron is thrust and sharpened at the point. This with a miserable iron hammer, wedge, and crowbar, constitutes all the apparatus that the native miner has to depend upon. It is plain that with such tools no hard rocks can be penetrated nor can the softer ones be worked with much facility, and to this fact may be attributed the universal smallness of the passages throughout the mines, as the native miner can have his passage no larger than the rock which encloses the ore and its matrix will admit of. Proper pickaxes and steel gads (wedges) should therefore be substituted instead of the inefficient tools in use, and when blasting may be required the necessary materials should be provided. The miners work during the day, using torches made of dry pine, and clear out on an average from ten to twelve maunds of ore.

The ore is removed from the mine by boys, who pick up the
Removing the ore. stuff with their hands and put it into skins, which they drag along the floor by means of a rope and cross handle attached to their neck to the entrance of the mine. In most mines the greater part of this work must be done in a creeping posture, the string from the skin being fastened around the waist of the dragger. In place of this method wheel-barrows

¹ Paras. 19, 20, Captain Herbert's report, already quoted.

or sledges on four wheels and shovels should be used when the passages are enlarged and properly supported with sawn timber.

The ore or *dāua* being delivered at the mouth of the mine is reduced to a small size either by the water-

Crushing the ore.

mill or by the manual labour of women. A

large stone is placed on the ground on which they lay the ores; they then, either with a stone or a large hammer, and more frequently the former, proceed to pulverize the ore and pick out the impurities. In this way a woman may manage one to two maunds (82½ avd.) a day, according to the hardness of the ores. In Cornwall a woman will pulverize from 10 to 15 cwt. per day, according, as in the former case, to the nature of the ores. The method in practice there is, first, to dispense with the picking; secondly, to have the ores elevated, so as to enable the individual to stand while working, and to have a plate of iron about a foot square and two inches thick on which the ores are broken with a broad flat hammer. The impurities are then finally separated by a peculiar mode of dressing the ores with a sieve, by which a boy gets through with from one and a half to two tons per day. The ores are conveyed to the women, and from them to the boys by a man who attends for that purpose.

The washing of the ore in Kumaon also is performed by women,

Cleaning the ore.

who carry the stuff in baskets from the

entrance of the mine to a stream, where they

contrive by dabbling it with their hands to wash off the mud and finer particles of the earth. They then proceed to pick out all the pieces of ore they can get hold of; or, in the case of what may be submitted to the water in a commuted state, they work this against the stream, so as to gather it clean at the head of a small pit by handfuls; but, from the bad construction of the pits, it is with difficulty that this is performed. After picking up any larger pieces of ore which may have gone back with the stream, they scoop out the refuse with their hands, and then proceed with another charge. In Cornwall, one woman provided with a wheelbarrow and shovel for the conveying and washing of the ores, and a boy with a sieve for dressing them, as formerly mentioned, would accomplish a task equal to that of ten women on the system described.

The drainage of the mine is managed in a proper manner by an adit. But whenever any attempt is made to go below it, as is the case in most

Drainage.

if not all of the mines, the water is then raised in wooden buckets, handed from one man to another until they reach the adit into which they are emptied. In this manner six, ten, or even more men may be employed, whilst only an inferior number can be spared for excavating the ore. At the Sira mine, for instance, six men were found constantly engaged in lifting up the water, and there were only two at the ores : the work done by these six men could be effected with a hand-pump by one man ; but in order to keep the pump constantly going, two men might be required, and the remaining four added to the number of those who are excavating.

The furnace of the *Dhanauriya* or smelter is very simple, and is made of common stone and clay faced with

Smelting the ore.

slabs of quartzose schist, luted with a compost of chaff and clay. It is about $3\frac{1}{2}'$ long by $2\frac{1}{2}'$ broad, with an ash-pit about six inches square, all of which are built inside a house about $12'$ by $14'$, of which the roof is composed of planks. (Figs. A. B.) The operation of smelting takes about $28\frac{1}{2}$ hours, during



Fig. A. - Kumaur furnace!




Fig. B. - Vertical Section, from the front.

which time the fire is kept up, and after that the facing slabs and luting require renewal. The implements used are a crowbar, poker, shovel, and a pair of buffalo hides, dressed whole, to form the bellows, the neck of which forms the nozzle, and the buttock the valve for the ingress of air. The hides for making them are worth Rs. 12 apiece.¹ The furnace being freshly luted, the ash-pit is filled with charcoal dust and chaff, and a fire being lit, six baskets of iron ore, each containing about thirty sers (the ser = 2½ lb. 2oz. avd.), are placed round the fire. The blast is then commenced, one bellows being inflated while the other is undergoing depletion. In about half an hour the slag commences to flow from the floss-hole, which is kept open by a poker. In about two hours more, the ore having subsided considerably, two more baskets of ore and a corresponding supply of charcoal is given with a new luting for the bellows nozzle. In another two hours, this having also subsided, the charge is deemed ready. The fire is then raked out through the flosshole, and the charge, consisting of a pasty mass called *phalka* or *jhaui*, is shoved out with a crowbar by the smelter. The same operation is repeated until seven blooms are obtained, consuming thirty-eight baskets of ore, thirty-one of which are converted into the seven blooms, and the remainder, comprising the partially roasted ore, become the property of the smelter. The charcoal consumed weighs 340 sers, or a little more than the seven blooms, which weigh 327 sers, or about one-third of the ore expended (930 sers). Each bloom consists of three qualities of metal, all intermixed with earthy particles. These are kept separate, and are broken into small pieces before being sent to the *khatauniya* or refiner.

The furnace of the refiner is smaller than that of the smelter, and the implements required are a pincers, Refining the ore. poker, two or three sledge-hammers, an anvil, and bellows. The fire being lit, a mixture of one-sixth of first quality, one-sixth of second quality, and the remaining two-thirds of third quality, in all about six sers of bloom metal, is placed on the hearth opposite the bellows, with the larger pieces nearest the fire. The blast having commenced, in a quarter of an hour the slag begins to flow, and in another quarter of an hour the

¹ Beckett, III., Sel. Rec., N.-W.P., 25.

metal (now a porous, pasty mass) is taken out of the fire and subjected to the blows of two or more sledge-hammers; the blows being slight at first, to prevent the metal flying into pieces, but as it becomes more solid, they are given with the full force of the workmen. Meanwhile a fresh supply of bloom-metal is placed on the hearth, as at first. The hammered mass, after several hammerings, assumes the shape of a small bar, weighing one and a quarter ser; it is thick in the middle and tapering to either extremity, and six sers of charcoal have been used in its formation. This bar is now fit for the market, and is called by the workmen *phala*, but by the plains-people *pain*. The charcoal used by the refiner is made from the dry trunks of fir trees which have been felled for two or more years, while that made use of by the smelter is made from small green wood. The refiner class is subdivided into another, called Bhadeliya, who, instead of making the iron into bars, manufacture it at once into cooking utensils. Nine hundred and thirty sers of ore produce 327 sers of bloom-metal, which in its turn produce 82 sers of marketable bar-iron, or only 8·8 per cent. The bloom operation consumes 340 sers of charcoal and the refining process 667 sers, so that for every ser of iron produced 8·2 sers of charcoal are consumed. The Swedish furnace only consumes 1·83 times the weight of the iron produced.

The mines are leased for a term of years to contractors for a certain sum, and the lessee collects for the season from the different classes of workmen at the following rates: from each son or miner Rs. 2½; from each gang of smelters Rs. 4½; from refiners of the Khatauniya class Rs. 4½, and from those of the Bhadeliya class Rs. 6. The miner is originally sole proprietor of the ore, which he takes to the smelter to reduce into blooms, giving him for his trouble one basket of ore (30 sers) and one basket of charcoal (5 sers) for each bloom turned out; also for each set of seven blooms 16 sers of grain, and food for one man for four days; and at the end of the season a suit of clothes. Sometimes, however, owing to the smelter being largely in debt to the miner, he does not receive any charcoal from him. The smelter can only work for certain miners, generally five in number, not being allowed to work for any other miners; or, in other words, each party of five miners employ one family of smelters exclusively.

Each party of smelters must consist of at least five persons, but they generally count eight to ten persons. The share of each party of refiners is one half of the bloom-metal made over to them to refine, no further remuneration being allowed them. Refiners, unlike the smelters, are not bound to work for any particular person, but may work for any one that chooses to patronize them.

In the roasting and smelting of the ore Captain Herbert

Improvement needed in recommends a system of reverberatory furnaces for these two different processes. An appliances.

excellent material is at hand in the indurated talc known as potstone, which, though soft, is infusible. The simple blast furnace in use in Chili would also be an improvement. It is of a circular shape, similar to a lime-kiln, covered with a dome to confine and concentrate the heat. The ore is arranged in it in alternate layers with the fuel, which is wood, and being lighted it continues burning for a considerable time. When required, the heat is urged by a double pair of bellows worked by a crank turned by a water-mill.

Chili furnace.

The methods of reduction practised in England, where the subject is best understood, vary with the ore, and even with the establishment. But the English system.

differences are trifling and only affect the minor details. The two great objects to be effected are, first, by a proper calcining heat to drive off the volatile ingredients sulphur and arsenic, and to oxidate the iron, thereby promoting the fusibility of the ore and consequent separation of the metal from the scoria when in fusion; and, secondly, by an intense and properly continued fusing heat to effect the vitrification of all the impurities which thus form a slag at the top and are skimmed off while the metal sinks down in a comparatively pure state. To promote this vitrification of the ingredients occasional additions are made to the ore as the case may seem to require, though in general the run of the ores is such as to require little beyond a few slags of an old smelting. The operations of roasting and smelting are repeated several times, each smelting being followed by a roasting, to expedite which effect in the case of copper the ore is, after each smelting but the last, let into water to be granulated. This separation of the metal into such small parts assists the calcining power of the furnace, and the work is more speedily effected than if

performed on the mass. After the last smelting comes the process of refining or poling, which consists in keeping the copper in a melted state covered with charcoal, and introducing from time to time a wooden pole into the melted mass to produce the evolution of gaseous matters. Lead is sometimes used both in Hungary and England to expedite the previous operations of the refinery. The oxides of this metal are amongst the most powerful vitrifiers known. As such they are effectual in the assay and refinery of the precious metals, and as such they may be also used with copper. But the process requires attention, for if not stopped in time, or if too much lead be added, the copper itself will be oxidated and vitrified.

The process of manufacturing iron from the ores is different from that of copper, inasmuch as none but the oxides or carbonated oxides of the former metal are ever employed. In the copper ores, that is in those which occur in any quantity, the metal is combined with sulphur, which can only be driven off by repeated roastings. In the iron ores the metal is united to oxygen and mixed with various earthy impurities. In reducing these ores, then, there are three distinct points to be attended to: first, the provision of a substance which shall effectually take the oxygen from the ore, leaving the metal mixed only with its earthy constituents; second, the proportioning the flux used to those earthy ingredients so as to insure a complete vitrification of them and separation from the metallic particles; and third, a sufficient heat to fuse the latter, that the separation and reduction may be more complete. The first point is attained by using a sufficient quantity of charcoal in the reduction of the ores; the second by adding, as the ore may require it, limestone or other flux; and the third point is only to be effected by using a powerful blast furnace.

It is not easy to give the outturn from the mines, the arrangements are so intricate and the returns so imperfect. In 1868 about 29 maunds of copper were raised from the Kumaon mines, and in 1869 the same mines yielded the same amount, of which 21 maunds were exported. The Dhanpur mines in Garhwál yielded 10 maunds of copper in 1869, but every year since the produce has decreased. In 1868 the Kumaon iron mines yielded about 2,000 maunds of metal, and the Garhwál mines about 1,732 maunds, while the returns of 1869 give

5,153 maunds for Kumaon and 529 maunds for Garhwál. Besides this an immense quantity of copper is imported into Kumaon in the shape of manufactured vessels for culinary purposes: about 2,000 maunds of iron also are imported from the plains against 155 maunds exported. No reliance can be placed upon the estimates of outturn in recent years, as the mines have been leased for a term of years, and the lessees are not inclined to have their affairs too closely examined.

Previous to the Gorkhálí conquest of Garhwál the copper mines of Nágpur are said to have yielded Rs. 5,000 Gk. a year, or about Rs. 3,800 of our money. The entire mineral revenue of Kumaon and Garhwál, including mint dues¹ on the coinage of copper pice, had fallen in 1812 to Rs. 4,800 Gk., equivalent to Rs. 3,600 British currency.¹ This was mainly due to the neglect of the Gorkhálí Government, under which the mines had fallen in and become choked with rubbish. Their suspicious policy prevented them from trusting their own officers, whilst their want of probity precluded any private person from venturing to sink the capital necessary to re-open the mines. In 1815 the Nágpur mines were leased for Rs. 10, and in the following year for Rs. 15, and with the villages attached to them seldom brought in more than Rs. 1,850 a year, whilst those in Kumaon were leased at Rs. 850 a year. Up to the year 1826 the revenue of the Kumaon mines was included in the assessment of pargana Rámgarh, and that of the Garhwál mines in pargana Dhanpur, and subsequently was accounted for in the returns of the pargana within which they are actually situated. Between the years 1815 and 1840 the revenue derived from mines averaged as follows:—

	Kumaon.	Garhwál.	Total.
	Rs.	Rs.	Rs.
Copper ...	801	2,086	2,887
Iron ...	1,905	226	2,131
	<u>2,706</u>	<u>2,312</u>	<u>5,018</u>

The highest mineral revenue of the province for any one year amounted to Rs. 5,417. This return was not altogether due to the

¹ These mint dues were collected for a few years under British rule at the mines of Dhanpur and Gangoli, and at one-half per cent. yielded a revenue of Rs. 300 a year. To Board, dated 6th August, 1831.

smelting of ore, and included the land revenue of villages attached to the mines for the location and support of labourers. Mr. Beckett in his report¹ on the settlement of Garhwál gives the revenue of each mine from 1839-40 to 1863-64. The revenue every fifth year from each class of mine during this period was as follows:—

Class of mine.	1839-40.	1844-45.	1849-50.	1854-55.	1859-60.	1863-64.	Total revenue from 1839-39 to 1863-64.
	Ra.	Ra.	Ra.	Ra.	Ra.	Ra.	Ra.
Iron ...	609	727	779	480	272	131	3,764
Copper ...	1,990	2,442	2,138	1,305	81	627	21,304
Lead	5	3	3	...	10	64
Total ...	2,599	3,174	2,920	1,788	353	771	25,132
N u m b e r of mines worked,	24	25	23	17	11	30	76

In 1865 there were 24 iron, 9 copper, and 2 lead mines worked in Garhwál, and 33 iron, 35 copper, and 3 lead mines had been abandoned. The lead mines have since been abandoned, and the revenue from copper and iron mines in 1878-79 was as follows:—

	Copper.	Iron.	Total.
	Ra. s. p.	Ra. s. p.	Ra. s. p.
1878 86 8 0	136 12 0	223 4 0
1879 89 0 0	153 8 0	242 8 0

There are no statistics of outturn for these years.

In his Kumaon settlement report Mr. Beckett gives the revenue of each mine from the year 1844-45 to the year 1872-73. The revenue every fifth year from each class of mine during this period was as follows:—

Class of mine.	1848-49.	1853-54.	1858-59.	1863-64.	1868-69.	1872-73.	Total revenue from 1848-49 to 1872-73.
	Ra.	Ra.	Ra.	Ra.	Ra.	Ra.	Ra.
Copper ..	100	48	Nil.	120	67	30	2,831
Iron ..	2,274	1,751	1,532	870	929	1,420	46,126
Total	2,374	1,799	1,532	990	996	1,450	48,957

¹ Allahabad, 1866.

The following quotation from the Commissioner's report in 1874 gives the opinion of those best acquainted with the subject on the future of the mining industry in Kumaon:—"Iron and copper abound, but at the present value of labour the mines are worth very little. The sons or miners have, as a rule, given up their old trade and taken to contracts. The great attraction to miners in former times was the cheapness of grain in the Khetari valley, where iron was most extensively manufactured. This advantage no longer exists, for the market at Ránikhet has doubled the price of grain, and the miners would be no longer content to exchange their labour for the small profits on iron. Copper mines are in no greater favour. Formerly some villages where the miners reside were included in the mining leases of Kumaon and Garhwál. These villages have been settled with the miners, therefore they are no longer servants of the contractor. Tea gardens and other labour markets offer much better terms than a contractor, who, at the least possible expenditure, tries to make the greatest possible profit. These contractors know nothing about the science of mining, and they have no money to expend in penetrating beyond the worked-out galleries. In fact the mines have collapsed, and without considerable outlay no reasonable profit can be expected. Labour is expensive, and English copper can be bought at a cheaper rate in the Almora bazar than the local miners can produce it with profit. I expect nothing more from native petty contractors than a pittance which they can realize by the resident miners working when convenient to themselves, when they give half of the ore to the contractors and keep the other half. No doubt there is abundance of copper in Gangoli; but any mines, copper or iron, that are now worked barely produce sufficient for local consumption. Agricultural instruments are made for the people of the surrounding country, and a few copper vessels; but all the mines in the interior are in remote places, and too far removed from a good market to be of much value."

The gold exported from Kumaon is either obtained from the
 Gold. streams within the province, or is brought
 down by the Bhotiya traders from Tibet.
 Although no mine of this metal has been discovered in the province,
 there are indications of its existence in Garhwál. The sands of the

Alaknanda, Pindar, and Sona furnish a small amount of gold-dust. The Ganges also is auriferous as far as Lachhman Jhūla, and the Rámanga for a short distance below its junction with the Sona. The washing is nowhere a profitable occupation, and scarcely gives an average of four annas a day for each workman. The gold obtained by washing the sands of rivers paid a small duty during the Gorkhāli rule, and was leased with the forest duties for a short time after the British occupation, but the amount was too trifling to render its continuance expedient, and it was accordingly abolished by Mr. Traill. Undoubtedly a greater return might be had from this source by the use of mercury, as in Australia, for the purpose of separating the gold from the sand; for as this is afterwards recovered by a simple process of distillation, the expense would be very little more than it now is. Captain Herbert found gold in a matrix of granite near the Alaknanda.

The gold imported from Tibet by traders is chiefly taken in exchange for grain or cloth to balance their accounts, as rupees are taken by them at the hill fairs for the same purpose. The principal gold mines in Tibet, *sār-chaka*, are ten days' journey beyond the borax fields further north and north-east in a district otherwise uninhabitable, named *Sār-bachyád*. These are farmed or managed by a *sār-pan* or gold commissioner on a triennial contract direct from Lhasa. The lessee in 1845 was also Garpan Urku-wa at Gartoh, and paid Rs. 17,000 per annum for the lease. He had 170 miners at work, for whose subsistence he used to send supplies from Pruang. It would also appear to be sometimes the custom to sublet 'claims' at a tax of a *sārjao* or *jao* of gold, about $7\frac{1}{2}$ mashas, or ten rupees.¹ The gold mines are worked by pits and shafts under ground, where the gold is found in its pure native state, and undergoes no other process than washing and shifting, and after that requires little or no refining. In this state, tied up in little bags called *sār-shu* (H. *phatang*), weighing about 90 grains, it forms the heavy currency of the country. A superstitious belief holds ground that no large nugget should be removed, as it belongs to the genii of the place, but the Lāma of Gnari is said to have one weighing nearly a ser.

¹ H. Strachey, J. A. S. Ben., XVIII. (2), 5; J. D. Cunningham, *ibid*, XIII; Herbert to Commissioner of Kumaon, 19th January, 1826.

Gold is sold at the same fairs as the borax, and is imported to the value of about Rs. 10 to 12,000 annually.

The gold in the bags commonly current has usually not more than 7·73 specific gravity. Even the picked

Thok Jalang mines.

yellow grains have only a specific gravity of 11·96, showing that they are alloyed with some other metal. The grosser impurities appear to consist of iron more or less oxidised. One of Montgomery's pandits visited the gold mines of Thok Jalang in Rohtoh in 1867,¹ and describes it as a great excavation from 10 to 200 paces in width and 25 feet in depth, access to the bottom being by means of steps and slopes, the earth dug out being thrown on either side. The digging is carried on with a long-handled kind of spade or an iron hoe, the iron for which comes from Ladák. A very small stream runs through the gold field, and the bottom is consequently a quagmire during the daytime. The diggers dam up the water, leaving a sloping channel for an escape. A cloth is then spread at the bottom of this channel, and the channel is supplied by one man with the auriferous earth, and another gives water, so that the gold sinks to the bottom and is caught in the cloth at the end. Some nuggets weigh up to two pounds. The diggers come from the Tsang province round Shigatze or Digarcha. There are numbers of abandoned gold fields in different directions about Thok Jalang, and probably a whole series of them from Rohtoh to Lhasa. The Sárpan levies a tax of about half a tola (*saishu*), or two-fifths of an ounce, from each digger. There is no wood, and water can only be had from melted ice. A cold wind blows at all seasons, and, in consequence, the tents of the diggers are pitched in excavations in the ground to protect them from this wind. The dried dung of yaks, ponies, and sheep afford fuel. The Tibetans cook and eat three times a day, their food consisting chiefly of boiled meat, barley cakes, buttermilk, and tea stewed with butter : they also smoke a great deal. They always sleep with their knees close up to the head and rest on the knees and elbows, huddling all the clothing on to their backs. The price of gold at Thok Jalang was about 5½ to 6 rupees per *saishu*, or 30 rupees per ounce.

¹ Rec. G. L. H. D., 36.

Silver was brought down to these provinces from Tibet in former times. It was imported into that country from those surrounding it (probably China), and does not seem to be found in Tibet itself in any quantity. It was sent into Tibet in a crude state in lumps called *doja* or *thaka*, of a general value of Rs. 165 each. Importations from that source have ceased for some time, owing possibly to the great and growing influx of silver in the shape of rupees from British territory. Formerly all borax, salt, &c., was bartered for grain, cloth, &c., but now, while a large amount is still disposed of in that way (probably to procure actual necessities), still, whether it proceed from the increase of trade and the portability of coin for hoarding purposes, or from the existence of a greater demand for silver in Tibet, by far the largest amount of borax is disposed of here for British money. The Bhotiyas, too, state that our coin is largely current in Gartok and the other large towns, and is preferred by the inhabitants there to the coinage of other countries. They ask for the *Cheharáddr Rupaya*, or face-printed money.¹ The difference in the exchange now made up in Government rupees cannot be less than eighty thousand to one lakh of rupees per annum. This trade in rupees dates from about 1820, when they began to displace the Srinagari and Ladáki rupees.

The mines of copper in Kumaon and Garhwál have never been of much practical value either as a source of supply for local consumption or as offering a valuable return to labour and capital. They are still, however, deserving of notice, and we shall now describe each in succession, commencing with the Gangoli mines in Kumaon.

The Rái mine in pargana Gangoli is the most important in Kumaon. The ore is chiefly pyrites in a matrix of indurated and sometimes slaty talcose and steatitic schists inclosed in dolomite. In some places the one; and in some places the other, forms the roof and sides of the mine. The dolomite has a large crystalline grain and great tenacity, and forms a perfectly durable work when excavated. The schists when massive may be depended on, and can be easily worked,

¹ Lawder in Rec. Geol. Sur.

but, as a rule, they occur of such inferior consistence, having much the appearance of re-united débris, that they require support, and often occasion much inconvenience and even danger. The ore occurs in the schists in numerous strings, having every appearance of being leaders, as they are called, to solid ore, and forming a distinct lode. The strike or direction of the strata is nearly W.-N.-W. to E.-S.-E., dipping at an angle of about 45° to the N.-N.-E. The copper ore is accompanied by iron pyrites which are occasionally found in the pentagonal dodecahedron form, but most commonly in such irregular and anomalous forms as can with difficulty be described. There are a few specimens of grey copper, but the working ore is undoubtedly pyrites. On visiting the mine in 1836, Captain Drummond found the lode about two feet wide, containing a good yellow copper ore, but with a large proportion of its matrix talcose, twenty per cent. only being metalliferous. The ore is extracted in the usual way by means of drifts slightly inclining upwards, to allow for drainage. The adit at Captain Drummond's visit was driven on the course of one of the lodes which continues west about 60 feet, when it falls in with another lode that alters its direction to 15° and afterwards to 30° north, inclining nearly 50° to the east of north. At that time the adit had penetrated some 348 feet from the entrance. The ore had been taken away from beneath as far as the miners could excavate it, and the hollow had been filled up with rubbish. From above, too, the ore was taken away as far as it was found productive. The passage varied from two to four feet in height and from two to two and a half feet in width, being bounded by the hard dolomitic rock which the miners did not know how to remove. In 1868-69 these mines fell in and became flooded with water. About a couple of hundred yards to the north, and in the same hill, is another similar deposit of copper. This used to be laid open to the surface during the rainy season, and was then allowed to fall in, so soon as the water employed by the miners to carry off the talcose mud from the ore ceased to be plentiful. This also has ceased to be worked for some time.

In 1815 one specimen of fused copper from the Gangoli mine and several specimens of the ore in matrix were sent to the Mint at Calcutta for assay. The report showed that the ore was mixed with arsenic and sulphur,

Ore assayed.

and produced 25 per cent. of malleable metal, but the specimens were too small to allow of any exhaustive examination of them.¹ In 1826 Captain Herbert valued the outturn at 35 per cent. of the pure ore, and in 1836 Captain Drummond gave the general result from the pyrites in their perfectly pure state as about 30 per cent. of metallic copper.² Pyrites, though not a rich ore, is the most important of any, from its abundance and from being generally more to be depended on for continuance than the richer varieties. In England more copper is obtained from it than from all the other ores together. The Gangoli, Sira, and Sor mines were farmed from the conquest until 1828, when they were leased for one year to the miners,³ and were again farmed at a reduced rent in 1833. In 1815 they yielded a revenue of Rs. 850, increased to Rs. 1,201 in 1819 and 1820, and to Rs. 1,215 in 1821 and 1822, but in 1874 the whole of the copper mines of Kumaon brought in a revenue of only Rs. 30 a year.

The Sira mines in Patti Bárabási in pargana Sira are situated on the northern side of a hill somewhat higher than the one at Rái. The ore here too consists of copper pyrites, accompanied by iron pyrites in a gangue formed of dolomitic and talcose rocks. In 1816, a specimen of copper ore from the Sira mine was sent to the Mint at Calcutta for assay, with the result that it was found to contain only 24 per cent. of malleable metal, so that it was thought that this mine would not repay the working.⁴ Captain Drummond found that nearly thirty-three fathoms from the entrance the adit struck on a copper lode on which a level passage was driven that continued westward, its course being about 10° south of west, and the dip northerly from 45° to 50°. The ore was harder and more mixed with iron pyrites than the ore at Rái. At the end of the level a second lode yielding a poor ore was met, and beyond it a pit was sunk which seemed to have yielded in former times fair returns.

The Gaul mine in Patti Kharáhi and the Sor Garang are similarly situated, but the ore produced is in very small quantities, consisting of grey

¹ To Government, dated 14th June, 1815. From Government, dated 18th January, 1816. ² See further Stat. Kum., p. 359 : J. A. S. Ben., VII., 935. ³ To Board, dated 2nd January, 1829. ⁴ From Government, dated 18th January, 1816.

copper, copper pyrites, and carbonate of copper. Stentite and limestone are the neighbouring rocks, the stentite forming the gangue. The Sor mine had not been worked up to 1833, although a lease at a small rental had been taken out in 1821.¹ Captain Herbert notices that all these ores are free from the presence of arsenic, which, above all other metals, deteriorates the quality of the copper and is most difficult to remove.

In digging the foundation of a house at Hawallāgh the workmen came on a vessel containing three crystallized Bournonite. specimens of bournonite, the triple sulphuret of copper, antimony, and lead, and the only trace hitherto discovered of its existence in these hills (1826). Copper pyrites also exist near Gannai and Phudialī in Patti Athghion in pargana Gangoli; at Bujāl and Rathūyat in Patti Bel in a matrix of stentite and foldspar; and at Tāmbarūn in Patti Gangoli in a matrix of talcose rocks. There are small mines in Patti Giwār, at Chīn ka Kālī, Belor, Sor, and at Kenuakhet, on the east bank of the Ladhiya river in Kālī Kumaon.

The copper mines of British Garhwāl are more extensive and have always borne a higher reputation than Copper mines of Garhwāl. those of Kumaon. The principal are situated at Dhanpur and Dhubri in pargana Dewalgark. These mines yielded a considerable outturn in former times, but of late years operations have not been so vigorously carried on, owing to the intricacy of the workings, and the idea prevailing among the miners that very little ore remains in the mines.

The Dhanpur mine is situated on the north side of a high and precipitous range in compact dolomite.² The Dhanpur mine. ores are principally copper pyrites and grey or vitreous copper ores, with the red oxide and green carbonate in smaller quantities, the latter being scarce. The ores are found in a bed about fifty to sixty feet wide, which runs nearly north and south, and underlies east about one foot in the fathom. It is divided by a bed of potstone or indurated talc, which runs through the copper formation longitudinally, conforming to the strata and having a frith

¹ To Board, dated 14th February. 1829.

² Mr. Wilkin's report, J. A. S., Pen., XII., 454; Beckendorf's report. *ibid*, XIV., 471; Captain Herbert to Commissioner, 10th January. 1826.

or flukan on the western side. The seams of ore are sometimes one foot in thickness, but seldom more than one inch. In his report on Dhanpur, in 1816, Mr. Traill declared that these mines were incapable of much improvement, and that the ore produced was not rich. The lease of the mines, including the twenty-two villages attached to supply the requisite labour and grow grain for the miners, was fixed at Rs. 1,850 a year. Up to 1829 there was little improvement, as the lessees were too poor to undertake the cleaning out of the mines, and no capitalist would venture to take them.¹ In 1838, the best seam or vein seen by Mr. Wilkin was not more than half an inch thick, and in 1841 the best lode worked was about two inches. The veins are very close together, and being softer than the matrix, the ore used to be first removed, and then the miners burned the rock with wood and threw water on it to facilitate removal. The red dolomite is of such a consistence as to seldom require props for its support, thus enabling the miners to dispense with wooden framework and to work all the year round, while the situation of the mine on the top of a hill admits of adits for drainage. The interior consists of a network of chambers from end to end.

The Dhobri mine is situated on the south side of the Dhanpur range in very nearly the same kind of rock as the Dhanpur mine, but in this mine most of the veins are horizontal, running along the side of the hill. At the surface they are very small, containing oxide of iron and green stains of copper, and occasionally copper pyrites. The present working mine is not extended very far from the outside of the precipice or surface, the ores being much the same near the surface as at a distance from it. When the miners find their passages growing long and tedious they begin outside on a new vein. There are several old mines west of the Dhobri village ; one of them is very extensive, and the ores seem to have been most abundant when the horizontal vein was crossed by perpendicular ores ; but the whole of the horizontal vein has been taken away. The most western of these veins is said to have been very rich, but it fell in about the time the Gorkhâlis entered the province, and has not been opened since. The ores of these mines are principally copper pyrites, containing 25 per cent. of copper. There is water for machinery about a mile and a

¹ To Government, dated 14th February, 1829.

half below the mine, and wood for all purposes near that place. There is another mine on this range at Maulgiri, said to be in the same rock as the Dhobri mine. There are other mines of both copper and iron in Dhanpur, but few of them are worked, and they are for the most part of little value. The mines were leased in 1872 to a European, but even then western intelligence and energy could not make them a remunerative investment. There is a copper mine in Patti Lobha at Agar Sera¹ in the face of a precipice on the right bank of the Rámanga river which was leased in 1872 for three rupees. The lessee, however, makes little profit, as the shafts have been sunk so deep that men are afraid to enter them, and the rock is too hard to allow of fresh shafts being driven, unless at great expense.

The Pokhri copper mines early attracted the attention of the Government of the country. For many years they had been worked by the Garhwál Rájas, and subsequently by the Gorkhális. They consist of several separate mines; that known as the Chaumattiya is situated in talc which rests on dolomitic limestone. The Chaumattiya mine. lode after crossing the ridge east of the mine enters a very compact basin, in which is situated the Duined mine. This has not been worked much, owing to the softness of the talc and the abundance of water, but it is said to have a good lode in one part of it. The lode then crosses the hill near Deothán, a small village above the mine, and is found near Gúgli and Keswara, where some ores have been extracted, but have never proved very profitable in working.²

The Rája's mine is situated about 450 yards north of the Chaumattiya mine in common dolomite which rests on talcose schist. A shaft of 70 fathoms was dug by the early workers meeting an adit which must have been driven over 100 fathoms through dead ground. Several other adits were driven, and when they fell together, about one hundred years ago, there were three places where copper was found—the Gaja Chauk, Kuvera Chauk, and Bhartwál Kda, all of which have now fallen in. The produce was about 300 sers of ore a day, yielding 25 per cent. of copper. Two-thirds were claimed

¹ Described by Beckett in *Sel. Rec., N.-W. P., III., N. S., 371.* ² For details of these mines see Mr. Wilkin's report in *J. A. S. Ben., XII., 484*, and Beckett's *ibid.*, XIV., 471.

by the Rájá, and the remainder was left to the miners, who had also grants of land free of revenue. The experiments of Mr. Wilkin already noticed were confined to these two mines and a new mine which he opened close by. In the new mine the lode was very promising and yielded good specimens of ores near the surface, but at a depth of fifteen fathoms it became poor, and was consequently abandoned. During the time the experiment lasted the expenditure on the Chaumattiya mine amounted to Rs. 2,847, and the return in copper to Rs. 231, besides about three to four hundred rupees worth of ore. Rs. 347 were expended on the Rájá's mine, and the experiment was then abandoned, and the new mine cost Rs. 246 before the operations were closed. The entire net cost of the undertaking when operations ceased was for labour, in working the mines, less sale proceeds of copper, Rs. 2,585, and for European superintendence Rs. 4,800, or a total of Rs. 7,385. These mines were then leased to a native contractor for Rs. 500 a year.

In addition to the three mines mentioned above there are several others in the vicinity of Pokhri, some of which were worked by the former rulers, and some again have never been opened. Mr. Wilkin noticed the principal mines, and described them as follows:—

The Nola mine is situated about two and a half miles north-west of the Pokhri mines in talc, which rests on dolomitic limestone. The lode is a bed of yellow or buff-coloured talc, about four feet wide, dipping north-west at 50°; it rests immediately on the dolomite limestone, and has a sulphuric effervescence on the surface. This mine is said to have been rich; it is situated on the western side of an extensive basin or valley, on the eastern side of which ores have been turned up by the plough, but no mine has been worked. This is an extensive field for mining, as the lode may be productive throughout the basin or valley. There is wood and water for all purposes near this mine.

The Thála mine is situated about a mile north-west of the Nola mine, probably on the same lode, in an extensive plain or comparatively level surface. It was first worked in 1810, and again in 1825, but there being no good facility for adits, the water prevented its being worked to any considerable depth. The miners who worked it state the ores to be

copper pyrites disseminated in a lode of two feet wide, one-fifth of which was metalliferous. There is plenty of wood for all purposes in the neighbourhood of this mine.

The Danda mine is situated on the hill, about 500 yards above the Thála mine, in chlorite slate and talc, which on the north-western side comes in contact with common dolomite. This mine has been worked to a considerable extent, and is said to have yielded Rs. 52,000 profit in one year. The ores are of good quality, and found in three or four different beds or holes which dip into the hill at an angle of 30° . The chlorite slate in which the beds of talc and ores are found is so hard as to stand without timber ; it also contains finely disseminated copper in small quantity. The lodes run into a fine fall or basin westward, in which Mr. Wilkin thought they would be found productive. There is abundance of wood near this mine, but no water for machinery nearer than the Thála mine.

The Tálapungla mine is situated about a mile north-east of the Danda mine in talc, which rests on dolomitic limestone. The strata in which the ores are found is about six fathoms wide, dipping south-west at various angles. The bed is extensive, but the ores are scarce ; however, this might improve at a distance from the surface. Ores have been found in a precipice, east of this mine, near the village of Bangtál, but at present the outcrop is covered with rubbish ; it is in the talcose formation, and has good facilities for working.

The Kharna mine is situated in the ravine below Bangtál, near its junction with the Nágál river in talc ; it was discovered by the water of the ravine washing away the strata, and leaving a quantity of ores exposed to view ; these ores were taken away by the Pokhri miners and the mine was worked five or six fathoms under the surface, beyond which they were prevented from going by the water. They say that the lode at the bottom of the mine for two fathoms in length is one foot wide, of solid copper pyrites. Of late years nothing has been done at this mine beyond washing among the surface layer, which contains a small quantity of copper pyrites. There is plenty of wood in the neighbourhood of this mine and water for machinery, but no facility for adits.

Mr. Beckett has described the mode of mining and preparing the ore of the Agar Sera¹ mine from which the following account has been extracted. The gangue consists of white and yellow quartz much encrusted with green carbonate of copper, and is so difficult to work that not more than 40 to 60lb. of ore can be excavated by one man in a day. The workings are dry and the lode has a dip of about 30° below the horizon with a north-westerly direction. The ore is pounded and moistened with water and receives an admixture of five parts in six of limestone as a flux. The charge, consisting of about 6lb. of unmixed ore, takes about half an hour to melt and is placed from time to time in handfuls on the furnace, and covered with oak charcoal which is occasionally moistened with water. When the fire falls in after the last supply of ore, the charge is ready to be taken out. The door of the furnace is then taken away and the remains of the fire being raked out, there appears at the bottom a melted mass which, after being stirred about a minute or two to allow the heavier particles to settle down, is sprinkled with water to harden the surface. Three or four of these charges being taken away, the melted copper is found at the bottom in a small mass weighing about 2½ ounces, for which twelve pounds of charcoal have been used. Thus from every 100 parts of ore about 2½ths parts of copper are procured, having consumed 200 parts of charcoal or, in other words, 137lb. of charcoal are required for the production of a little over two pounds of pure copper, which sells at about one rupee a pound.

The iron ores found in Kumaon all belong to either of two varieties, the rhombohedral or the prismatic. The
 Iron mines.

first is a peroxide of iron containing in its best defined type 70 per cent. of iron and 30 per cent. of oxygen. The workable ore, however, often contains earthy impurities which reduce the proportion so low as 50 per cent. of metal. This is the common species. A variety of this, known as red hæmatite, also occurs in many places, and frequently contains small grains of specular iron ore of a highly splendid lustre. At Rāingarh it passes into the variety known as scaly iron ore, consisting of loosely cohering glimmering particles of a steel-grey or iron-black colour, strongly soiling and feeling unctuous to the touch. Captain Herbert

¹ See Sel. Rec., N.-W. P., III., N.S. p. 34. and Glenn. in Science, I., 230; As. Res. XVIII, (1), 227 (Herbert); Traill and Batten, Statistics of Kumaon, Agra, 851.

considers these beds connected with those at Dhaniya Kot on the Kosi. Both yield very good iron. The prismatic species or hydrated peroxide is only known to occur in the Chaugarkha pargana.

The following are the principal iron mines in Kumaon. In Patti

Kumaon mines. **Agar of the Rámgarh pargana,** as noted, the iron is of the species known as scaly, some-

what laminated in structure, slightly micaceous, and influencing the magnetic needle. The names of the principal mines are the Lú-gani, Nathúa Kán, Gulla and Satbúnga mines, the last of which has a rich hæmatite. In Patti Rámgarh, also, there are several mines that are largely worked. For the first 18 years of British rule these mines were leased to the headman of the Agarís at a nominal rent,¹ which up to 1826 included all the iron mines in Kumaon. In 1833 the mining industry in Rámgarh received its first check in the emigration of the miners to Khetsári in Páli, and has never since recovered its early importance. In the Chaugarkha pargana, the ore of the Muniya mine in Patti Lakhanpur is of the prismatic species. It is of two varieties, the ochry and compact. The former sometimes contains octahedral crystals and magnetic iron ore, and in the neighbourhood of the mine on the summit of a small hill there occur rolled pieces consisting of grains of quartz and small octahedral crystals of this mineral cemented together. These pieces are magnets, and have each two poles. The ores, too, contain manganese in notable proportion, and would consequently afford a very good steel; as it is to the alloy of this metal that the superiority of the steel manufactured from brown ore is attributed. In Patti Dárun there are mines at Digarhia and Jhiratoli of the same nature; in Rangor Patti at Jalal and Digarhia; in Patti Kharáhi at Lob. A specimen of iron ore brought from the neighbourhood of Milam, called by the natives of Malla Juhár '*buldúnga*,' seems a crystalline variety of red hæmatite. It is used there for a red dye, the colour being extracted by rubbing the stone on a hard surface while wet.

The mines of Patti Giwar in pargana Páli are found at Chiteli,

Giwar mines. Sirauli, Khetsári, Simalkhet, Gudi, Bailgaon, Bonigarh, and close by at Mehachauri and

Tilwára. The valley in which the iron is produced runs nearly north and south, and extends from Dwárahát on the south to Pandúa

¹ To Government, 24th December, 1833.

Khal on the north. It is formed by the rivers Kotlar and Khetsari, both flowing into the Rám-ganga, the bed of the former being about nine miles long, and that of the latter six miles. The ores lie on the east side of the valley and occur along a range of hills about thirteen miles in extent. The Simalkhet mine¹ is the largest, and has four entrances penetrating upwards of 350 feet into the mountain. The ores here consist of red hæmatite of a good quality in a gangue of clay slate branching in every direction and at all angles. The lodes range from 3' to 18' in width and $2\frac{1}{2}$ ' to 15' in height, the average being from 3' to 5' by 4' to 6'. There is no water, and the surrounding rock is compact, requiring few supports.

In Sayalgarh of pargana Kotauli there is some iron ore not at present worked. At Manglalekh in Talla Rao the ore is much esteemed for its quality, and is raised in some quantity. At Dehchauri, Rámgarh, and Khurpa Tál the mines are in the hands of the Kumaon Iron Works Company. The Lugthan mine is in Malla Katyúr, and there is another in Baraon Patti in pargana Gangoli. The burrows at Khairna are now unworked, likewise those at Simalkha and Uchakot.

In the Garhwál district the iron mines in Patti Painkhanda exhibit specimens of granular iron pyrites imbedded in veins of quartz which occur in a dark-greyish talcose schist. They are apparently not very rich in ore. In Patti Sili Chandpur the Rajbúnga mine gives a rich hæmatite which is slightly attracted by the needle, and is still worked. The Khúsh mine in the same patti gives a micaceous ore, scaling off easily and showing minute crystals resembling garnets on the edges of some specimens. The adjacent beds seem to be chloritic schists; this ore affects the needle. In Talli Chandpur magnetic ore is found with hæmatite, and a specimen from Patti Talli Kálíphát resembles specular iron ore.

Specimens from the Búkhandá mine in Patti Bichhla Nágpur are also of a micaceous nature, and seem to contain in parts minute crystals of quartz and pyrites, otherwise they much resemble graphite, and soil the fingers when touched. They do not influence the compass needle. The Jákhtoli mine in the same patti give an ore which

¹ III., *Sel. Rec.*, N.-W. P., 23.

is probably a clay ironstone. It is of a light coffee colour and of little specific gravity. The Gilet mine close by gives an ore of a similar quality. In Malla Nágpur the ore is probably hæmatite. A vein of iron pyrites runs along the Alaknanda in this patti near the village of Hath. The people call them '*sona ke pathar*,' or gold stone, and sell them to the pilgrims to Badrináth at high rates. The stones, in the form of powder, are used as an orpiment, and the stone itself as a flint. A specimen from Nágpur itself is probably a carbonate of iron.

The Mok mine in Patti Malli Dasauli yields an ore of which Dasauli and Bachhansyún. specimens appear to be magnetic, rich in metal, black in colour and crystalline, and laminated in structure. It possesses highly magnetic properties. The Charbang mine in the same locality is of a similar character, very rich in iron, and, according to Mr. Lawder, exhibits its polarity in the direction of the planes of lamination. The Dúngara mine in Patti Bachhansyún gives specimens which may possibly be an earthy hydrated oxide of iron. Its colour varies from ochry to dusky black, streak the same. It is of little specific gravity, the clay seemingly predominating. The iron of Bachhansyún, however, has a wide reputation for hardness and toughness. Sledge-hammers (*gan*) made from it have been found to last out those of the best English metal.

The ore of Pipali mine in Patti Iriya Kot is probably a Iriya Kot. hydrous form of sesqui-oxide of iron, the clay largely predominating. The Danda Toli mine in the same patti seems to give an argillaceous variety of brown hæmatite. Lohba affords a rich hæmatite, raised in large quantities. The Chalya mine in Patti Painún gives a hard and brittle ore possessing the iron-black colour and metallic lustre of magnetic iron, but specimens of it failed to affect the compass needle in the manner characteristic of that ore. It may possibly on analysis be found to contain manganese.

The deposits of lead are fairly numerous. The ore is found at Táchhíra in pargana Dhanpur mixed with Lead. a little silver. A large mine also existed at Ghirti in the snowy range between Milam and Níti, but this has lately been closed by a landslip. The mines at Ralum and Banskum on the banks of the Gori river and at Baidli Baghur

are unworked. The ore is galena, and the matrix is siliceous, with varying proportions of feldspar and calcspar. The Nágpur mine is a fair one, but somewhat inaccessible. Near the Gaul mine in Patti Kharáhi and at Sor Gurang copper mines there are deposits of lead ore. The former is a galena traversing a siliceous limestone, but neither are regularly worked. When the villagers require lead they burn the rock, and the lead, more or less sulphuretted, trickles from the crevices. A large nodule of lead, the size of an eighteen-pound shot, was found in Patti Maundarsyún, on the banks of the Nayár river. It consisted of pure galena, but though search was made no more could be discovered. In Jaunsár there are mines at Maiyár and Borela on the left bank of the Tons river and at Aiyár on the right bank. At Maiyár and Aiyár the matrix is clay slate, and at Borela, limestone supposed to be a bed in the clay slate.¹ The revenue from these mines is now nominal.

Yellow arsenic, known as *haritál*, is found in the northern parts of Kumaon, near Múnsyári. A small portion is brought down every year by the Bhotiyas for sale at the Bageswar fair in January.

Indications of lignite appear near Ránibágh close to Haldwáni, the Barakheri pass near Bhāmuri, and in the streams of the sub-Himálaya near Najibabad in the Bijnor district. They do not give promise of any workable fuel, and, judging from the experience obtained in other parts of the hills, it is questionable whether any lignite deposits will ever be discovered of such extent and quality as to repay the expense of mining them. An analysis of a specimen of the Ránibágh lignite gave carbon 60·0, volatile matter 36·4, ash 3·6. The percentage of ash, however, contrasts favourably with Bengal coal.² Traces of a true peat are found at Bhím Tál. In 1833 Mr. E. Ravenshaw reported³ the existence of coal in the bed of the Dhela river near Láldhang in Garhwál, where it occurred in thin seams varying from one inch to four inches broad. Similar traces were discovered in the beds of the Chala and Phika streams, but none of any commercial value. The specimens received in Calcutta were nearly all of the same character, "strongly impregnated with

¹ *Gleanings in Science*, I., 230.
Ben., 11., 294.

² *As. Res.*, XVI., 387, 397.

³ *J. A. S.*

sulphuret of iron which forms their fibres, streaking some of them and passes into thick masses of pyrites, decomposing in others. A clean lump had a specific gravity of 1.968 in consequence, and the residual ash was principally iron oxide."

Graphite¹ (plumbago) crops out at the Kalimatiya hill to the north of Almora and on the spur of Banini Devi facing Almora on the Lohughāt road.

In 1850, specimens were sent to England and subjected to examination, when it was found that it could be made serviceable as graphite. Excavations were also made by Major Drummond at Garjoli near Balti, and Palsimi, about three miles from Balti and the same eastward of Almora. The following is the report of Mr. Rose, the mineralogist who tested the specimens sent from Almora :—
 " Graphite is applied to several purposes. When very fine, compact, and of a sufficient cohesion, it is cut up for drawing-pencils. When the texture is loose, or it is otherwise of inferior quality, it is ground down and deprived of foreign substances by washing, as ores of metals are prepared for smelting. The powder thus purified is then used for various purposes, such as crucibles (being refractory or infusible by heat) for burning iron, and reducing the friction of machinery. A new method is now adopted for making artificial pencils, which are scarcely if at all inferior to those sawn out of the finest blocks. The dust of such fine material as your specimens Nos. 4, 8, and particularly No. 10, properly prepared, is subjected to vast hydraulic pressure (several hundred tons), and thus acquires the compactness and solidity necessary for the best purposes. The best kinds of graphite may be known by a pale lead-blue colour, high lustre, unctuousity, and inferior specific gravity. The first nine specimens will answer for pencils, most of them sufficiently pure and compact to be divided for that purpose. All the varieties sent may be used, even No. 13, though connected with much matrix, as it can be deprived of this by grinding and washing. All the varieties of this substance must continue in demand and bring remunerative prices if the expense of mining and conveyance should not be too great." It is also found in Patti Loliba of Garhwāl on the Karnprayág road, and is there used as a dye.

¹ See, further, *Gleanings in Science*, III., 280; *J. A. S.*, XXIV., 203; *Ind. Rec.* (H. D.), XVII., 58; *Rec.*, N.-W. P., III., N. S., 371.

Sulphur is found both in Kumaon and Garhwál. In the former district it occurs in the tract called Mún-syári, and mechanically mixed with carbonate of lime in the beds of the Rám-ganga and Ganjiya rivers. It is also found as green sulphate of iron, and could be obtained in any quantity from the iron pyrites of the copper mines. There are also some sulphureous springs, as those at Naini Tal, Nargoli, and Káthgodám. There are two sulphur springs in Garhwál: the first lies close to the snowy range to the north-east of the temple of Madh Maheswar in pargana Nágpur; the other is on the left bank of the Biri river, two miles above its junction with the Alaknanda. The water of this last is so very strongly impregnated that its existence can be discovered by the smell long before arriving at the spring itself. Neither are made use of any way. Sulphur is also found in the galleries of the lead mines at Maiyár on the Tons in Jaunsár.

Borax or tincal, a native saline compound of boracic acid and soda. The borax and salt fields of Ginari or Húndes, *thá-thaka* or *tháti-tháka*, lie to the north of Bonghwa Tál, across mountains that round the north-east side of the valley of the Shajan river, parallel to the Gangri range, and in the eastern part of the Zjang of Rohto (Rudukh) and at the Chapakani lake. The two salts are obtained from different spots in the same vicinity, and are both worked in the same way by lixiviation from the earth taken from the surface of the ground in which the salts are developed by natural efflorescence. These salt fields are open to all who choose to adventure their labour in them on payment of one-tenth of the produce to the Lhasa Government, who have an excise establishment on the spot.¹ The borax is collected from June to September and sold at the different fairs—Ganpa, Gartoh, Siblam, Shájna, Taklakhar, Dabakhar. It is brought down by the Bhotiya traders and purchased by the merchants of Rámnagar, where it is refined. The process is as follows:—The borax is pounded and placed in shallow tubs, and then covered with water to the extent of a few inches; to this is added a solution of about two pounds of lime dissolved in two parts of water, for every ten maunds (820 pounds) of borax, and the whole mass is well stirred every six hours. Next day it is drained on sieves or

¹ Strachey: J. A. S. Ben., XVII. (2), 57.

cloth, and after this is again dissolved in $2\frac{1}{2}$ times its weight of boiling water, and about sixteen pounds of lime added for the above quantity. It is then filtered, evaporation takes place, and subsequently it is crystallized in funnel-shaped vessels, usually of *kansa*, an alloy of copper and zinc or lead. The loss in weight is about 20 per cent. Borax is used in medicine and the arts. Dry borax acts on the metallic oxides at a high temperature, melting and vitrifying them into beautiful coloured glasses. It is also used as a flux for soldering in goldsmith's work and as a varnish combined with shell-lac. Its principal use is, however, in the manufacture of coloured glass, enamel, and glazed substances.¹

Gypsum is found in pargana Chhakháta. Perhaps the best bed is near the Nihál bridge on the road between Káládhúngi and Naini Tál. In 1850 the late Mr. Tregear, of the Bareilly College, made some very good plaster of Paris from it, which might be found useful in external plastering, as it has the property of expanding on cooling. Gypsum is found in Garhwál, on the banks of the Alaknanda near Panai and Nagrasu. There is also a dark-green variety which the people sometimes make into saucers and bowls.² Captain P. T. Cautley noticed³ the occurrence of gypsum at Sansardhára and Salkot near Dehra and described its appearance and origin, which was further discussed by the Rev. R. Everest, but these papers have now little practical value.⁴

A white saponaceous stone resembling and used for the same purpose as pipeclay is produced in many places. In Garhwál various vessels are turned from it, which when polished have the appearance of marble. They retain liquid, but being extremely brittle are little used.

Asbestos has lately been discovered in a hill to the north of and at a short distance from Ukhimath in Garhwál. It is said to be of very good quality, but it is too far inland to be profitably worked. The people use it medicinally for dressing wounds and burns, and as a wick for oil-lamps, but it may yet be turned to a profitable account as a packing for steam-joints and the like.

¹ Ure, I., 221.
229.

² An. Rec. XVIII. (1), p. 216.

³ J. A. & Rec., I.,

⁴ *Ibid*, 430, and Sherwill on Darjiling gypsum, *ibid*, XXI., 538.

*Silajit*¹ or *Salajit*, a native sulphate of alumina, is found both in Kumaon, Nepál and parganas Painkhandá and Nágpur in Garhwál. It is much sought after and used as a dressing for wounds. It occurs in small light lumps, colour brownish white, externally anhydrous, internally semi-crystalline, fracture slightly fibrous, with a lustre resembling asbestos, porous, containing small cavities lined with scarcely perceptible needle-like crystals, adheres a little to the tongue. Taste acidulous saline, soluble in twice its weight of distilled water, friable. Mr. Stevenson's analysis² gives as its component parts : sulphate of alumina, 95 ; peroxide of iron, 3 ; insoluble matter (silicic acid), 1, and loss, 1. This analysis would appear to point to a specimen of greater purity than those commonly met with in the bazars which, as a rule, have seldom more than 66 per cent. of sulphate of alumina. The lumps generally have an admixture of red sand and frequently portions of micaceous stone are found embedded in them. Some of them have the smooth surfaces of stalactites and are not unlike those deposits. All are readily soluble in water, and when touched with the tongue give the taste of common alum. Dr. Campbell has described³ the Nepálese trade in *Salajit*.

Limestone is found all over the division, both in immense masses exhibiting various shades of colour and structure, and as local tufa deposits. There are three distinct ranges of limestone hills in Garhwál : the first north of the Alaknanda in Nágpur, the second running from Lohbá Patti to the Pindar, and again to the Alaknanda in Patti Bachhansyún, and the third running parallel to the plains and south of the Nayár river. There are also small patches of limestone scattered throughout the district, but not in such large quantities as in the abovementioned ranges. Lime is manufactured at Naini Tal, at Jyúli in the Kharáhi range, half-way between Bágeswar and Almora, at Chiteli, north of Dwárahát, at Simalkha, Baitálghát, and Dhikuli for Ránikhet, and on the new cart-road to Rámnagar. Lime is also made in Borarau, Sor, Sira, Dhyúnirao, and Charál. Two kinds of limestone are used in the Taráí district, the one being obtained from the quarries at the foot of the Kumaon hills, which

¹ Derived from '*sil*,' a stone, and '*jít*,' principle or essence. Ben., II. 321.

² J. A. S., Ben., II., 482.

³ J. A. S.,

give by far the best kind of lime ; the other is the tufa deposit obtained in the small nálas of the tract itself ; this latter kind, however, is of a very inferior quality. First-class limestone costs at the quarries five to eight rupees per 100 maunds ; the tax levied by the Forest Department is eight rupees on that amount, and cartage may be averaged at half a rupee per mile for 100 maunds. Thus the stone is landed at most points in the district for 30 rupees per 100 maunds, and including the expense of burning, a maund of lime costs 10 to 12 annas. This lime will bear two or three portions of pounded brick or *surki*. Second-class lime ready for use now costs 25 rupees, and delivered in Naini Tál, 50-100 rupees per 100 maunds ; it will, however, only bear a proportion of one part of pounded brick to two parts of lime.

Good building stone can be procured in most parts of the hills. At Almora fine-grained, evenly-bedded quartzites and mica-schist form the hill itself, and supply material not to be excelled for durability and facility of dressing. Mica-schist seems to form the principal beds for some distance to the east and west of Almora, reaching to Dwárahát and Mási on the west, Páli, Ránikhet, Síyáhi Devi, Dol, and towards Káli Kumaun to the east, and also in the formation of the Jageswar and Binsar ranges to the north. At Naini Tál the stones used are limestone and clay schist. At Ránikhet a pale-coloured gneiss forms both a handsome and a lasting building stone having the property of hardening by exposure. Sandstone is abundantly found in the lower hills. Gneiss and chlorite-schists are used frequently as building stones in the district. In the Bhábar split boulders are found to answer the purpose of bricks. The Tarái is the only portion of the Kumaon division where bricks are extensively used for building purposes. Nine-inch bricks cost about Rs. 750 per lakh, and the small native bricks Rs. 100. Stone is sometimes carted from the foot of the hills for the better kind of work, but owing to the great expense is, so far as possible, dispensed with.

At Chiteli near Dwárahát there are roofing-slate quarries, now unworked ; also at Dhári in the Bel Patti of Gangoli ; in Borarao Patti, Sult Patti, and at Naini Tál. In Lohba of Garhwál the thin dark-blue slate is procurable, but these last appear to be much inferior to the Chiteli quarry.

Roofing slate.

Alum, known as *phitkari*, is found in different parts of the province, and in abundance in the aluminous shales near Ják village, on the road from Naini Tál to Khairna and as an efflorescence on the micaceous schist in the bed of the Kosi below Almora. The shales contain minute particles of pyrites disseminated throughout their mass, which on becoming decomposed promote the formation of alum and the lixiviation produced by water leaves an encrustation of alum on the rock. A dark-coloured talc called *jalyoshi* is exported to the plains and used as a tonic and febrifuge in medicine. In the neighbourhood of Kotgaon and Giwarsiu near Paori in Garhwál a bitumen or mineral resin is found.

There is no doubt but that in both quantity and quality the metalliferous deposits in this division are good, but the absence of coal and the competition of sea-borne metal have hitherto rendered mining an unproductive speculation, nor does there seem any probability of it attaining any important position among the industries of the province.

REFERENCES.

The following list gives a reference to some of the works and papers on the geology and mineralogy of the North-Western Himálaya :—

1. Report of the mineralogical survey of the Himálaya mountains lying between the rivers Satlaj and Káli : with a geological map, by Captain J. D. Herbert. J. A. S. Ben., XI., 1, p. x.
2. On the mineral productions of the Himálaya mountains, by the same. Trans. A. S., XVIII., i., 236.
3. Journal of a tour from Almora through parts of the province of Kumaon and Garhwál, chiefly in the centre of the hills, by the same. Edited by J. H. Batten. J. A. S. Ben., XIII., 734.
4. Some inquiries in the province of Kumaon relative to geology and other branches of natural science, by Assistant Surgeon J. McClelland, Calcutta, 1835.
5. Geological specimens from Kumaon and Garhwál, by the same. J. A. S. Ben., II., 633.
6. Notice of some fossil impressions occurring in the transition limestone of Kumaon, by the same. *Ibid*, VI., 653.
7. On the geology of part of the Himálaya mountains and Tibet, by Captain R. Strachey. J. Geo. Soc., VII. (1851).
8. Geology of Dehra Dún, by Dr. Royle. J. A. S. Ben., I., 96, 249.
9. Geological sketch of Mussooree and Landour in the Himálaya, together with an abstract of the thermometrical register kept at Landour during the year 1831, by F. H. Fisher. *Ibid*, I., 193.
10. Geological observations made on a journey from Mussooree to Gangotri by Rev. R. Everest. *Ibid*, IV., 690.

11. On the origin of the Kumaon Lakes, by V. Ball. *Rec. Geol. Sur.*, XI., 2, 174.
12. Note on a trip over the Milam pass, Kumaon, by T. W. H. Hughes, with a description of the fossils by Dr. Waagen. *Ibid*, 182.
13. A Manual of the Geology of India, by H. B. Medlicott and W. T. Blandford, Calcutta, 1879.
14. *Sivatherium giganteum*, a new fossil ruminant genus from the valley of the Markanda in the Siwálík branch of the Sub-Himáláyan mountains, by Dr. H. Falconer and Captain P. T. Cantley. *As. Res.*, XIX., I, and *J. A.S. Ben.*, V., 38. VI, 152.
15. Note on the fossil hippopotamus of the Siwálík hills, by the same. *As. Res.*, XIX., 89.
16. The fossil gharial of the Siwálík hills, by the same. *Ibid*, 82.
17. The fossil bear of the same, by the same. *Ibid*, 193.
18. The fossil crocodile of the same, by the same. *Ibid*, 25.
19. The fossil tiger of the same, by the same. *Ibid*, 135.
20. The fossil camel of the same, by the same, *ibid*, 115; and by Lieutenant W. E. Baker. *J. A. S. Ben.*, IV., 694 : IX, 620.
21. An additional fossil species of the order quadrumana, by the same. *Ibid*, VI, 354.
22. Specimens of the hippopotamus and other fossil genera in the Dádúpur collection by H. M. Durand, *As. Res.*, XIX., 54, and selected specimens of the Sub-Himáláyan fossils in the Dádúpur collection, by Lieutenant W. E. Baker. *J. A. S. Ben.*, IV. 565, and Durand, V., 291, 486, 661, 739.
23. Discovery of fossils in the Siwálíks, by Captain Cantley. *Ibid*, III., 592.
24. Fossil elephant's tooth from the same, by Lieutenant Baker. *Ibid*, 638.
25. The fossil elk of the Himáláya, by the same *Ibid*, IV., 506.
26. Synopsis of fossil genera and species in the collection of Dr. Falconer and Captain Cantley. *Ibid*, IV., 706.
27. Collections of fossils presented by Colonel Colvin. *Ibid*, V., 58, 179.
28. Note on the teeth of the mastodon, by Captain Cantley. *Ibid*, 294, 768.
29. Fossil remains of the smaller carnivora from the Sub-Himáláya, by Lieutenants Baker and Durand. *Ibid*, 585.
30. Notice of the skull (fragment) of a gigantic fossil batrachian from the Nahan field, by Dr. T. Cantor. *Ibid*, VI., 538.
31. Note on the occurrence of fossil bones in the Siwálík range eastward of Hardwár, by Dr. H. Falconer. *Ibid*, VI., 233.
32. Note on a fossil ruminant species allied to Giraffids in the Sewálík hills, by Captain P. T. Cantley. *Ibid*, VII., 658.
33. Note relating to Siwálík fauna, by H. B. Medlicott. *Ibid*, XXXIV., ii., 63.
34. A sketch of the history of the fossil vertebrate of India, by R. Lyddeker. *Ibid*, XLIX., ii., 8.
35. *Fauna antiqua Sivalensis*, by Falconer and Cantley. London, 1846-49-56.
36. Palaeontological Memoirs and notes of H. Falconer, by C. Murchison. London, 1868.
37. Palaeontologia Indica, published by the Geological Survey, Calcutta.
38. Memoirs of the Geological Survey of India, by the same.
39. Records of the Geological Survey of India, by the same.

CHAPTER VII.

SCIENTIFIC BOTANY.

CONTENTS.

A list of the *flora* of Garhwál, Jaunsar-Báwar and the Dehra Dún, by Dr. King. Tract to which the list refers. The Siwálíks. Sub-Siwálík tract. Dehra Dún. Himálayan tract. Characteristic vegetation from the plains to Mussooree. From Mussooree northwards. A list of the *flora* of Kumaon, and the Kumaon Bhábar, by Dr. Watson.

LIST¹ OF THE PLANTS OF GARHWÁL, JAUNSAËR-BÁWAR AND THE DEHRA DÚN.

THE tract of country of which the vegetation is to be considered in the following pages presents very varied conditions in respect of physical configuration, soil, and climate. Its boundaries are—on the west the river Tons, a tributary of the Jumna, and on the east the Bhágirathi branch of the Ganges; on the north the snowy range of the Himálaya, and on the south the Siwálík range.

The Siwálíks run parallel to the Himálaya, and are separated from the latter by the Dehra Dún, a flat valley having an elevation of from 1,800 to 2,250 feet above the level of the sea. The highest peaks of the Siwálíks do not rise much above 3,000 feet, and the majority are much lower, but their outline against the sky is jagged and picturesque in the extreme, and the part of them with which we have to deal is about seven miles wide. Their southern slopes are ploughed by the force of the south-western monsoon into numerous deep narrow valleys, and the steeply scarped cliffs have a singularly bold beauty, totally different from anything to be seen in the outer parts of the Himálaya behind them. On their northern aspect, the Siwálíks slope gently into the valley of the Dún, and the scenery, though beautiful, is tamer than on the southern side. The rainfall on the range during the south-west monsoon is about 60 inches, but from the porous character

¹ This list has been kindly prepared by Mr. G. King, M.B., F. L. S., Director of the Royal Botanical Gardens, Calcutta, for this work.

of the soil and the steepness of the slopes, water drains off with extraordinary rapidity. The drainage passes away by innumerable channels, which, except for a few days in the year, are wide wastes of dry gravel and boulders. After each heavy shower these suddenly become filled with rushing torrents, which subside as speedily as they rose. Except during the rainy season, the range is very dry, and water is so scarce as to make it almost uninhabitable.

Where the Siwálíks merge into the plains on the southern side, there are tracts of forest and grass equivalent to those covering the Blábar and Taráí of the Himálaya in the districts where there is no Siwálik range.

Sub-Siwálik tract.

The vegetation of the sub-Siwálik belts is to a great extent but a reproduction in miniature of that of the latter. The flora of the southern face of the Siwálíks comprises many plants of the plains, with the addition of such species as *Hypericum cernuum* and *Cochlospermum Gosypium*. Perhaps the most striking tree found in the Siwálíks is the long-leaved pine (*Pinus longifolia*), a Himálayan species, for which the Siwálíks form an outlying station. A large proportion of the arboreal vegetation of the Siwálíks consists of species which occur both on the lower slopes of the Himálaya and on the hilly districts of central and southern India, thus affording a good illustration of the width of distribution of Indian plants. As examples may be mentioned *Bombax malabaricum*, *Buchanania latifolia*, *Spondias mangifera*, *Boswellia serrata*, *Garuga pinnata*, *Spatholobus Roxburghii*, *Albizia stipulata*, *Anogeissus latifolia*, *Ulmus integrifolia*, and others. Epiphytal orchids are absent, and ferns are but few in the Siwálíks. One labiate plant (*Eremostachys superba*) is confined to one or two spots in the Siwálíks, and is one of the best examples known to me in India of a truly local plant. *Sál* (*Shorea robusta*) of very stunted growth is abundant in the southern sub-Siwálik tract and on both sides of the range, as also over the whole of the Dún. This tree is here near its western limit; a little way to the west of the Jumna it disappears altogether.

The Dehra Dún is a charming valley. The Siwálíks on the south shut it in from the plains, and, until recently, not only most effectually protected it from the hot winds in summer, but gave it a higher winter temperature than the plains outside. The extensive and ruthless

Dehra Dún.

clearances of the Siwálik forest of recent years are, however, believed (and apparently with good reason) to have caused an increase of the summer temperature of the Dún. The soil of the Dún consists chiefly of sandstone débris washed down from the Himálaya. The drainage is excellent, especially near the base of the Himálaya where the larger débris has been deposited, and the water level is everywhere far below the surface. The drainage water passes off by the Asan into the river Jumna, and by the Son and Súswa into the Ganges. The watershed of these streams runs across the valley in the line of the station of Dehra. The rainfall is about 80 inches, but, although only a small proportion of this total falls during any other than the rainy season, vegetation is kept wonderfully green during the whole year by the moisture prevalent in the atmosphere. Formerly the greater part of the Dún was covered by forest, of which the prevailing tree was *adi*, but of late years much of the forest has disappeared. A certain proportion of the cleared land has been devoted to village and some to tea cultivation, but much of it has been invaded by tall, coarse grasses useless alike to man and beast. Like the Siwálíks, the Dún is ravaged every hot season by fires from which the arboreal vegetation suffers to a degree which can only be appreciated by one who has lived in the forest. Unless these fires are checked with a high hand, the utter destruction of all forest vegetation in the Dún is but a question of time. The flora of the Dún presents an interesting mixture of species common on the plains outside, and of plants from the lower hills. Notable amongst the latter are the pretty little *Gentiana pedicellata*, and in the rains, here and there, *Platystemma violoides*; whilst amongst shrubs and climbing plants *Lespedeza*, *Indigofera pulchella*, *Jasminum pubescens*, and *Conibretum decandrum* are perhaps the most noteworthy. The dense canebrakes of the Tará and Bhábur of the trans-Gangetic sub-Himálayan tract are represented in the Dún by a few patches of a single species (*Calamus Royleanus*), and this is not found westward of the Nálapani swamp, three miles from the station of Dehra. Epiphytal *Scitamineæ* are unknown in the Dún: a few orchids, however, occur, the most prominent among which are *Aerides affine* and *odoratum*, *Saccolabium Gharwalicum*, and *Oberonia iridifolia*. *Peperomia reflexa* is found here and there on trees in damp spots, and there are some epiphytal ferns, such as *Polypodium lineare*.

The remainder of the region under review consists of a large irregularly shaped tract of the Himálaya proper, narrowed at its southern limit, but expanding towards the Himálayan tract. snowy range, part of which it includes.

The physical features of this area are so varied that it would be out of place to attempt to describe them here. It must suffice to remind the reader that, while the lower and outer ranges have a climate not very different from that of the plains of India, the higher peaks, rising as they do to 18,000 feet above the sea-level, are covered with perpetual snow. As regards vegetation, the tract may be divided into three zones: the *sub-tropical* covering the slopes of the lower and outer hills, and following the courses of the deep hot valleys far into the interior of the range; the *temperate* covering the middle elevations; and the *arctic* confined to the greater heights and bounded on the north by the snow line itself. The transition between these zones is gradual. In consequence mainly of its greater distance from the sea, the rainfall of the western Himálaya is much less than that of the eastern part of the chain. But not only does the climate steadily increase in dryness from east to west, but the rainfall on the outer ranges is much heavier than in the interior. At Mussooree, for example, it amounts to about 80 inches, while at Harsil, in the northern end of the Bhágirathi valley, the rainy season is represented by a short period of misty weather, and actual showers are but rare. The intervening country has a rainfall graduated between these two extremes.

In ascending the Himálaya from the Dún to Mussooree, new species meet the eye step by step. Among Characteristic vegetation from the plains to Mussooree. trees, *Bauhinia retusa*, *Engelhardtia Colebrookiana*, *Rhododendron arboreum*, and the Oaks are the most striking new forms. *Pinus longifolia*, already met with on the Siwálíks, becomes abundant. Berberries, various *Bahmerius* and other *Urticaceæ*, *Hamiltonia*, *Leptodermis*, various species of *Viburnum*, *Clematis* and *Rosa* at first mingle with, and finally replace, the sub-arboreal woody vegetation of the base of the hills. During the rains the change in vegetation, as one ascends, is even more striking. On steep banks and faces of rock by the roadside the eye is delighted by such charming species as *Chirita bifolia*, *Didymocarpus aromaticus* and *macrophyllus*, *Platystemma violoides*,

Selaginella Jacquemontii, *Androsace sarmentosa* and *incisa*, *Fragaria indica*, *Saxifraga ligulata*, *Argostemma sarmentosum*, and *Begonia picta*. Ferns too abound both on the ground and on trees. A few epiphytal orchids, the *Peperomia* already mentioned, and *Remusatia vivipara* are found on the trees. But the epiphytal vegetation here is but poor and scanty when compared with that of the eastern Himálaya, thus illustrating in the most forcible way the comparative dryness of the atmosphere throughout the year. On these lower slopes a dwarf palm (*Phoenix acaulis*) and a pretty little bamboo called *ringdl* (*Arundinaria falcata*) are not unfrequent; a larger species of bamboo (*Dendrocalamus strictus*), which occurs in plenty, is common also in the Dún and Siwálíks.

At the level of Mussooree temperate forms begin to appear and the botanist revels in plants belonging to such familiar European genera as *Ranunculus*, *Thalictrum*, *Anemone*, *Geranium*, *Potentilla*, *Rubus*, *Rumex*, *Pedicularis*, and *Habenaria*. Umbelliferous plants and *Polygonaceæ* are common, and terrestrial orchids of northern

From Mussooree north-wards. forms are not unfrequent during the rains.

As the higher elevations are approached, the vegetation assumes the characteristic Alpine type. *Astragali*, *Corydalis* of different species, numerous *Caryophylleæ*, *Saxifrages*, *Sedums*, Primroses, Gentians and *Carices* abound, and in the carpet of rich green grass are here and there studded the magnificent Aconites, *Meconopsis* and *Pæonia emodi*. Prominent amongst the forest vegetation are the stately cedar, the rigid silver spruce and the graceful weeping pine. The twisted cypress towers aloft, and the solemn yew stretches out its sombre arms. Lichens, mosses and *fungi* abound, and *algæ* are numerous in the streams.

The following list of the flowering plants and ferns does not profess to be complete. It contains only the names of species of which there are specimens in the Herbarium of the Calcutta Botanic Garden, or in my own collection. The former Herbarium ought to contain every species occurring within the British Empire in India. It has, however, been brought together principally by desultory private effort, and is consequently unequal and defective. As a rule, there are included in this list indigenous plants only; the few others that occur are marked either as *naturalized* or *introduced*. Certain plants which are believed to be natives, but which are

found only in cultivation, are marked as *cultivated*. For much assistance in revising this list I am indebted to Mr. W. Waterfield, C. S.

Natural Order 1.—Ranunculaceae.

- Clematis Nepaulensis*, DC.
 " *montana*, Ham.
 " *barbellata*, Edgw.
 " *grata*, Wall.
 " *Gouriana*, Roxb.
 " *puberula*, Hf. & T.
 " *graveolens*, Lindl.
 " *orientalis*, L.
 " *nutans*, Royle.
 " *acuminata*, DC.
 " *connata*, DC.
 " *Buchananiana*, DC.
Anemone *vitifolia*, Ham.
 " *obtusiloba*, Don.
 " *rupestris*, Wall.
 " *rivularis*, Ham.
 " *polyanthes*, Don.
 " *narussiflora*, L.
 " *tetrasepala*, Royle.
 " *elongata*, Don.
Thalictrum *elegans*, Wall.
 " *platycarpum*, Hf. & T.
 " *cultratum*, Wall.
 " *Chelidoni*, Hf. & T.
 " *reniforme*, Wall.
 " *pauciflorum*, Royle.
 " *rostellatum*, Hf. & T.
 " *Punduanum*, Wall.
 " *saniculiforme*, DC.
 " *Javanicum*, Bl.
 " *foliolosum*, DC.
 " *minus*, L.
Callianthemum cachenirianum, Camb.
Adonis festalis, L.
Ranunculus aquatilis, L.
 " *Lingua*, L.
 " *pnichellus*, C. A. Mey.
 " *lobatus*, Jacquem.
 " *hyperboreus*, Rottb.
 " *affinis*, Br.
 " *hirtellus*, Royle.
 " *sceleratus*, L.
 " *diffusus*, DC.
 " *lætus*, Wall.
 " *pensylvanicus*, L.
 " *arvensis*, L.
Oxygraphis polypetalæ, Hf. & T.
Caltha palustris, L.
Trollius acaulis, Lindl.
Isopyrum thalicteroides, L.
 " *grandiflorum*, Fisch.
Nigella sativa, L. *naturalized*.
Aquilegia vulgaris, L.
Delphinium denudatum, Wall.
 " *coruleum*, Jacquem.
 " *elatum*, L.
 " *vestitum*, Wall.

Natural Order 1.—Ranunculaceae—
(concluded).

- Delphinium Cashmirianum*, Royle.
 " *Ajacis*, L., *introduced*.
Aconitum Lycoctonum, L.
 " *palmatum*, Don.
 " *ferox*, Wall.
 " *Napellus*, L.
 " *heterophyllum*, Wall.
Actæa spicata, L.
Cimicifuga foetida, L.
Pæonia emodi, Wall.

Natural Order 2.—Magnoliaceæ.

- Michelia Champaca*, L., *introduced*.
Schizandra grandiflora, Hf. & T.
 " *propinqua*, Hf. & T.

Natural Order 3.—Anonaceæ.

- Miliusa velutina*, Hf. & T.

Natural Order 4.—Menispermaceæ.

- Tinospora cordifolia*, Miers.
Cocculus laurifolius, DC.
 " *villosus*, DC.
Stephania elegans, Hf. & T.
 " *rotunda*, Lour.
Cissampelos Pareira, L.

Natural Order 5.—Berberidæ.

- Holboellia latifolia*, Wall.
Berberis nepalensis, Spreng.
 " *vulgaris*, L.
 " *aristata*, DC.
 " *Lycium*, Royle.
 " *asiatica*, Roxb.
Podophyllum emodi, Wall.

Natural Order 6.—Nymphaeaceæ.

- Nelumbium speciosum*, Willd., *introduced*.
Nymphaea Lotus, L.

Natural Order 7.—Papaveraceæ.

- Papaver somniferum*, L., *cultivated*.
 " *dubium*, L.
Argemone mexicana, L., *naturalized*.
Meconopsis aculeata, Royle.
 " *robusta*, Hf. & T.
 " *nepalensis*, DC.
Stylophorum lactucoides B. & Hf.

Natural Order 8.—Fumariaceæ.

- Dicentra* Roylei, Hf. & T.
 " scandens, Walp.
Corydalis rufaefolia, Sibth.
 " cachemirianna, Royle.
 " crithmifolia, Royle.
 " elegans, Wall.
 " Gervasianna, Wall.
 " sibirica, Pers.
 " pernuta, Royle.
 " chierophylla, DC.
 " meifolia, Wall.
 " fiabellata, Edgw.
Fumaria parviflora, Lamk.

Natural Order 9.—Cruciferaæ.

- Nasturtium* officinale, Br.
 " palustre, DC.
 " montanum, Wall.
Barbarea vulgaris, Br.
Arabis glabra, Crantz.
 " alpina, L.
 " amplexicaulis, Edgw.
Cardamine hirsuta, L.
 " impatiens, L.
 " macrophylla, Willd.
Allyssum canescens, DC.
Draba alpina, L.
 " incana, L.
 " lasiophylla, Royle.
 " fladnitzensis, Wulf.
Sisymbrium mollissimum, C. A. Meyer.
 " himalaicum, Hf. & T.
 " Thalianum, Gay & Monn.
 " rupestre, Edgw.
 " strictum, Hf. & T.
 " Wallichii, Hf. & T.
 " Sophia, L.
 " Columnne, Jacq.
 " Alliaria, Scop.
Eutrema primulaefolium, Hf. & T.
Erysimum hieraciifolium, L.
 " altaicum, C. A. Meyer.
 " Thomsoni, Hf.
Braya rosea, Bunge.
Brassica nigra, Koch.
 " campestris, L.
 " juncea, Hf. & T.
Fruca sativa, Lamk.
Capsella Bursa-pastoris, Moench.
Lepidium sativum, L.
 " Draba, L.
 " capitatum, Hf. & T.
Megacarpus polyandra, Benth.
Thlaspi arvense, L.
 " alpestre, L.
Iberidella Andersoni, Hf. & T.
Raphanus sativus, L., *cultivated*.
Goldbachia laevigata, DC.
Chorispora tenella, DC.

Natural Order 10.—Capparidææ.

- Cleome* viscosa, L.
Gynandropsis pentaphylla, DC.

Natural Order 10.—Capparidææ—(concl.)

- Mærua* arenaria, Hf. & T.
Capparis spinosa, L.
 " horrida, L. f.
Cratæva religiosa, Forst.

Natural Order 11.—Violaceæ.

- Viola* biflora, L.
 " Patrinii, DC.
 " serpens, Wall.
 " kunawarensis, Royle.
Ionidium suffruticosum, Ging.

Natural Order 12.—Bisincæ.

- Cochlospermum* Gossypium, DC.
Flacourtia Ramontchii, L'Herit.
 " sepaiaria, Roxb.
Xylosma longifolium, Clos.

Natural Order 13.—Pittosporææ.

- Pittosporum* floribundum, W. & A.
 " eriocarpum, Royle.

Natural Order 14.—Polygalææ.

- Polygala* triphylla, Ham.
 " crotalarioides, Ham.
 " abyssinica, Fresen.
 " leptalea, DC.
 " persicariæfolia, DC.
 " chinensis, L.
 " sibirica, L.
Salomonis cantoniensis, Lour.

Natural Order 15.—Caryophyllææ.

- Gypsophila* cerastioides, Dom.
Saponaria Vaccaria, L.
Silene inflata, Sm.
 " conoidea, L.
 " Fatconeriana, Benth.
 " Griffithii, Boiss.
 " gallica, L.
 " Moorcroftiana, Wall.
 " Webbiana, Wall.
 " Stracheyi, Edgw.
Cucubalus bacciferus, L.
Lychnis apetala, L.
 " brachypetala, Hort. Berol.
 " multicaulis, Wall.
 " inflata, Wall.
 " indica, Benth. *var. fimbriata*, Wall.
 " nutans, Benth.
 " pilosa, Edgw.
Cerastium dahuricum, Fisch.
 " vulgatum, L.
 " Thomsoni, Hf.
Stellaria crispata, Wall.
 " paniculata, Edgw.
 " semivestita, Edgw.

Natural Order 13.—Caryophyllæ—
(concluded).

- Stellaria aquatica*, Scop.
 " *media*, L.
 " *Webbiana*, Wall.
 " *latifolia*, Benth.
 " *longissima*, Wall.
 " *saxatilis*, Ham.
 " *glauca*, With.
 " *decumbens*, Edgw.
Arenaria foliosa, Royle.
 " *festucoides*, Benth.
 " *serpyllifolia*, L.
 " *orbiculata*, Royle.
 " *glanduligera*, Edgw.
 " *Stracheyi*, Edgw.
 " *holosteoides*, Edgw.
 " *Benthani*, Edgw.
Sagina procumbens, L.
Spergula arvensis, L., in fields.
 " *pentandra*, L., ditto.
Drymaria coriata, Willd.
Polycarpea corymbosa, Lamk.

Natural Order 16.—Portulacæ.

- Portulaca oleraceæ*, L.

Natural Order 17.—Tamariscinæ.

- Tamarix gallica*, L.
 " *dioica*, Roxb.
Myricaria germanica, Desv.
 " *elegans*, Royle.

Natural Order 18.—Elatinæ.

- Bergia ammannioides*, Roxb.

Natural Order 19.—Hypericinæ.

- Hypericum cornutum*, Roxb.
 " *cordifolium*, Choisy.
 " *lysimachioides*, Wall.
 " *patulum*, Thunb.
 " *perforatum*, L.
 " *elodeoides*, Choisy.
 " *nepaulense*, Choisy.

Natural Order 20.—Ternstroemiaceæ.

- Eurya japonica*, Thunb.
Actinidia callosa, Lindl.
Saurauja nepaulensis, DC.
Camellia Thea, Link. Tea, introduced
 and cultivated.

Natural Order 21.—Dipterocarpaceæ.

- Shorea robusta*, Gaertn., S&L.

Natural Order 22.—Malvaceæ.

- Malva verticillata*, L.
 " *silvestris*, L.
 " *rotundifolia*, L.
 " *parviflora*, L.
Skia rhombifolia, L.
 " *cordifolia*, L.
Abutilon polyandrum, Schl.
 " *indicum*, G. Don.
 " *graveolens*, W. & A.
Urena lobata, L.
 " *repanda*, Roxb.
Hibiscus Trionum, L.
 " *furcatus*, Roxb.
 " *Solandra*, I'Herit.
 " *ficulneus*, L.
 " *purgens*, Roxb.
 " *Abelmoschus*, L.
 " *cancellatus*, Roxb.
 " *esculentus*, L., naturalized.
Theopesia Lampas, Dalz. & Gibs.
Gossypium herbaceum, L., introduced.
Kydia calycina, Roxb.
Bombax malabaricum, DC.

Natural Order 23.—Sterculiaceæ.

- Sterculia villosa*, Roxb.
 " *Roxburghii*, Wall.
 " *fulgens*, Wall.
Helicteres laora, L.
Pterospermum acerifolium, Willd., rare.
Melochia corchorifolia, L.
Waltheria americana, L.
Abroma augusta, L., probably introduced.

Natural Order 24.—Tiliaceæ.

- Grewia oppositifolia*, Roxb.
 " *vestita*, Wall.
 " *tiliaefolia*, Vahl.
 " *asiatica*, L.
 " *sapkila*, Roxb.
 " *scabrophylla*, Roxb.
 " *polygama*, Roxb.
 " *laevigata*, Vahl.
Corchorus fascicularis, Lamk.
 " *acutangulus*, Lamk.
Triumfetta pilosa, Roth.
 " *rhomboides*, Jacq.
 " *annua*, L.

Natural Order 25.—Linææ.

- Linum usitatissimum*, L.
 " *mysorensis*, Heyne.
Reinwardtia trigyna, Planch.

Natural Order 26.—Malpighiaceæ.

- Hiptage Madagblota*, Gaertn.
Aspidopterys Wallichii, Hf.
 " *nutans*, Hf.

Natural Order 27.—Zygophyllaceæ.

Tribulus terrestris, L.

Natural Order 28.—Geraniaceæ.

Geranium palustre, L.
 " *collinum*, M. Bieb.
 " *Wallichianum*, Sweet.
 " *nepalense*, Sweet.
 " *polyanthes*, Edgew. & Hf.
 " *rotundifolium*, L.
 " *molle*, L.
 " *Robertianum*, L.
 " *lucidum*, L.
 " *ocellatum*, Camb.
Oxalis corniculata, L.
 " *Acetosella*, L.
Biophytum sensitivum, DC.
 " *Reinwardtii*, Walp.
Impatiens Royloi, Walp.
 " *Thomsoni*, Hf.
 " *sulcata*, Wall.
 " *amplexicaulis*, Edgew.
 " *Balsamina*, L., Mujethi.
 " *scabrida*, DC.
 " *bicornuta*, Wall.
 " *amphorata*, Edgew.

Natural Order 29.—Rutaceæ.

Bœninghausenia albiflora, Reichb.
Dictamnus albus, L.
Xanthoxylum alatum, Roxb.
 " *oxyphyllum*, Edgew.
Toddalia aculeata, Pers.
Skimmia Laureola, Hf.
Glycosmis pentaptylla, Corr.
Limonia acidissima, L.
Murraya exotica, L.
 " *Kanigii*, Spreng.
Clausena pentaptylla, DC.
Citrus medica, L.
 " *Aurantium*, L.
Ægle Marmelos, Corr.
Pteronia Elephantum, Corr.

Natural Order 30.—Simarubaceæ.

Ailanthus excelsa, Roxb.
Picrasma quassoides, Bourn.

Natural Order 31.—Ochnaceæ.

Ocina pumila, Ham.

Natural Order 32.—Bursaceæ.

Bowellia serrata, Roxb.
Garuga pinnata, Roxb.

Natural Order 33.—Meliaceæ.

Melia Azadirach, L., Bukain.
 " *composita*, Willd.
Azadirachta Indica, Juss. Nim.
Adreia serrata, Royle.
 " *Touua*, Roxb.

Natural Order 34.—Olacineæ.

Olax nana, Wall.

Natural Order 35.—Mliaceæ.

Ilex dipyrrena, Wall.
 " *excelsa*, Wall.
 " *odorata*, Ham.

Natural Order 36.—Celastrineæ.

Euonymus pendulus, Wall.
 " *echinatus*, Wall.
 " *grandiflorus*, Wall.
 " *Hamiltonianus*, Wall.
 " *lanceus*, Ham.
 " *tingens*, Wall.
Celastrus paniculata, Willd., Makagni.
Gymnosporia rufa, Wall.
 " *montana*, Roxb.
 " *Boyleana*, Wall.
Ehmodendron glaucum, Pers.

Natural Order 37.—Rhamnaceæ.

Ventilago calyculata, Tulasac.
Zyzyphus Jujuba, Lamk.
 " *Oenoplia*, Mill.
 " *vulgaris*, Lamk.
 " *xylopyra*, Willd.
 " *nummularia*, W. & A.
 " *oxyphylla*, Edgew.
Rhochemia floribunda, Wall.
Rhamnus purpurea, Edgew.
 " *procumbens*, Edgew.
 " *virgatus*, Roxb.
 " *triqueter*, Wall.
Sageretia oppositifolia, Brongn.
 " *thecana*, Brongn.
Scutia indica, Brong.
Gouania leptostachya, DC.
 " *nepalensis*, Wall.
Hovenia dulcis, Thunb.

Natural Order 38.—Ampelideæ.

Vitis capreolata, Don.
 " *latifolia*, Roxb.
 " *vinifera* L., *introduced*.
 " *Limul*, Wall.
 " *adnata*, Wall.
 " *lanata*, Roxb.
 " *himalayana*, Brand.
 " *parvifolia*, Roxb.
 " *obtecta*, Wall.
 " *divaricata*, Wall.
Loea sambucina, Willd.
 " *robusta*, Roxb.
 " *alata*, Edgew.
 " *aspera*, Wall.

Natural Order 39.—Sapindaceæ.

Æsculus indica, Colebr.
Schleichera trifuga, Willd.
Sapindus trifoliatus, L.
 " *Mukorosi*, Gaertn., *probably introduced*.

Natural Order 39.—Sapindaceæ—
(concluded).

- Sapindus* *Danura*, Voigt, *probably introduced*.
Acer *oblongum*, Wall.
 „ *castaneum*, Wall.
 „ *lavigatum*, Wall.
 „ *pictum*, Thunb.
 „ *villosum*, Wall.
 „ *caudatum*, Wall.
Mcianthus *major*, L., *introduced*.
Staphylea *Emodi*, Wall.

Natural Order 40.—Sabiaceæ.

- Sabia* *paniculata*, Edgw.
 „ *campanulata*, Wall.
Meliosma *simplicifolia*, Bl.
 „ *dielleniifolia*, Bl.

Natural Order 41.—Anacardiaceæ.

- Rhus* *vernicefera*, DC.
 „ *acuminata*, DC.
 „ *seminalata*, DC.
 „ *Cotinus*, L.
 „ *succedanea*, L.
 „ *parviflora*, Roxb.
Spondias *Mangifera*, Pers.
Mangifera *indica*, L.
Buchanania *latifolia*, Roxb.
Oxlinia *Wodier*, Roxb.
Nemecarpus *Anacardium*, L.

Natural Order 42.—Coriariæ.

- Coriaria* *nepalensis*, Wall.

Natural Order 43.—Moringæ.

- Moringa* *pterigosperma*, Gaertn.

Natural Order 44.—Leguminosæ.

- Piptanthus* *nepalensis*, Don.
Thermopsis *barbata*, Royle.
Crotalaria *prostrata*, Roxb.
 „ *alata*, Roxb.
 „ *sericea*, Retz.
 „ *juncosa*, L.
 „ *tetragona*, Roxb.
 „ *albida*, Heync.
 „ *neglecta*, W. & A.
Argyrobium *flaccidum*, J. et S.
 „ *roscum*, J. et S.
Parochetus *communis*, Ham.
Trigonolla *Pænum-Gruceum*, L.
 „ *gracilis*, Benth.
Medicago *lupulina*, L.
 „ *laciniata*, All.
 „ *sativa*, L., *cultivated*.
Melilotus *parviflora*, Desf.
 „ *alba*, Lamk.
 „ *officinalis*, L.

Natural Order 44.—Leguminosæ—
(continued).

- Trifolium* *repens*, L.
 „ *pratense*, L.
Lotus *corniculatus*, L.
Indigofera *linifolia*, Retz.
 „ *trifoliata*, L.
 „ *trita*, L.
 „ *tinctoria*, L.
 „ *pulchella*, Roxb.
 „ *Dosua*, Ham.
 „ *atropurpurea*, Ham.
 „ *hiranta*, L.
 „ *Girardinia*, Wall.
 „ *heterantha*, Wall.
Tephrosia *purpurea*, Pers.
 „ *tenuis*, Wall.
Millettia *auriculata*, Baker.
Caragana *brevispina*, Royle.
Colutea *nepalensis*, Sims.
Astragalus *graveolens*, Ham.
 „ *trichocarpus*, Grah.
 „ *adesmisiifolius*, Benth.
 „ *leucocephalus*, Grah.
 „ *hoenckioides*, Benth.
 „ *rhizanthus*, Royle.
 „ *Webbianus*, Grah.
 „ *multiceps*, Wall.
Oxytropis *microphylla*, DC.
Æschynomene *indica*, L.
Snitthia *sensitiva*, Ait.
 „ *ciliata*, Royle.
Zornia *diphylla*, Pers.
Ononis *dalbergioides*, Benth.
Desmodium *oxyphyllum*, DC.
 „ *gangeticum*, DC.
 „ *huxiflorum*, DC.
 „ *gyrans*, DC.
 „ *podocarpum*, DC.
 „ *tiliæfolium*, Dou.
 „ *floribundum*, G. Don.
 „ *concinnum*, DC.
 „ *polycarpum*, DC.
 „ *parvifolium*, DC.
 „ *triflorum*, DC.
Urtia *picta*, Desv.
 „ *lagopus*, DC.
 „ *lagopodoides*, DC.
Alysicarpus *virginialis*, DC.
 „ *huplenrifolius*, DC.
Lespedeza *elegans*, Camb.
 „ *sericea*, Miq.
 „ *erucarpa*, DC.
 „ *juncosa*, Pers.
Vicia *sativa*, L.
 „ *rigidula*, Royle.
 „ *hiranta*, Koch.
 „ *tenara*, Grah.
 „ *tenuifolia*, Roth.
 „ *tetrasperma*, Munch.
Lens *esculenta*, Moench, *cultivated*.
Lathyrus *sativus*, L.
 „ *luteus*, Buk.
 „ *Aphaca*, L.

Natural Order 44.—Leguminosæ—(concl'd.) *Natural Order 45.—Rosaceæ—(concl'd.)*

- Pisum sativum*, L., *introduced*.
Abrus precatorius, L.
Clitoria Ternatea, L.
Dumasia villosa, DC.
Shuteria involucreta, W. & A.
Glycine Soja, S. et Z.
Erythrina arborescens, Roxb.
 " *resupinata*, Roxb.
 " *suberosa*, Roxb.
Mucuna pruriens, DC.
 " *atropurpurea*, DC.
Butea frondosa, Roxb.
Spatholobus Roxburghii, Benth.
Pueraria tuberosa, DC.
Phaseolus Mungo, L.
 " *aconitifolius*, Jacq. } *cultivated*.
Vigna Catjang, End.
Dolichos Lablab, L.
Cajanus indicus, L.
Atylosia scarabaeoides, Benth.
 " *mollis*, Benth.
Rhynchosia minima, DC.
 " *Falconeri*, Baker.
Flemingia semialata, Roxb.
 " *strobilifera*, R. Br.
 " *vestita*, Benth.
Dalbergia Sissoo, Roxb.
 " *lanceolaria*, L.
 " *hircina*, Benth.
 " *volubilis*, Roxb.
Pongamia glabra, Vent.
Sophora mollis, Wall.
Cassalpinia Bonducella, Flemm.
 " *sepiaria*, Roxb.
Cassia fistula, L.
 " *Tora*, L.
 " *occidentalis*, L.
 " *Abus*, L.
 " *pumila*, Lamk.
 " *Wallichiana*, DC.
Bauhinia racemosa, Lamk.
 " *variegata*, L.
 " *retusa*, Roxb.
 " *Vahlia*, W. & A.
 " *purpurea*, L.
Mimosa rubicaulis, Lamk.
 " *pudica*, L. Sensitive plant, *naturalized*.
Acacia Catechu, Willd.
 " *pennata*, Willd.
 " *cassia*, W. and A.
 " *Farnesiana*, Willd, *naturalized*.
 " *eburnea*, Willd.
Albizia lucida, Benth.
 " *Lebbek*, Benth.
 " *odoratissima*, Benth.
 " *procera*, Benth.
 " *Julibrissin*, Boiv.
 " *stipulata*, Boiv.
Natural Order 45.—Rosaceæ.
Prunus Puddum, Roxb.
 " *Padua*, L.

- Prunus domestica*, L. Alucha, Plum, *introduced*.
 " *armeniaca* L., Zarl Alu, Apricot, *cultivated*.
Amygdalus persica, L., Aru, Peach, *introduced*.
Prinsepia utilis, Boyle.
Spiraea canescens, Don.
 " *callosa*, Thunb.
 " *Kamtschatica*, Pall. ?
 " *sorbifolia*, L.
 " *Aruncus*, L.
Rubus acuminatus, Sm.
 " *paniculatus*, Sm.
 " *parvifolius*, L.
 " *concolor*, Wall.
 " *niveus*, Wall.
 " *fruticosus*, L.
 " *flavus*, Ham.
 " *lasiocarpus*, Sm.
 " *rosifolius*, Sm.
 " *biflorus*, Ham.
 " *asper*, Don.
Geum elatum, Wall.
Fragaria indica, Andr.
 " *vesca*, L.
Agrimonia Eupatorium, Linn.
Potentilla fruticosa, L.
 " *microphylla*, Don.
 " *multifolia*, L.
 " *Leschenaultiana*, Sm.
 " *fulgens*, Wall.
 " *nepalensis*, Hook.
 " *atrosanguinea*, Lodd.
 " *supina*, L.
 " (*Sibbaldia*) *procumbens*, L.
Rosa Lyellii, Lindl.
 " *moschata*, Mill.
 " *Brunonii*, Lind.
 " *macrophylla*, Lind.
 " *Webbianæ*, Wall.
 " *sericea*, Lind.
Pyrus Aria, Ehrh.
 " *baccata*, L., Gwála Mehal.
 " *variolosa*, Wall, Mehal.
 " *communis*, L., Nashpáti, Pear, *cultivated*.
 " *Malus* L., Seo, Apple, *cultivated*.
 " *ursina*, Wall.
 " *foliolosa*, Wall.
 " *Cydonia*, L., *introduced*.
Crataegus Pyracantha, Pers.
Cotoneaster acuminata, Lindl.
 " *microphylla*, Wall.
 " *bacillaris*, Wall.
 " *nummularia*, F. et M.
Natural Order 46.—Saxifraga.
Astilbe rivularis, Don.
Saxifraga ligulata, Wall.
 " *Stracheyi*, Hf. & T.
 " *imbricata*, Boyle.

Natural Order 46.—Saxifraga—(concl.)

- Saxifraga* *ramulosa*, Wall.
 " *granulata*, L.
 " *odontophylla*, Willd.
 " *micrantha*, Edgw.
 " *pallida*, Wall.
 " *flagellaria*, Willd.
 " *Brunonis*, Wall.
 " *filiculis*, Wall.
 " *brachypoda*, Don.
 " *diversifolia*, Wall.
 " *Jacquemontiana*, Dene.
Chrysosplenium nepalense, Dene.
Parnassia *Wightiana*, Wall.
 " *nubicola*, Wall.
Hydrangea altissima, Wall.
Deutzia corymbosa, R. Br.
 " *staminea*, R. Br.
Philadelphus coronarius, L.
Itea nutans, Royle.
Ribes *Grossularia*, L.
 " *glaciale*, Wall.
 " *rubrum*, L.
 " *nigrum*, L.

Natural Order 47.—Crassulaceæ.

- Tillsea pentandra*, Royle.
Crassula indica, Dene.
Byrophyllum calycinum Salisb., in *gardens*.
Kalanchoe varians, Haw.
Cotyledon Oreades, Benth. et Hf.
Sedum cronulatum, Hf. & T.
 " *quadrisidum*, Pall.
 " *Himalense*, Don.
 " *crassipes*, Wall.
 " *elongatum*, Wall.
 " *linearifolium*, Royle.
 " *pauciflorum*, Edgw.
 " *trifidum*, Wall.
 " *rosulatum*, Edgw.
 " *adenotrichum*, Wall.
 " *Ewersii*, Ledeb.
 " *multicaule*, Wall.
 " *Rhodiola*, DC.

Natural Order 48.—Droseraceæ.

- Drosera peltata* Var. *lunata*, Ham.

Natural Order 49.—Haloragaceæ.

- Myriophyllum spicatum*, L.

Natural Order 50.—Combretaceæ.

- Terminalia bellerica*, Roxb. Baharia.
 " *Chebala*, Retz. Har.
 " *citrina*, Roxb. Harina.
 " *Arjuna*, Bedd. Arjan.
 " *tomentosa*, Roxb. Saj.
Anogeisus latifolia, Wall.
Combretum decandrum, Roxb.

Natural Order 51.—Myrsinæ.

- Eugenia Jambolana*, Lamk., Jéman.
 " *caryophylli folia*, Lamk.
Careya arborea, Roxb.
Barringtonia acutangula, Gærtn.

Natural Order 52.—Melastomaceæ.

- Osbeckia stellata*, Don.
 " *angustifolia*, Don.

Natural Order 53.—Lythraceæ.

- Ammannia rotundifolia*, Ham.
 " *vesicatoria*, Roxb.
Woodfordia floribunda, Salisb.
Lawsonia alba Lamk., Mehul. *gardenia*.
Lagerstrœmia parviflora, Roxb.
Punica Granatum, L., *wild and cultivated*.

Natural Order 54.—Onagraceæ.

- Epilobium tomentosum*, Boiss.
 " *roseum*, L.
Oenothera rosea, Sims., } *naturalized*.
 " *sinuata* L., }
 " *grandiflora*, }
Circœa Lutetiana, L.
 " *alpina*, L.

Natural Order 55.—Samydaceæ.

- Casearia tomentosa*, Roxb.

Natural Order 56.—Cucurbitaceæ.

- Trichosanthes cucumerina*, L.
 " *palmata*, Roxb.
Luffa cylindrica, Naud.
Cucumis Melo, L., *cultivated*.
 " *sativus*, L.
 " *Hardwickii*, Royle.
Citrullus vulgaris, Schr.
Bryonia laciniosa, L.
Mukia scabrelin, Arn.
Zehneria umbellata, Thw.

Natural Order 57.—Begoniaceæ.

- Begonia picta*, Wall.
 " *amazona*, Wall.

Natural Order 58.—Datiscaceæ.

- Datisca cannabina*, L.

Natural Order 59.—Ficoides.

- Trianthema pentandra*, L.
 " *crystallina*, Vahl.
Mollis hirta, Thunb.
 " *pentaphylla*, L.
 " *cerviana*, Ser.

Natural Order 60.—Umbellifera.

- Hydrocotyle asiatica*, L.
Sanicula europæa, L.
Bupleurum tenac. Ham.
 longicaule, Wall.
 Candollei, Wall.
 falcatum, L.
 marginatum, Wall.
Apium graveolens, L., *introduced*.
Carum Copticum, Benth. et Hf. Ajwain,
 cultivated.
Acronema tenerum, Edgw.
Pimpinella diversifolia, DC.
Cherophyllum villosum, Wall.
Oenanthe stolonifera, Wall.
Selinum Candollei, Bth. et Hf.
 Brunonis, DC.
 Govanianum, DC.
 angelicoide, DC.
Heracleum candicans, Wall.
 nepalense, Don.
Peucedanum graveolens, L. Sawa.
Coriandrum sativum, L., *cultivated*.
Daucus Carota, L., *cultivated*.
Torilis Anthriscus, Gmel.

Natural Order 61.—Araliaceæ.

- Aralia cachemirica*, Dene.
Heteropanax fragrans, Seem.
Hedera Helix, L.
Heptapleurum venulosum, Seem.

Natural Order 62.—Cornaceæ.

- Marlea begoniifolia*, Roxb.
Cornus macrophylla, Wall.
 oblonga, Wall.
 capitata, Wall.
Alangium Lamackii, Thw.

Natural Order 63.—Caprifoliaceæ.

- Abelia triflora*, R. Br.
Lycosteria formosa, Wall.
Lonicera Myrtillus, Hf. & T.
 parvifolia, Edgw.
 purpureascens, Hf. & T.
 hypoleuca, Dene.
 orientalis, Lamk.
 alpigena, L.
 heterophylla, Dene.
 angustifolia, Wall.
 spinosa, Jacquem.
 quinquelocularis, Hardw.
Viburnum cotinifolium, Don.
 stellulatum, Wall.
 integerrimum, Wall.
 nervosum, Don.
 coriaceum, Bl.

Natural Order 64.—Rubiaceæ.

- Adina cordifolia*, Benth. et Hf.
Stephegyne parvifolia, Benth. et Hf.
Hymenodictyon excoleum, Wall.

Natural Order 64.—Linbiaceæ—(concl.)

- Wendlandia exserta*, DC.
 puberula, DC.
Argostemma sarmentosum, Wall.
 verticillatum, Wall.
Hedyotis Burmanniana, R. Br.
 aspera, Heyne.
 brachypoda, DC.
 gracilis, Wall.
Randia dumetorum, Lamk.
 tetrasperma, Roxb.
 uliginosa, DC.
Knoxia corymbosa, L.
Pavetta tomentosa, Sm.
Gardenia turgida, Roxb.
Coffea bengalensis, Roxb.
Hamiltonia suaveolens, Roxb.
Leptodermis lanceolata, Wall.
Spermacoce articularis, L.
Galium asperifolium, Wall.
 acutum, Edgw.
 rotundifolium, L.
 sepylloides, Royle.
 confertum, Royle.
 aparinæ, L.
Rubia cordifolia, L.
Asperula cynanchica, L.

Natural Order 65.—Valerianaceæ.

- Valeriana Wallichii*, DC.
 Hardwickii, Wall.

Natural Order 66.—Dipsacæ.

- Morina longifolia*, Wall.
 breviflora, Edgw.
Dipsacus inermis, Wall.

Natural Order 67.—Compositæ.

- Vernonia anthelmintica*, Willd.
 cinerea, Less.
Elephantopus scaber, L.
Adenostemma viscosum, Forst.
 var. elatum, Don.
 var. laefolium, Don.
Eupatorium longicaule, Wall.
 Wallichii, DC.
Solidago Virga-aurea, L.
Cyathocline lyrata, Cass.
Dichrocephala gracilis, DC.
Grangea madraspatana, Poir.
Myriactis nepalensis, Less.
 Wallichii, Less.
Aster alpinus, L.
 mollisculus, Benth.
 juncæus, Benth.
 Thomsoni, C. B. Clarke.
 (= *Calimeris flexuosa*, Royle.)
 peduncularis, Wall.
Erigeron acris, L.
 bollioides, Benth.
Conyza veronifolia, Wall.

Natural Order 67.—Composite (contd.)

- Conyza absinthifolia*, DC.
 " *viscidula*, Wall.
Blumea lacera, DC.
 " *runcinata*, DC.
 " *bieracifolia*, DC.
Sphaeranthus lirtus, Willd.
Leontopodium alpinum, Cass.
Anaphalis cinnamomea, Benth.
 " *triplinervis*, Sims.
 " *nubigena*, DC.
 " *nubigena*, var. *polycephala*,
 C. B. Clarke.
 " *chionantha*, DC.
 " *araneosa*, DC.
Gnaphalium multiceps, Wall.
 " *indicum*, L.
Cæsulia axillaris, Roxb.
Inula vestita, Wall.
 " *nervosa*, Wall.
 " *barbata*, Wall.
 " *Cappa*, DC.
Carpesium cernuum, L.
 " *abrotanoides*, L.
Siegesbeckia orientalis, L.
Eclipta erecta, L.
Bidens tripartita, L.
Allardia tridaetylites, Hf. & T.
Cotula anthemoides, L.
Tanaetum thibeticum, Hf. & T.
 " *longifolium*, Wall.
Artemisia vestita, Wall.
 " *vulgaris*, L.
Senecio densiflorus, Wall.
 " *graciliflorus*, DC.
 " *laciniosus*, Wall.
 " *coronopifolius*, Desf.
 " *alatus*, Wall.
 " *rufinervis*, DC.
 " *Kunthianus*, Wall.
 " *Candolleanus*, Wall.
Echinops niveus, Wall.
 " *echinatus*, Roxb.
Arctium Lappa, Willd.
Saussurea obvallata, Wall.
 " *taraxacifolia*, Wall.
 " *Kunthiana*, Wall.
 " *depressa*, Wall.
 " *graminifolia*, Wall.
 " *Candolleana*, Wall.
 " *carthamoides*, Benth.
 " *albescens*, Hf. & T.
 " *candicans*, DC.
 " *Roylei*, DC.
Jurinea macrocephala, Benth.
Serratula pallida, DC.
Tricholepis elongata, DC.
Leucomeris spectabilis, DC.
Ainalisa pteropoda, DC.
 " *aptera*, DC.
Gerbera lanuginosa, Benth. et Hf.
Berniera nepalensis, DC.
Tragopogon janczumi, Wall.
Youngia lyrata, Cass.

Natural Order 67.—Composite—(concluded).

- Lactuca graciliflora*, Wall.
 " *Brunoniana*, Wall.
 " *hastata*, DC.
 " *macrantha*, Benth. et Hf.
 " *longifolia*, DC.
 " *auriculata*, DC.
 " *sagittata*, Hf. & T.
 " *Tatarica*, Benth.
Prenanthes hispidula, DC.
Taraxacum Dens-leonis, Desf.
Crepis foetida, L.
Mulgedium Tataricum, DC.
Microhynchus sarmentosa, DC.
Gynura nepalensis, DC.
Emilia sonchifolia, DC.
Doronicum Roylei, DC.

Natural Order 68.—Campanulacea.

- Cyananthus integer*, Wall.
 " *linifolius*, Wall.
Wahlenbergia agrestis, A. DC.
Campanula latifolia, L.
 " *sylvatica*, Wall.
 " *canescens*, Wall.
 " *colorata*, Wall.
Lobelia trigona, Roxb.
 " *pyramidalis*, Wall.

Natural Order 69.—Ericacea.

- Pernettya repens*, Zoll.
Andromeda ovalifolia, Wall.
Rhododendron barbatum, Wall.
 " *campanulatum*, Don.
 " *arborescens*, Sm.
 " *lepidotum*, Wall.
 " *anthopogon*, Don.
Cassiope fastigiata, Don.

Natural Order 70.—Primulacea.

- Primula floribunda*, Wall.
 " *petiolaris*, Wall.
 " *purpurea*, Royle.
 " *rosea*, Royle.
 " *minutissima*, Wall.
 " *involuta*, Wall.
 " *sibirica*, Jacq.
 " *denticulata*, Wall.
Androsace sarmentosa, Wall.
 " *Jacquemontii*, Duby.
 " *rotundifolia*, Hardw.
 " *incisa*, Wall.
 " *elegans*, Duby.
Lysimachia lobeloides, Wall.
 " *pyramidalis*, Wall.
 " *ovalis*, Wall.
 " *japonica*, Thumb.
Anagallis arvensis, L.
Micropyxis pamila, Duby.
Samolus Valerandi, L.

Natural Order 71.—Myrsinæ.

- Mars indica*, A. DC.
 " *argentea*, Wall.
Embellia robusta, Roxb.
Myrsine semiserrata, Wall.
 " *blfaria*, Wall.
Ardisia humilis, Vahl.
 " *floribunda*, Wall.

Natural Order 72.—Sapotacæ.

- Bassia butyræa*, Roxb.
 " *latifolia*, Roxb., at low elevations : planted.

Natural Order 73.—Ebenacæ.

- Diospyros esculpta*, Ham.
 " *Melanoxylon*, Roxb.
 " *montana*, Roxb.
 " *Embryopteris*, Pers.

Natural Order 74.—Stryacæ.

- Symplocos eratægoides*, Ham.
 " *spicata*, Roxb.
 " *racemosa*, Roxb.
 " *ramosissima*, Wall.

Natural Order 75.—Jasminæ.

- Jaaminum laurifolium*, Roxb.
 " *pubescens*, Willd.
 " *punctatum*, Wall.
 " *arborescens*, Roxb.
 " *latifolium*, Roxb.
 " *dispermum*, Wall.
 " *revolutum*, Sims.
 " *pubigerum*, Don.
 " *officinale*, L.
 " *grandiflorum*, L.
 " *Zambac*, Ait., cultivated.
Nyctanthes Arbor-tristia, L.

Natural Order 76.—Oleacæ.

- Fraxinus floribunda*, Wall.
Ligustrum nepalense, Wall.
 " *compactum*, Hf. & T.
Olea cuspidata, Wall.
 " *glandulifera*, Wall.
Chionanthus macrophylla, Wall.
Syringa emodi, Wall.

Natural Order 77.—Apocynæ.

- Carissa diffusa*, Roxb.
Ophtoxylon serpentinum, Willd.
Tabernaemontana coronaria, R. Br.
Vinea pusilla, Murr.
Vallis dichotoma, Wall.

Natural Order 77.—Apocynæ—(conold).

- Wrightia mollissima*, Wall.
Holarrhena antihysenterica, Wall.
Alstonia scholaris, R. Br.
Nerium odoratum, Solander.
Chonemorpha macrophylla, G. Don.
Ichneocarpus fragrans, Wall.
 " *frutescens*, R. Br.

Natural Order 78.—Asclepiadacæ.

- Cryptolepis elegans*, Wall.
Vincetoxicum caucasicum, Dene.
 " *montanum*, Dene.
Calotropis procera, R. Br.
Diemia extensa, R. Br.
Periploca calophylla, Falc.
Tylophora carnosa, Wall.
Marsdenia Roylei, Wight.
Pergularia odoratissima, L.
Orthanthem viminea, Wight.
Hoya parasitica, Wall.
Ceropegia longifolia, Wall.
 " *lanceolata*, Wight.

Natural Order 79.—Leguminæ.

- Gardneria angustifolia*, Wall.
Buddleia paniculata, Wall.
 " *asiatica*, Lour.

Natural Order 80.—Gentianacæ.

- Exacum tetragonum*, Roxb.
 " *polunculatum*, L.
Canacora decussata, R. et S.
 " *diffusa*, R. Br.
Gentiana tenella, Fries.
 " *pedicellata*, Wall.
 " *argentea*, Royle.
 " *capitata*, Ham.
 " *decussata*, Ham.
 " *marginata*, Griseb.
 " *venusta*, Wall.
 " *depressa*, Don.
 " *Kurroo*, Royle.
Crawfordia fasciculata, Wall.
Ophelia coriata, Don.
 " *livida*, Don.
 " *purpurascens*, Don.
 " *nervosa*, Wall.
 " *pulchella*, Don.
 " *angustifolia*, Don.
 " *Chiraya*, Griseb.
Halenia elliptica, Don.
Sweetia speciosa, Wall.

Natural Order 81.—Bignoniacæ.

- Calosanthus indica*, Bl.
Stereospermum chelonoides, DC.
 " *suaevoleus*, DC.

Natural Order 88.—*Orubanchea*.

Phelipaea indica, G. Don.
Aginetia indica, Roxb.

Natural Order 89.—*Geraneaceae*.

Dilymocarpus anabaternans, Wall.
 " *atomatus*, Wall.
 " *macrophyllus*, Wall.
 " *pedicellatus*, R. Br.
 " *lanuginosus*, Wall.
Chirita bifolia, Don.
Rhynchos glossum obliquum, Bl.
Platystemma violoides, Wall.
Lysionotus ternifolius, Wall.

Natural Order 90.—*Acanthaceae*.

Thunbergia grandiflora, Roxb.
 " *coccinea*, Wall.
Elbermaiera glauca, N. ab. E.
Hygrophila polycarpa, T. Anders.
 " *spinosa*, T. Anders.
Petalidium barlerioides, N. ab. E.
Hemigraphis Pavala, T. Anders.
Strobilanthes auriculatus, N. ab. E.
 " *glutinosus*, T. Anders.
 " *alatus*, N. ab. E.
 " *Wallichii*, N. ab. E.
 " *isophyllus*, T. Anders.
 " *anisophyllus*, T. Anders.
Aechmanthera Wallichii, N. ab. E.
Dactylacanthus nervosus, T. Anders.
Barleria cristata, L.
Lepidagathis cuspidata, N. ab. E.
 " *purpuricaulis*, N. ab. E.
 " *hyalina*, N. ab. E.
 " *fasciculata*, N. ab. E.
Philagacanthus thyrsoflorus, N. ab. E.
Justicia Adhatola, L.
 " *peplodes*, T. Anders.
 " *procumbens*, L.
Rungia pectinata, N. ab. E.
 " *repens*, N. ab. E.
Dicliptera Roxburghiana, N. ab. E.
Peristrophe bicalyculata, N. ab. E.
 " *speciosa*, N. ab. E.

Natural Order 91.—*Verbenaceae*.

Verbena officinalis, L.
Lippia nodiflora, Rich.
Lantana alba, Mill.
Premna viburnoides, Wall.
 " *lanceolata*, Roxb.
 " *barbata*, Wall.
 " *herbacea*, Roxb.
Callicarpa arborea, Roxb.
 " *macrophylla*, Vahl.
Clerodendron serratum, Spr.
 " *infortunatum*, L.
 " *Siphonanthus*, R. Br.

**Natural Order 91.—*Verbenaceae*.
—(continued).**

Caryopteris Wallichiana, Schauer.
Gmelina arborea, Roxb.
Vitex Negundo, L.
Holuskiodia sanguinea, Retz.

Natural Order 92.—*Labiatae*.

Ocimum canum, L.
 " *sacrum*, L., *naturalized*.
Orthosiphon rubicundus, Benth.
Plectranthus scrophularioides, Wall.
 " *striatus*, Benth.
 " *Cretia*, Don.
 " *ternifolius*, Don.
 " *cordifolius*, Don.
Colinus barbatus, Benth.
Pogostemon pleuranthoides, Desf.
Dysophylla cruciata, Benth.
 " *pumila*, Benth.
Colobrookia oppositifolia, Sm.
Elsholtzia polystachya, Benth.
 " *flava*, Benth.
 " *incisa*, Benth.
 " *eristachya*, Benth.
 " *strobilifera*, Benth.
Mentha arvensis, L.
 " *incana*, Willd., var. *Royleana*, Benth.
Origanum noronae, Don.
Melissa calamintha, L.
Miconeria biflora, Benth.
Hedeoma nepalensis, Benth.
Salvia glutinosa, L.
 " *Moorecroftiana*, Wall.
 " *lunata*, Roxb.
 " *plebeja*, R. Br.
Nepteta spicata, Benth.
 " *distans*, Royle.
 " *ciliaris*, Royle.
 " *ruderalis*, Ham.
 " *leucophylla*, Benth.
 " *Goumiana*, Benth.
Lallemantia Royleana, Benth.
Amsoules ovata, R. Br.
Prunella vulgaris, L.
Scutellaria grossa, Benth.
 " *repens*, Ham.
 " *linearis*, Benth.
 " *scandens*, Don.
Carmotome versicolor, Benth.
Leonurus Royleanus, Benth.
 " *pubescentia*, Benth.
 " *cardiatus*, L.
Lamium amplexicaule, L.
 " *petiolatum*, Royle.
Colubonium coccinea, Wall.
Stachys scircea, Wall.
 " *splendens*, Wall.
Roylea elegans, Wall.
Leucas urticifolia, R. Br.
 " *lunata*, Benth.
 " *molissima*, Wall.

Natural Order 92.—Labiata
—(concluded).

- Leucas diffusa*, Benth.
" *hymopifolia*, Benth.
" *cephalotes*, Spr.
Leonotis nepetifolia, R. Br.
Phlomis lamiiifolia, Royle.
Bremontachys superba, Royle.
Ajuga remota, Benth.
" *bracteosa*, Wall.
Teucrium Royleanum, Benth.
" *quadrifarium*, Ham.

Natural Order 93.—Plantagineæ.

- Plantago major*, L.
" *Isinghula*, Roxb.

Natural Order 94.—Chenopodiaceæ.

- Chenopodium album*, L.
" *Botrys*, L.

Natural Order 95.—Basellaceæ.

- Basella rubra*, L. *cultivated and naturalized.*

Natural Order 96.—Amarantaceæ.

- Deeringia celosioides*, Moq.
Rodetia Amherstiana, Moq.
Celosia argentea, Moq.
Amaranthus cernuus, L. *cultivated and naturalized.*
Amaranthus paniculatus, Moq.
" *Gaugeticus*, L.
" *trumentaceus*, Roxb. *Ramdana*, *Anardana*, *cultivated and naturalized.*
" *spinosus*, L.
Aerva scandens, Wall.
" *lanata*, Juss.
Achyranthes aspera, L.
Digera arvensis, Forsk.
Pupalia lappacea, DC.
Cyathula prostrata, Bl.
Alternanthera sessilis, R. Br.

Natural Order 97.—Nyctaginæ.

- Boerhaavia diffusa*, L.
Mimbilis Jalapa, L., *introduced and naturalized.*

Natural Order 98.—Phytolaccææ.

- Piceunia latbena*, Moq.

Natural Order 99.—Polygonaceæ.

- Rumex Wallichii*, Meissn.
" *nepalensis*, Spr.
" *hastatus*, Don.

Natural Order 99.—Polygonaceæ.
—(concluded).

- Rumex vesicarius*, L.
Koenigia islandica, L.
Rheum emodi, Wall.
" *himalense*, Royle.
Polygonum Roxburghii, Meissn.
" *plebeium*, R. Br.
" *herniarioides*, Del.
" *recumbens*, Royle.
" *aviculare*, L.
" *barbatum*, L.
" *amphibium*, L.
" *laphifolium*, L.
" *sphaerostachyum*, Meissn.
" *amplexicaule*, Don.
" *vaccinaefolium*, Wall.
" *affine*, Don.
" *nepalense*, Meissn.
" *sphaerocephalum*, Don.
" *capitatum*, Ham.
" *chinense*, L.
" *pterocarpum*, Wall.
" *rumicifolium*, Royle.
Fagopyrum esculentum, Moench.
" *cymosum*, Meissn.
" *tataricum*, Gaertn.

Natural Order 100.—Laurinæ.

- Cinnamomum Tamala*, N. ab E., *var. albiflorum*, Tejpat.
Phoebe lanceolata, N. ab E.
" *pallida*, N. ab E.
Machilus odoratissimus, N. ab E.
Tetranthem Roxburghii, N. ab E.
" *monopetala*, Roxb.
Litsea consimilis, N. ab E.
" *lanuginosa*, N. ab E.

Natural Order 101.—Thymelacææ.

- Daphne papyracea*, Wall.
Wikstroemia virgata, Meissn.

Natural Order 102.—Loranthacææ.

- Loranthus ligustrinus*, Wall.
" *pulvernulentus*, Wall.
" *vestitus*, Wall.
" *longiflorus*, DC.v.
" *cordifolius*, Wall.
Viscum album, L.
" *articulatum*, Burm.
" *attenuatum*, D.C.

Natural Order 103.—Elaeagnææ.

- Hippophaë salicifolia*, Don.
Elaeagnus umbellata, Thunb.
" *latifolia*, L.

Natural Order 104.—Santalacææ.

- Osyris arborea*, Wall.

*Natural Order 105.—Aristolochiacea.**Aristolochia bracteata*, Retz.*Natural Order 106.—Euphorbiacea.*

- Phyllanthus nepalensis*, Müll.
 " *parrifolius*, Wall.
 " *velutinus*, Müll.
 " *Miruri*, L.
 " *Embllica*, L.
Antidesma diandrum, Thuc.
Securinega obovata, Müll.
 " *Leucopyrus*, Müll.
Putranjiva Roxburghii, Wall.
Andrachne cordifolia, Müll.
Bischofia javanica, Bl.
Briedelia montana, Willd.
 " *stipularia*, Bl.
Croton oblongifolius, Roxb., *introduced*.
Trewia nudiflora, Willd.
Mallotus philippinensis, Müll. (Rottlera
tinctoria, Roxb.).
Ricinus communis, L., *cultivated*.
Honcucya riparia, Lour.
Baliospermum polyandrum, Wight.
Excoecaria insignis, Müll.
Euphorbia indica, Lamk.
 " *pilulifera*, L.
 " *nerifolia*, L.
 " *Sikkimensis*, Boiss.
 " *longifolia*, Don.
 " *Helioscopia*, L.
 " *nepalensis*, Boiss.
 " *Nivulia*, Ham.
 " *Koylenna*, Boiss.
Sarcococca pruniformis, Lindl.

Natural Order 107.—Cupulifera.

- Quercus semecarpifolia*, Sm.
 " *hex*, L.
 " *dilatata*, Lindl.
 " *lauguina*, Don.
 " *incana*, Roxb.
 " *annulata*, Sm.

*Natural Order 108.—Myricacea.**Myrica sapida*, Wall.*Natural Order 109.—Betulacea.*

- Betula Bhojpattri*, Wall.
 " *acuminata*, Wall.
 " *nitida*, Don.
Alnus nepalensis, Don.

Natural Order 110.—Corylacea.

- Carpinus viminea*, Wall.
 " *faginea*, Lindl.
Corylus Columna, Linu.

Natural Order 111.—Juglandea.

- Juglans regia*, L.
Engelhardtia Colebrookiana, Lindl.

Natural Order 112.—Salicina.

- Salix tetrasperma*, Roxb.
 " *babylonica*, L., *introduced*.
 " *elegans*, Wall.
 " *daphnoides*, L.
 " *strobilaris*, Anders.
 " *Lindleyana*, Wall.
Populus ciliata, Wall.

Natural Order 113.—Urticacea.

- Ulmus integrifolia*, Roxb.
 " *erosa*, Roth.
Celtis caucasica, Willd.
 " *australis*, L.
Sponia politoria, Planch.
 " *orientalis*, Planch.
Cannabis sativa, L.
Urtica parviflora, Roxb.
Girardinia heterophylla, Dens.
Pilea scripta, Wedd.
Lecanthus peduncularis, Wedd.
Elatostemma sessile, Forst.
 " *diversifolium*, Wedd.
Stracheyanum, Wedd.
Bomheria rugulosa, Wedd.
 " *macrophylla*, Don.
 " *platyphylla*, Don.
Pouzolsia indica, Gaud.
 " *vinen*, Wedd.
Debregeasia longifolia, Wedd.
 " *bicolor*, Wedd.
Memorialis pentandra, Wedd.
 " *hirta*, Wedd.
Villebrunea frutescens, Bl.
Maoutia Puya, Wedd.
Cudrania javanensis, Trec.
Streblus asper, Lour.
Morus atropurpurea, Roxb., *introduced*.
 " *laevigata*, Wall.
 " *indica*, L.
 " *alba*, Willd., *cultivated*.
Ficus bengalensis, L., Bar.
 " *infectoria*, Willd.
 " *religiosa*, L., Pipal.
 " *cordifolia*, Roxb.
 " *nemorialis*, Wall.
 " *caudata*, Wall.
 " *scandens*, Roxb.
 " *foveolata*, Wall.
 " *tuberculata*, Wall.
 " *Roxburghii*, Wall.

Natural Order 113.—Urticaceæ.
—(concluded).

- Ficus hispida*, L. f.
" *Cunia*, Ham.
" *glomerata*, Willd., Oular.
" *Carica*, L. *introduced*.
" *virgata*, Roxb.

Natural Order 114.—Piperaceæ.

- Peperomia reflexa*, Dietr.

Natural Order 115.—Gustaceæ.

- Ephedra vulgaris*, Rich.

Natural Order 116.—Coniferae.

- Pinus longifolia*, Roxb.
" *Gerardiana*, Wall.
" *excelsa*, Wall.
Abies Wobbiiana, Lindl.
" *Smithiana*, Forbes.
Cedrus Deodara, Loud.
Cupressus torulosa, Don.
Juniperus communis, L.
" *excelsa*, Bieb.
" *recurva*, Ham.
Taxus baccata, L.

Natural Order 117.—Palmaeæ.

- Phoenix aculeata*, Roxb.
" *sylvestris*, Roxb.
Chamaerops Martiana, Wall.
Calamus Royleanus, Griff.
Wallichia densiflora, Mart.

Natural Order 118.—Aroideæ.

- Arisaema Jacquemontii*, Bl.
" *utile*, Hf. & T.
" *costatum*, Mart.
" *speciosum*, Mart.
" *curvatum*, Kunth.
Saurumatum guttatum, Schott.
Remusatia vivipara, Schott.
Anomorphallus campanulatus, Bl.
Colocasia, species.
Scindapsus officinalis, Schott.
" *decursiva*, Schott.
Typhonium, species.
Pothos scandens, L.
Acorus calamus, L.

Natural Order 119.—Lemnaceæ

- Lemna minor*, L.
" *polyrrhiza*, L.

Natural Order 120.—Typhaceæ.

- Typha elephantina*, Roxb.

Natural Order 121.—Juncaginaceæ.

- Potamogeton pectinatus*, L.
Aponogeton monostachyus, Roxb.

Natural Order 122.—Seltamineæ.

- Globoia Orixensis*, Roxb.
Zingiber capitatum, Roxb.
" *caesium*, Roxb.
" *ligulatum*, Roxb.
Curcuma angustifolia, Roxb.
" *Zerumbet*, Roxb.
Roscoeia elata, Sm.
" *alpina*, Roxb.
" *spicata*, Sm.
" *purpurea*, Sm.
Amomum subulatum, Roxb.
Hedycheilum coronarium, L., *in gardens*.
" *spicata*, Sm.
" *coccineum*, Ham.
Costus speciosus, L.
Musa paradisiaca, L. *plantain, cultivated*.
" *nepalensis*, Wall.

Natural Order 123.—Orchidaceæ.

- Oberonia iridifolia*, Lindl.
Dienia cylindrostachya, L.
Liparis obovata, Vahl.
Cœlogyne cristata, Lindl.
" *ocellata*, Lindl.
" *ovata*, Lindl.
Eria convallarioides, Lindl.
" *alba*, Lindl.
" *flava*, Lindl.
Dendrobium erioflorum, Griff.
" *alpestre*, Royle.
Otocilus, species.
Apaturia Smithiana, Lindl.
Eulophia campestris, L.
" *harbacea*, Lindl.
Cyrtopora fusca, Wight.
Yanda cristata, Lindl.
Saccolabium guttatum, Lindl.
" *Gharwalicum*, Reichb. f.
Aerides affine, Wall.
" *odoratum*, Lowe.
" *Hystrix*, Lindl.
" *difforme*, Wall.
Calanthe puberula, Lindl.
Orchis latifolia, L.
Gymnadenia cucullata, Rich.
" *spathulata*, Lindl.
Aceras angustifolia, Lindl.
Platanthera orchidis, Lindl.
" *clavigera*, Lindl.
" *obcordata*, Lindl.
" *fusca*, Lindl.
" *candida*, Lindl.
" *acuminata*, Lindl.

Natural Order 123.—Orchidaceae
—(concluded).

- Gamoplexis orobanchoides*, Falc.
Epipactis latifolia, Sw.
 " *maurustachya*, Lindl.
Spiranthes australis, R. Br.
Habenaria pectinata, Lindl.
 " *intermedia*, Don.
 " *marginata*, Lindl.
 " *longifolia*, Lindl.
Peristylus Goodyeroides Lindl.
Hermannium Monorchis, Sw.
 " *speciosa*.
Satyrium nepalense, Don.
Monochilus flavus, Wall.
Tripleura pallida, Lindl.
Goodyera marginata, Lindl.
Cypripedium cordigerum, Don.

Natural Order 124.—Iridaceae.

- Iris sulcata*, Wall.
 " *decora*, Wall.
 " *Ramsonensis*, Wall.
 " *nepalensis*, Wall.

Natural Order 125.—Amaryllidaceae.

- Crinum toxicarium*, Roxb.
 " *amcnum*, Roxb.
Ourculigo recurvata, Roxb.
Hypoxis aurea, Lour.
 " *orchoides*, Ks.

Natural Order 126.—Dioscoreae.

- Dioscorea deltoidea*, Wall.
 " *angittata*, Royle.
 " *pentaphylla*, L.

Natural Order 127.—Liliaceae.

- Tulipa stelleri*, Hook.
Gagea thesioides, Fisch.
 " *elegans*, Wall.
Iphigenia indica, Kth.
Lloydia serotina, Besqhb.
Fritillaria cirrrosa, Don.
 " *verticillata*, Royle.
Lilium giganteum, Wall.
 " *Wallichianum*, R. et S.
 " *nepalense*, Wall.
 " *Thomsonianum*, Hf.
 " *roseum*, L.
Gloriosa superba, L.
Allium Victoriale, L.
 " *Wallichii*, Kth.
 " *tuberosum*, Roxb.
 " *Govanianum*, Wall.
 " *consanguineum*, Kth.
 " *rubens*, Schrad.

Natural Order 127.—Liliaceae.—(concl.)

- Urginea indica*, Kunth.
Asphodelus clavatus, Roxb.
Asparagus racemosus, Willd.
 " *filicinus*, Ham.
Polygonatum multiflorum, All.
 " *sibiricum*, Rod.
 " *verticillatum*, All.
Flüggen intermedia, Sch.
 " *Wallichiana*, Kth.
Smilax maculata, Roxb.
 " *elegans*, Wall.
 " *macrophylla*, Roxb.
 " *ovalifolia*, Roxb.
Paris polyphylla, Sm.
Trillidium Govanianum, Kth.
Disporum Pitatum, Don.
 " *calcaratum*, Don.

Natural Order 128.—Alismaceae.

- Alisma Plantago*, L.
 " *reniforme*, Don.
Sagittaria angittifolia, L.

Natural Order 129.—Pontederaceae.

- Monochoria hastata*, Presl.

Natural Order 130.—Commelynacae.

- Cyanotis axillaria*, R. et S.
 " *barbata*, Don.
Commelyna salicifolia, Roxb.
 " *bengalensis*, L.
 " *obliqua*, Ham.
Ancilema herbaceum, Wall.

Natural Order 131.—Juncaceae.

- Juncus concinnus*, Don.
 " *leucomelos*, Royle.
 " *cantharus*, L.
 " *bufonius*, L.
 " *effusus*, L.
 " *leucanthus*, Don.

Natural Order 132.—Rustiacae.

- Eriocaulon quinquangulare*, L.
 " *trilobum*, Ham.
 " *sexangulare*, L.

Natural Order 133.—Cyperaceae.

- Carex rigida*, Good.
 " *Thomsoni*, Boott.
 " *mitis*, Boott.
 " *nubigena*, Don.
 " *foliosa*, Don.

Natural Order 133.—Cyperaceæ—
(concluded.)

- Carex longipes*, Don.
 „ *Moorecroftii*, Falc.
 „ *alopecuroides*, Don.
 „ *Wallichiana*, Presc.
 „ *setigera*, Don.
 „ *eruenta*, Nees.
 „ *uncinoides*, Boott.
 „ *vesiculosa*, Boott.
 „ *filicina*, Nees.
 „ *conduplicata*, Nees.
 „ *bengalensis*, Roxb.
Rhynchospora aurea, Nees.
 „ *Wallichiana*, Nees.
Fimbristylis quinquangularis, Nees.
 „ *diehotoma*, Vahl.
 „ *æstivalis*, Vahl.
 „ *conduplicata*, Link.
Scirpus junceoides, Roxb.
 „ *atkinsii*, Roth.
 „ *mucronatus*, L.
Abilgaardia monostachya, Vahl.
Eleocharis uniglumis, R. Br.
 „ *palustris*, R. Br.
 „ *ovata*, R. Br.
Isolopis setacea, R. Br.
 „ *trifida*, Nees.
 „ *supina*, R. Br.
 „ *Roylei*, Nees.
Kyllingia monocephala, L.
 „ *brevifolia*, Roth.
 „ *triceps*, Nees.
Eriophorum comosum, Wall.
Cyperus pygmaeus, Vahl.
 „ *alopecuroides*, Roth.
 „ *compressus*, L.
 „ *niveus*, Retz.
 „ *difformis*, L.
 „ *Iris*, L.
 „ *rotundus*, L.
Mariacus panicus, Vahl.

Natural Order 134.—Graminæ.

- Oryza sativa*, L., Rice, *cultivated*.
Zea Mays, L., Makai, Bhuta, Indian
 Corn, *introduced and cultivated*.
Coix Lachryma, L.
 „ *gigantes*, Koen.
Milium effusum, L., *introduced*.
Paspalum pedicellatum, Nees.
 „ *serobiculatum*, L., *cultivated*.
 „ *Royleanum*, Nees.
Coridothloa seminata, Nees.
Panicum sanguinale, L.
 „ *ciliare*, Retz.
 „ *millaceum*, L., *cultivated*.
 „ *maximum*, Jacq.
 „ *ovalifolium*, Poir.
 „ *longipes*, W. & A.
 „ *millare*, Lamk.
 „ *vestitum*, Nees.
 „ *plicatum*, Roxb.

Natural Order 134.—Graminæ (contd.)

- Panicum prostratum*, Lamk.
 „ *repens*, L.
Oplismenus colonus, Kth.
 „ *compositus*, R. & S.
 „ *frumentaceus*, Kunth, *cultivated*.
 „ *stagninus*, Kth.
Setaria glauca, Beauv.
 „ *italica*, Kth., *naturalised; cultivated*.
 „ *verticillata*, Beauv.
Pennisetum spicata, Lamk, Bajra, *cultivated*.
Pennisetum cenchroides, Rich.
 „ *triflorum*, Nees.
Gymnothrix flaccida, Munro.
Arundinella setosa, Trin.
Sporobolus diandra, Beauv.
 „ *elongatus*, R. Br.
Agrostis alba, L.
 „ *ciliata*, Trin.
Milkenbergia sylvatica, Trin.
 „ *geniculata*, Nees.
 „ *viridissima*, Nees.
Polypogon littoralis, L.
Calamagrostis Epigejos, Roth.
Orthoraphium Roylei, Nees.
Piptatherum equiglume, Munro.
Alopecurus pratensis, L.
Phleum arenarium, L.
Phragmites Roxburghii, Kunth.
Chloris barbata, Swartz.
Eleusine indica Gaertn.
 „ *Coracana*, Gaertn., *cultivated*.
 „ *verticillata*, Roxb.
Cynodon Dactylon, Pers., Dab.
Avena fatua, L.
 „ *pratensis*, L.
Poa annua, L.
 „ *pratensis*, L.
 „ *bulbosa*, L.
Eragrostis plumosa, Link.
 „ *pilosa*, L.
 „ *verticillata*, Beauv.
 „ *unioides*, Nees.
 „ *poroides*, Beauv.
 „ *narioides*, Trin.
 „ *diandra*, Roxb.
Melica ciliata, L.
Koeleria cristata, Pers.
 „ *phleoides*, Pers.
Dactylis glomerata, L.
Festuca duriuscula, L.
Bromus squarrosus, L.
 „ *arvensis*, L.
 „ *giganteus*, L.
Arundinaria falcata, Nees.
 „ *Falconeri*, Kx.
Thamnocalamus spathiflorus, Munro.
Bambusa arundinacea, L.
Dendrocalamus strictus, Nees.
Hordeum hexastichon L., Jan, Barley, *cultivated*.

Natural Order 134.—Gramina—(concl'd.)

- Hordium vulgare*, L., *cultivated*.
Elymus sibiricus, L.
Lolium speciosum, Stev.
 " *temulentum*, L.
Triticum vulgare, Vil., Gehán, Wheat,
cultivated.
Mnesithea laevis, Kth.
Rottboellia exaltata, L.
Ophiurus corymbosus, Grtn.
Manisuris granularis, Sw.
Andropogon tryllus, L.
 " *acicularis*, Retz.
 " *involutus*, Steud.
 " *montanus*, Roxb.
 " *ischaemum*, L.
Heteropogon contortus, R. & S.
 " *Roylei*, Nees.
Sorghum vulgare, Pers., Jowar, *cultivated*.
 " *Halepense*, L.
Apocypis Royleanus, Nees.
Batrachium molle, Nees.
 " *lanceifolium*, Trin.
 " *echinatum*, Nees.
Anatherum muricatum, Beauv.
Cymbopogon laniger, Desf.
 " *flexuosus*, Nees.
 " *Martini*, Mauro.
 " *Schoenanthus*, L.
 " *Nardus*, L.
Androscopia gigantea, Brongn.
Anthistiria aru dinneen, Nees.
 " *Anatherum*, Nees.
 " *scandens*, Roxb.
Apluda aristata, L.
 " *geniculata*, Roxb.
Imperata arundinacea, Cyrill.
Saccharum spontaneum, L.
 " *procerum*, Roxb.
 " *officinarium*, L., *Sugarcane*
 cultivated, introduced.
 " *Sara*, Roxb.
 " *filifolium*, Nees.
Erianthus Ravennae, Beauv.
 " *japonicus*, Beauv.

CRYPTOGAMS.

Filices or Ferns.

- Woodsia elongata*, Hk.
 " *lancea*, Hk.
Dieksonia appendiculata, Wall.
 " *scabra*, Wall.
Davallia pulchra, Don.
 " *divaricata*, Bl.
 " *immersa*, Wall.
 " *nodosa*, Hk.
Adiantum lunulatum, Burm.
 " *candatum*, L.
 " *Capillus-Veneris*, L.

Filices or Ferns—(continued.)

- Adiantum venustum*, Don.
Cheilanthes Dalhousie, Hk.
 " *farinosa*, Kaulf.
Onychium auratum, Kaulf.
 " *japonicum*, Kunze.
Pelica calomelanos, Link.
Pteris longifolia, L.
 " *eretica*, L.
 " *quadrisaurita*, Retz.
 " *aquilina*, L.
 " *Wallichiana*, Agdh.
Woodwardia radicans, Sm.
Asplenium alternans, Wall.
 " *varians* Hk. and Gr.
 " *viridie*, Hudb.
 " *Tichomanes*, L.
 " *planicaule*, Wall.
 " *fontanum*, Bernh.
 " *bulbiferum*, Forst.
 " *tennisfolium*, Don.
 " *Filix foemina*, Bernh.
 " *polypodioides*, Mett.
 " *dilatatum*, Wall.
 " *esculentum*, Persl.
 " *Ceterach*, L.
 " *australe*, Brack.
Aspidium auriculatum, Sw.
 " *aculeatum*, Sw.
 " *Thomsoni*, Hk.
 " *aristatum*, Sw.
 " *falcatum*, Sw.
Nephrodium unitum, R. Br.
 " *prolixum*, Bak.
 " *sparsum*, Don.
 " *molle*, Desv.
 " *electarium*, Bak.
 " *polymorphum*, Bak.
 " *Filix-mas*, Rich.
 " *coriaceum*, Don.
 " *odoratum*, Bak.
Oleandra Wallichii, Hk.
 " *neriiformis*, Cav.
Polypodium cruceaeus, Wall.
 " *proliferum*, Presl.
 " *lineatum*, Cotebr.
 " *arvenum*, Wall.
 " *lachnopus*, Wall.
 " *malacodon*, Hk.
 " *trifidum*, Don.
 " *Lingua* Sw.
 " *stigmaeum*, Sw.
 " *fixum*, Bak.
 " *focculosum*, Don.
 " *lineare*, Thunb.
 " *normale*, Don.
 " *membranaceum*, Don.
 " *propinquum*, Wall.
 " *juglandifolium*, Don.
 " *guercifolium*, L.
 " *himalense*, Hook.
 " *leiorhizon*, Wall.
Acrostichum virens, Wall.
Gymnogramme vestita, Hk.

Filices or Ferns—(concl'd.)

- Gymnogramme* *Andersoni*, Bedd.
 " *javanica*, Bl.
 " *lanceolata*, Hk.
 " *involuta*, Don.
Osmunda *Claytoniana*, L.
Schizaea *digitata*, Sw.
Lygodium, *scandens*, Sw.
 " *japonicum*, Sw.
Botrychium *danceifolium*, Wall.
 " *Lunaria*, Sw.
 " *Virginianum*, Sw.

Lycopodiaceae.

- Selaginella* *Jacquemontii*, Spring.
 " *setacea*, Spring.
 " *subulifolia*, Spring.

Marsiliaceae.

- Marsilea* *quadrifolia*, L.

Characeae.

- Chara* *veticillata*, Roxb.

PLANTS OF KUMAON.¹

THE following list gives the names of all plants known to have been found in Kumaon, Garhwál and the Bhábar, three of the districts which are included in the Commissionership of Kumaon. Plants of the plains of India, plants growing in Tibet, and plants only found in the Independent States of Nepál and Tihri, do not appear in this list. The words, "common," "frequent," and "rare" have been employed to denote the comparative occurrence of the various species; but as the European stations of Almora, Naini Tál, and Muníkhet are all situated at a height of between 5,000 and 7,000 feet above the level of the sea, it will be easily understood that plants growing about that level are more likely to meet an observer's eye than plants which grow either at a very high, or at a very low elevation. Plants which are marked "rare" without any locality being given, are those which have been found by former observers, but which have never been seen growing within the limits of the division by Colonel Davidson, Mr. Waterfield or Dr. Watson. In the arrangement of the orders, genera, and species, Hooker's 'Flora of British India,' has been followed as far as published, that is to say, to the end of *Polypetales*. With regard to the other groups the books chiefly consulted have been Brandis' Forest Flora, Madden's Papers in the Journal of the Asiatic Society, Royle's Illustrations of Himalayan Botany, C. B. Clarke's Compositæ and Commelynaceæ, Steudel's Grasses, and Cyperaceæ, Hooker and Baker's Ferns, Mitten's Mosses.

Synonyms have rarely been given except where the name adopted in the list is different from that given by Royle. Royle's Illustrations, published in 1839, is still a standard work on Himalayan Botany, and most of the plants figured

Range of plants.

in it are found in the Commissionership of Kumaon. There are, however, in Royle's book a certain number of plants which are common near Mussooree, where the book was written, but which are not found east of the Alaknanda river, the

¹ This notice and list has been prepared for this volume by Mr. W. Watson, M.D., Deputy Sanitary Commissioner, Almora.

boundary between the Kumaon Commissionership and the Native State of Tihri. As examples of these Mussooree plants not found in Kumaon may be quoted—

Dictamnus albus, *Berberis Lycium*, *Rosa Lyelli*, *Sonerila tenera*, *Gentiana Kurroo*, *Picrorhiza Kurroo*, *Eremosylachys superba*.

On the other hand, the common *phalel* tree of Eastern Kumaon (*Bassia butyracea*) is apparently not found at Mussooree, as it is not mentioned by Royle as occurring there. The same remark applies to the very common and very beautiful *Silang* tree of Kumaon (*Olea acuminata*). This tree, conspicuous from its sweet scented yellow blossoms, is never alluded to by Royle, so that he probably never met with it near Mussooree. It is found in Kumaon at all elevations from 3,000 to 7,000 feet. There are fine trees of it at Debi Dhúra at a height of 6,800 feet, and at Kapkot at a height of 3,400 feet.

The Kumaon Division includes a tract of country varying much in elevation. On the south, the Bhábar strip along the foot of the hills has a mean height of about 1,400 feet, whilst the passes that lead into Tibet have an elevation exceeding 18,000 feet. It might be expected that European forms would be more abundant towards these passes than in the Bhábar, and this, on the whole, is the case. There are, however, some European plants that are only found at low elevations and are never seen above 6,000 feet. The most common of these

are :—*Ranunculus sceleratus*, *Geranium molle*, *Veronica Anagallis*, and *Verbena officinalis*, and it is difficult to assign an adequate reason for this phenomenon. Another remarkable fact connected with the montane flora is the eastern depression of many Himálayan plants. By this term is understood that plants which occur only at high elevations near Simla are found to grow at lower elevations at Mussooree and at still lower heights near Naini Tál. The following plants may be named as affording examples of eastern depression—

- (a) The shrub *Coriaria nepalensis*, which gives its name to the station of Mussooree, is said by Royle not to grow under 5,000 feet at Mussooree. In Kumaon, however, it descends as low as 2,000 feet.

- (b) *Hemiphragma heterophyllum*, one of the *Scrophulariaceæ*, is never found at Simla below 10,000 feet, whilst it occurs in Kumaon at 7,000 feet, and is common there at 8,000 feet.
- (c) The *Tilonj oak*, *Quercus dilatata*, is found at Simla about 7,000 feet, and near Naini Tál it grows abundantly down to 3,000 feet.
- (d) *Dipsacus inermis* is one of the most common plants at Almora (5,500 feet), but is not found towards Simla at elevations below 8,000 feet.
- (e) *Elsholtzia polystachya*, another common Almora plant, is found in Kumaon as low down as 4,000 feet, but near Simla it is not met with under 7,000 feet.
- (f) *Senecio densiflorus* and *S. rufinervis*, with auricled leaves, are common all over Kumaon in November as low as 4,000 feet, but are not found in the neighbourhood of Simla under 7,000 feet.

It does not follow, and is not apparently true, that the higher limit of these plants follows the same law as their lower limit, but of the existence of the law of eastern depression of the lower limit of Himálayan plants, there can be little doubt. There is no example, so far as is known, of the converse, that is to say, of a plant confined to the higher ranges in Kumaon and growing at a low level near Simla.

With regard to certain plants being confined to particular rock formations, some interesting observations were made by Major Madden; thus he found that *Trigonella gracilis*, *Argyrolobium roseum*, and *Argyrolobium flaccidum* were confined to limestone rock. *Thymus serpyllum*, on the contrary, is only found on micaceous rock formations.

One of the most interesting facts for investigation in the Flora of any country is the number of introduced plants which have become wild. Of these, two may be noticed in Kumaon. The *deodár* tree, which is really a native of the Himálaya west of the Alaknanda, but being a sacred tree, it has for many generations been planted round almost every Kumaon temple. There are now magnificent groves at Gangolihát,

Lohaghát, Champláwat, and Bála Jageswar, and there is a tree at Wán in Garhwál which is certainly many hundreds of years old. Still the *deodár* is undoubtedly not indigenous. The other plant is weed and is of comparatively recent introduction. It is the *Erigeron canadense*, known to the natives as the locust weed, (*sallo*) because it is believed to spring from soil where locusts have alighted. This plant is not mentioned by Major Madden, who published his papers in the year 1840, and it has therefore in all probability made its appearance since that date. It is now a most common and a most troublesome weed in cultivated fields.

Many of the Kumaon plants are collected by the villagers and sold to herbalists to be used in medicine. The chief marts are

Indigenous drugs. Haldwáni, Barmdeo, and Ramnagar in Kumaon, and Kotdwára in Garhwál.

The articles most in demand are the fruit of *Terminalia Chebula*, the roots of *Curcuma longa*, *Kyllingia triceps*, *Piper silvaticum*, and the various species of aconite; the bark of *Myrica aspidula* and *Cinnamomum Tamala*; the rind of the fruit of the pomegranate; the powder from *Mulotus philippinensis*, and the various species of *chiretta*, all of which will be noticed hereafter in the chapter on 'Economic Botany.'

Dr. Watson desires to record and acknowledge the assistance that he has received in the preparation of this list from his friends Dr. King, Director of the Royal Botanical Gardens, Calcutta; Colonel Davidson, R.E., and Mr. W. Waterfield of the Bengal Civil Service.

PLANTS OF KUMAON.

DIVISION I.—POLYPETALÆ.

No.	Scientific name.	Vernacular name.	Notes.
ORDER 1.—RANUNCULACEÆ.			
1	<i>Clematis</i> (<i>Cheilopsis</i>) <i>Nepaulensis</i>	December, near streams; rare.
2	" " <i>montana</i> ...	Ghantiālī ...	April, common.
3	" " <i>barbellata</i>	May; rare.
4	" (<i>Flammula</i>) <i>grata</i> ...	Ghantiālī ...	Common.
5	" " <i>Gouriana</i>	Bhābar and valleys; October.
6	" " <i>puberula</i>	Colonel Davidson.
7	" " <i>orientalis</i> [<i>parvifolia</i> Edgeworth.]	Above 10,000 feet.
8	" " <i>nutans</i>	May. Common.
9	" " <i>acuminata</i>	Rare.
10	" " <i>connata</i> [<i>amplexicaulis</i> <i>velutina</i> and <i>gracilis</i> , Edgeworth.] ...	Ghantiālī ...	August, September. Very common.
11	" " <i>Buchananiana</i> , ...	Ghantiālī ...	October. Very common.
12	" " <i>grewiaeflora</i>	Rare.
13	<i>Anemone</i> (<i>Eriocarpus</i>) <i>vitifolia</i>	June and July; common.
14	" (<i>Anemonanthes</i>) <i>obtusiloba</i> , ...	Kakriya ...	May. Very common.
	" [<i>Govaniana</i> and <i>discolor</i>]	Above 6,000 feet.
15	" " <i>rupestris</i>	June. Above 10,000 feet.
16	" " <i>rivularis</i>	Common.
17	" (<i>Omalocarpus</i>) <i>polyanthes</i> ,	Rare, 10,000 feet.
18	" " <i>elongata</i>	" "
19	<i>Thalictrum</i> (<i>Thysocarpus</i>) <i>elegans</i>	Rare. " Badrināth.
20	" " <i>platycarpum</i>	Rare.
21	" " <i>cultratum</i>	"
22	" " <i>Chelidonium</i>	"
23	" " <i>reniforme</i> [<i>neurocarpum</i> , Boyle.]	Common; high elevations, July.
24	" " <i>pauciflorum</i>	Frequent.
25	" (<i>Euthalictrum</i>) <i>Panduratum</i> ,	Common.
26	" " <i>saniculæ forme</i> ,	On trees; autumn.
27	" " <i>Javanicum</i>	"
28	" " <i>foliosum</i> ...	P'englī jārī ...	Very common, summer.
29	" " <i>minus</i> ...	" ...	Frequent.
30	<i>Callianthemum</i> <i>cachemirianum</i>	Above 10,000 feet.
31	<i>Adonis</i> <i>festivalis</i>	"
32	<i>Ranunculus</i> (<i>Batrachium</i>) <i>aquatilis</i>	Bhīm "Tāl.
33	" (<i>Hecatonia</i>) <i>aceleratus</i> ...	Shīm ...	Common up to 4,000 feet.
34	" " <i>hirtellus</i>	High elevations.
35	" " <i>diffusus</i>	Common.
36	" " <i>lætus</i> ...	Dainīya ...	Very common.
37	" (<i>Echinella</i>) <i>flaccidus</i>	Rare.
38	" " <i>arsenalis</i> ...	Ainwa ...	Common at low levels.

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 1.—RANUNCULACEÆ—(concluded).			
39	<i>Oxygraphis glacialis</i>	Above 15,000 feet.
40	" <i>polypetala</i>	" 10,000 "
41	<i>Caltha palustris</i> [Govaniana, Royle],	...	Common.
42	<i>Trollius acaulis</i>	Above 10,000 feet.
43	" <i>pumilus</i>	" 12,000 "
44	<i>Isopyrum grandiflorum</i>	" " "
45	<i>Aquilegia vulgaris</i>	Common. " "
46	<i>Delphinium denudatum</i> [pauciflorum, Royle.]	Mánila ...	"
47	" <i>cæruleum</i>	Above 15,000 feet.
48	" <i>elatum</i>	China hill 8,000 feet.
49	" <i>vestitum</i>	" 8,000 "
50	" <i>Cashmirianum</i>	" 10,000 "
51	" <i>Ajacia</i>	Mánila ...	Naturalized.
52	<i>Aconitum Lycoctonum</i>	Bish ...	Above 8,000 feet.
53	" <i>palmatum</i>	" " "
54	" <i>ferox</i>	" 10,000 "
55	" <i>Napellus</i>	" " "
56	" <i>heterophyllum</i>	Atis ...	" 8,000 "
57	<i>Actæa spicata</i>	Rare.
58	<i>Cimicifuga foetida</i>	"
59	<i>Pæonia emodi</i>	Bhúnýa madin,	Young shoots eaten.
ORDER 2.—DILLENIACEÆ.			
1	<i>Dillenia indica</i>	Rare.
ORDER 3.—MAGNOLIACEÆ.			
1	<i>Michelia Kisopa</i>	Ban champs,	Common: Beripul and Ukhimath.
2	<i>Schizandra grandiflora</i>	Silangti ...	Frequent; Gágar.
3	" <i>propinqua</i>	Sindráyan ...	Common.
ORDER 4.—ANONACEÆ.			
1	<i>Polyalthia Korinti</i> [Gutteria Korinti, Madden.]	...	Swamps. Bhubar.
2	<i>Anona squamosa</i>	Beha ...	Ditto "
3	<i>Millettia velutina</i>	Rare.
4	<i>Saccolpetalum tomentosum</i>	Kangri ...	Kotiwára.
ORDER 5.—MENISPERMACEÆ.			
1	<i>Tinospora cordifolia</i>	Gúcha ...	Common.
2	<i>Cocculus laurifolius</i>	Tilbarapany-i liand kakra.	" about 4,000 feet.
3	" <i>villosa</i>	Rare.
4	<i>Stephania elegans</i>	"
5	" <i>rotunda</i> [Roxburghiana]	Ganjára ...	Common.
6	<i>Clematis Pareira</i> [convolvulacea],	Pári ...	"

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 6.—BERBERIDÆ.			
1	<i>Holboellia latifolia</i>	Gophla ...	May, common, Rênikhet and Binsar.
...	" <i>rur. angustifolia</i>
2	<i>Berberis</i> (<i>Mahonia</i>) <i>nepalensis</i> ...	Gophla ...	Frequent, Thâkil and Binsar.
3	" (<i>Euberberis</i>) <i>vulgaris</i>	High elevations.
4	" " <i>umbellata</i>	Rare.
5	" " <i>aristata</i>	Chotra ...	Common. High elevations.
6	" " <i>Lycium</i>	Rare.
7	" " <i>asiatica</i>	Kilnora ...	Common.
8	" " <i>Wallichiana</i>	China hill, Colonel Davidson.
9	<i>Podophyllum emodi</i>	High elevations.
ORDER 7.—NYMPHACEÆ.			
1	<i>Nelumbium speciosum</i>	Kanwâl ...	Frequent, Buwâlî pokar.
ORDER 8.—PAPAVERACEÆ.			
1	<i>Papaver dubium</i> [<i>glabrum</i> , Royle] ...	Tâkuvia ...	Common.
2	<i>Argemone mexicana</i>	Bhar bhând,	Bhîbar; introduced.
3	<i>Meconopsis aculeata</i>	Kânda ...	Above 10,000 feet
4	" <i>robusta</i>	" ...	" " "
5	" <i>nipalensis</i>	" ...	" " "
6	<i>Stylophorum lactucoides</i>
ORDER 9.—FUMARIACEÆ.			
1	<i>Dicentra Roylei</i>	Rare.
2	" <i>scandens</i>	Common.
3	<i>Corydalis rutæfolia</i> [<i>diphylla</i>]	"
4	" <i>cachemiriæna</i>	Above 9,000 feet.
5	" <i>crithmifolia</i>	Rare.
6	" <i>elegans</i>	Above 10,000 feet.
7	" <i>Gossuliana</i>	" 8,000 "
8	" <i>ramosa</i>	" 12,000 "
9	" <i>sibirica</i> [<i>filiformis</i>]	Common.
10	" <i>cornuta</i>	Blutkes ...	Very common,
11	" <i>chamrophylla</i>	Common, Binsar.
12	" <i>nifolia</i>	Above 10,000 feet.
13	" <i>fiabellata</i>	" 9,000 "
14	<i>Fumaria parvifolia</i> [<i>Vaillantii</i>] ...	Khairawa ...	Very common.
ORDER 10.—CRUCIFERÆ.			
TRIBE ARABIDÆ.			
1	<i>Nasturtium officinale</i>	Piriyâ hâlim,	Cultivated.
2	" <i>palustre</i>	Common.
3	" <i>montanum</i>	"
4	<i>Barbarea vulgaris</i>	"
5	<i>Arabis glabra</i>	Frequent.

DIVISION 1.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 10.—CRUCIFERÆ—(continued).			
TRIBE ARABIDÆÆ—(concluded).			
6	<i>Arabis alpina</i>	Frequent.
7	" <i>amplexicaulis</i>	Itái ghás ...	Common.
8	<i>Cardamine hirsuta</i>
9	" <i>impatiens</i>	Sheliya ...	Very common.
10	" <i>macrophylla</i>	Common.
TRIBE ALYSSINÆÆ.			
11	<i>Alyssum maritimum</i>	Naturalized.
12	" <i>canescens</i>	Above 14,000 feet
13	<i>Barba (Chrysodroma) alpina</i>	" 12,000 "
14	" (<i>Leucodroma</i>) <i>incana</i>	" 10,000 "
15	" " <i>lasiophylla</i>	" 12,000 "
16	" " <i>fladnitzensis</i>	" 13 000 "
TRIBE SISYMBRIÆÆ.			
17	<i>Sisymbrium mollissimum</i>	Above 8,000 feet.
18	" <i>himalaicum</i>	" 10,000 "
19	" <i>Thalianum</i>	Common.
20	" <i>rupestris</i>	Above 8,000 feet.
21	" <i>striatum</i>	Common.
22	" <i>Wallichii</i>	"
23	" <i>sophia</i>	Jauá ghás ...	"
24	" <i>Columnæ</i>	Frequent.
25	" <i>Alliaria</i>	Rare.
26	<i>Eutrema prinnæefolium</i>	"
27	<i>Erysimum hieracifolium</i>	"
TRIBE BRASSICÆÆ.			
28	<i>Brassica nigra</i>	Káli sarson...	Cultivated.
29	" <i>campestris</i>	"
	" <i>var. dichotoma</i>	Jariya ...	"
	" <i>var. glauca</i> , Rox	Rara, píla sarson.	"
	" <i>var. glauca</i> , Royle	Dain toriya...	" up to 11,000 feet.
30	" <i>trilocularis</i>	Weed in fields.
31	" <i>juncea</i>	Rái ...	Cultivated.
32	<i>Eruca sativa</i>	Dúá ...	Common.
TRIBE LEPIDINÆÆ.			
33	<i>Capsella Bursa-pastoris</i>	Very common.
34	<i>Lepidium sativum</i>	Hálin ...	Common.
35	" <i>capitatum</i>	Above 10,000 feet.
36	" <i>ruderales</i>	Common, May and June, Almora.
TRIBE THLASPIDÆÆ.			
37	<i>Megacarpus polyandra</i>	Above 10,000 feet.
38	<i>Thlaspi arvense</i>	Dhapriya ...	Common.
39	" <i>alpestre</i>	Above 7,000 feet.
40	<i>Iberidella Andersoni</i>	" 12,000 "

DIVISION 1.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 10.—CRUCIFERÆ—(concluded).			
TRIBE RAPHANÆ.			
41	<i>Raphanus sativus</i> ...	Muli ...	Cultivated.
ORDER 11.—CAPPARIDACEÆ.			
1	<i>Cleome</i> [<i>Polanisia</i> , Madden] <i>viscosa</i>	Common, Bhābar and valleys.
2	<i>Gynandropsis pentaphylla</i>	Bhābar and valleys.
3	<i>Capparis spinosa</i> [<i>var. leucophylla</i>] ...	Ula kanta ...	Common.
4	" <i>horrida</i> ...	" ...	Bhābar.
5	<i>Mæna arcuaria</i> [<i>Niebulnia oblongifolia</i> , Hoyle, III.]	...	Rare.
ORDER 12.—RESEDACEÆ.			
1	<i>Reseda odorata</i>	Naturalized.
ORDER 13.—VIOLACEÆ.			
1	<i>Viola biflora</i> [reniformis]	Common.
2	" <i>Patrinii</i> [cæspitosa]	"
3	" <i>diffusa</i>	Rare.
4	" <i>serpens</i> [canescens, Wall.]	Common.
5	" <i>kunawarensis</i>	Above 11,000 feet.
ORDER 14.—BIXINEÆ.			
1	<i>Cochlospermum Gossypium</i> ...	Kumbi, kutira, ...	Common, Bhābar.
2	<i>Flacourtia Ramontelii</i> ...	Kantai ...	"
3	" <i>acicularis</i> ...	" ...	"
4	<i>Xylosma longitollum</i> ...	Katari ...	Rare.
ORDER 15.—PITTOSPORACEÆ.			
1	<i>Pittosporum floribundum</i>	Common, river banks.
2	" <i>eriocarpum</i> ...	Gar siling ...	"
ORDER 16.—POLYGALACEÆ.			
1	<i>Polygala triphylla</i> [glaucescens, Wall.]	Common, autumn.
2	" <i>crotolarifolia</i>	Very common.
3	" <i>abyssinica</i>	Rare.
4	" <i>leptocaula</i> [oligophylla, Wallich].	...	Common up to 4,000 feet.
5	" <i>peralcarisefolia</i>	Angust, frequent.
6	" <i>chinensis</i> , [Rothiana, W. and A.]	...	Common, Bhābar.
7	" <i>sibirica</i> [elegans, Wallich],	...	"
ORDER 17.—CARYOPHYLLACEÆ.			
TRIBE SILENÆ.			
1	<i>Gypsophila cerastioides</i>	Common about 10,000 feet.
2	<i>Saponaria vaccaria</i>	Introduced.
3	<i>Silene inflata</i>	Common.

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 17.—CARYOPHYLLACEÆ—(concluded).			
TRIBE SILENEÆ—(concluded).			
4	<i>Silene conoidea</i>	Common.
5	" <i>gallica</i>	Introduced.
6	" <i>Falconeriana</i>	Common.
7	" <i>Stuebelii</i>	Rare.
8	<i>Oenothera bacciferus</i>	"
9	<i>Lychula multicaulis</i>	"
10	" <i>inflata</i> , Pinkworth	Very rare.
11	" <i>indica</i> [fimbriata, Madden],	...	Very common.
TRIBE ALSINEÆ.			
12	<i>Corastium daluricum</i>	Rare.
13	" <i>vulgatum</i> [triviale, Madden].	...	Very common.
14	" <i>Thomsonii</i>	Rare.
15	<i>Stellaria</i> (<i>Schizostegium</i>) <i>crispata</i> [monosperma, Don].	...	Common.
16	" " <i>paniculata</i> [media, Wight.	...	Rare.
17	" (<i>Malachium</i>) <i>aquatica</i>	"
18	" (<i>Stellaria</i>) <i>media</i> [mono- gyna, Don].	...	"
19	" " <i>semivestita</i>	Painkhanda; common.
20	" " <i>Wetbliana</i> , [<i>Leuco- stemma angus- tifolia</i> , Wall].	...	Very common.
21	" " <i>latifolia</i> , [<i>leuco- stemma latifolia</i> , Wall].	...	Rare. Painpáni.
22	" (<i>Larrea</i>) <i>longissima</i> [pa- tens, Don].	...	"
23	" (<i>Adenonema</i>) <i>decumbens</i>	Above 11,000 feet.
24	<i>Arenaria</i> (<i>Alsine</i>) <i>foliosa</i>	Painkhanda; common, June.
25	" (<i>Erenogone</i>) <i>festucoides</i>	Rare.
26	" (<i>Euthalia</i>) <i>serpyllifolia</i>	Common, May & June.
27	" " <i>orbiculata</i>	"
28	" (<i>Lepyrodella</i>) <i>holosteoides</i> ,	"
29	" <i>Benthiana</i>	Rare.
30	<i>Spergula arvensis</i>	Introduced.
31	" <i>pentandra</i> [<i>Arenaria flacci- da</i> , Roxb.]	...	"
TRIBE POLYCARPÆ.			
32	<i>Drymaria cordata</i>	Common.
33	<i>Polycarpa corymbosa</i>	"
ORDER 18.—PORTULACÆ.			
1	<i>Portulaca oleracea</i>	Lániya-kulfa,	Common.

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 19.—TAMARISCINÆ.			
1	<i>Tamarix gallica</i>	Bhábar.
2	" <i>dioica</i>	"
3	<i>Myricaria gormanica</i>	Frequent, above 10,000 feet.
4	" <i>elegans</i>	Frequent, above 11,000 feet.
ORDER 20.—HYPERICINÆ.			
1	<i>Hypericum</i> (<i>Androsæmum</i>) <i>cernuum</i> , ...	Ulus-bena ...	Common. April.
2	" <i>cardifolium</i>	Rare, Kánni.
3	" <i>lysımachioides</i>	Frequent above 8,000 feet.
4	" <i>patulum</i> [uralum, Don].	Common, September.
5	" (<i>Euhypericum</i>) (<i>perforatum</i>).	" "
6	" <i>elodeoides</i>	" August, and September.
7	" <i>nepaulense</i>	" August, and September.
8	" (<i>Brathya</i>) <i>japonicum</i> [<i>pusillum</i>].	Low elevations.
ORDER 21.—TERNSTRÆMIACÆ.			
1	<i>Eurya acuminata</i>	Deura ...	Common
2	<i>Actinidia</i> [<i>Trochostylis</i> , Maiden] <i>callosa</i>	Bhábar and banks of Sarju.
2	<i>Saurauja nepaulensis</i>	Goganá ...	Common.
4	<i>Camellia thea</i>	Chá ...	Introduced.
ORDER 22.—DIPTEROCARPÆ.			
1	<i>Shorea robusta</i>	Sál Sákhu ...	Common up to 4,500 feet, March.
ORDER 23.—MALVACÆ.			
TRIBE MALVÆ.			
1	<i>Malva verticillata</i>	Frequent.
2	" <i>silvestris</i> , [<i>mauritiana</i> , Maiden]. ...	Tithoni ...	Gardens and wild.
3	" <i>rotundifolia</i>	Common.
4	" <i>parviflora</i>	Up to 2,000 feet.
5	<i>Sida rhombifolia</i>	Bhoa Kálaba'l ...	Common.
6	" <i>cordifolia</i>	"
7	<i>Abutilon polyandrum</i>	Bhábar and up to 3,000 feet.
8	" <i>indicum</i>	
TRIBE URENEÆ.			
9	<i>Urena lobata</i>	Sújya Chat-kura.	Common, banks of streams.

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 23.—MALVACEÆ—(continued)			
TRIBE HIBISCÆÆ.			
10	<i>Hibiscus</i> (<i>Furcaria</i>) <i>furcatus</i>	Introduced.
11	" (<i>Lagunea</i>) <i>solandira</i>	Common.
12	" (<i>Abelmoschus</i>) <i>ficulneus</i>	"
13	" <i>pungens</i> ...	Háo Kapása	"
14	" <i>cancellatus</i>	"
15	<i>Thespesia</i> <i>lampas</i> ...	Kapása	"
TRIBE BOMBACÆÆ.			
16	<i>Kydia</i> <i>calycina</i> ...	Patá	Common, September.
17	<i>Bombax</i> <i>malabaricum</i> [<i>Bombax</i> <i>heptaphyllum</i> , Roxb.]	Semal	" Bhábar, up to 4,800 feet.
ORDER 24.—STERCULIACÆÆ.			
1	<i>Sterculia</i> <i>urens</i> ...	Katira	Bhábar.
2	" <i>villosa</i> ...	Udyál	"
3	" (<i>Firmiana</i>) <i>fulgens</i> [Wallichii, Maiden.]	Bodálá	Common.
4	<i>Helicteres</i> <i>isora</i> ...	Maror phal,	Bhábar, May, September.
5	<i>Pterospermum</i> <i>acerifolium</i> ...	Muchkunda,	Introduced.
6	" <i>lanceifolium</i> [<i>P. suberifolium</i> , Maiden]	"	Bhábar.
7	<i>Melochia</i> <i>corchorifolia</i> , [<i>Riedelia</i> <i>corchorifolia</i> , Dec.]	...	Up to 4,000 feet.
8	<i>Waltheria</i> <i>indica</i> , [<i>Melochia</i> <i>corchorifolia</i> , Wall.]	...	" "
ORDER 25.—TILIACÆÆ.			
1	<i>Grewia</i> <i>oppositifolia</i> ...	Bhekla, bhen-gál.	Common.
2	" <i>tiliaefolia</i>	Frequent, valleys.
3	" <i>asiatica</i> [<i>elastica</i> , Royle, III.]	Pharsiya	Common, cultivated.
4	" <i>aspidia</i> [nana, Wall.]	Gar bheli	Common.
5	" <i>scabrophylla</i> [<i>sclerophylla</i> , Wall.]	...	"
6	" <i>laevigata</i> [<i>didyma</i> , Roxb.]	Bhimul	"
7	" <i>polygama</i>	Bhábar.
8	<i>Triumfetta</i> <i>pilosa</i> [<i>oblongata</i> , Wall.]	Leahwa, kumariya.	Common.
9	" <i>rhomboides</i> [<i>angulata</i> , Wall.]	Chat kura	Up to 4,000 feet.
10	<i>Corchorus</i> <i>fascicularis</i> ?	...	Introduced.
11	" <i>acutangulus</i> ?	...	"
12	<i>Eleocharis</i> <i>Varunua</i> ..	Ja'pai	Rare.
ORDER 26.—LINÆÆ.			
1	<i>Linum</i> <i>usitatissimum</i> ...	Alai	Introduced.
2	<i>Reinwardtia</i> <i>trigyna</i> ...	Pyra	Very common, March.
3	" <i>tetragyna</i>	Rare.

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 27.—MALPIGHIACEÆ.			
1	<i>Hiptage madablota</i>	March; common up to 3,000 feet.
2	<i>Aspidopterys Wallichii</i> [Hirsa nutans, Wall].	...	June; frequent up to 4,000.
ORDER 28.—ZYGOPHYLLÆ.			
1	<i>Tribulus terrestris</i>	Common.
ORDER 29.—GERANIACEÆ.			
1	<i>Geranium pratense</i>	Rare, Painkanda.
2	" <i>collinum</i>	"
3	" <i>palustre</i> ? [grandiflorum, Edgeworth]	...	" Māna Pāsa.
4	" <i>Wallichianum</i>	Common, Naini Tal and Binsar, July.
5	" <i>Grevilleanum</i> [erioctemon, Don]	...	"
6	" <i>nepalense</i>	Common, May, June.
7	" <i>polyanthos</i>	Rare.
8	" <i>rotundifolium</i>	"
9	" <i>molle</i>	February. Banks of Ganges up to 1,500 feet.
10	" <i>Robertianum</i>	"
11	" <i>lucidum</i>	Common.
12	" <i>ocellatum</i> [bicolor, Royle, III]	Ankri ...	Very common, April, May, Introduced.
13	<i>Tropæolum majus</i>	Very common.
14	<i>Oxalis corniculata</i>	Chalmor: ...	Rare, Pindari, April.
15	" <i>acetosella</i>	Common up to 4,000 feet.
16	<i>Biophytum sensitivum</i>	"
17	" <i>Reinwardtii</i>	Gardens.
18	<i>Impatiens</i> (<i>Brevicaulis</i>) <i>Balsamina</i> , (<i>Longicaulis</i>) <i>Boylei</i> var.	...	Common.
19	" <i>moschata</i> , Edge.	...	"
20	" <i>Thomsoni</i>	12,000 feet.
21	" <i>sulcata</i> [gigantea, Edge.],	...	Frequent.
22	" <i>amplicaulis</i>	Rare.
23	" <i>scabrida</i> [tricornis, Lind, Hamiltoniana, Don].	...	Common, May, June.
24	" <i>bicornuta</i>	Rare.
25	" <i>anphorata</i>	Common, autumn.
	" <i>var umbrosa</i> , Edge, [bicolor, Royle, III.]	...	"
26	" <i>racemosa</i>	Frequent.
27	" <i>laxiflora</i>	"
28	" <i>micranthemum</i>	Rare.
29	" <i>brachycentra</i>	Frequent, summer.
30	" <i>glauca</i>	Rare, Diwāl.

DIVISION 1.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 30.—RUTACEÆ.			
Tribe 1. UTEÆ.			
1	<i>Benninghausenia albiflora</i> , [Ruta albiflora, Wall.]	Pian ghās and upanyā ghās.	Very common, autumn.
Tribe 2. ZANTHOXYLÆ.			
2	<i>Zanthoxylum alatum</i> , [Z. hostile, Wall.]	Timur and tejbal.	Common.
3	" <i>acanthopodium</i> [hostile, Wall.]	Simur ...	Rare, Bnāh Jagewar.
4	" <i>oxyphyllum</i>	Frequent, Gāgar.
5	" <i>Budrunga</i>	Rare.
Tribe 3. TODDALIÆ.			
6	<i>Toddalia aculeata</i> ...	Khaseru ...	Bhābar and valleys.
7	<i>Skimmia</i> [Limonia, Wall. Anquetilla Deco.] <i>Lauricola</i> .	Nehar, knātūrī, gūral pata.	Very common.
Tribe 4. AURANTIÆ.			
8	<i>Glycosmis pentaphylla</i> ...	Potla ...	Common.
9	<i>Murraya exotica</i> ...	Juti ...	"
10	" (<i>Bergera</i>) <i>Königii</i> ...	Gani ...	Very common.
11	<i>Clausena pentaphylla</i>	Bhābar.
12	<i>Limonia acidissima</i> [L. crenulata, Roxb.]	...	"
13	<i>Citrus medica</i> ...	Bijaura ...	"
	var. <i>limonum</i> ...	Jamira ...	"
14	" <i>aurantium</i>	Rare.
15	<i>Feronia Kiephantum</i> ...	Kait ...	Bhābar.
16	<i>Ægle Marmelos</i> ...	Bel ...	"
ORDER 31.—SIMARUBEÆ.			
1	<i>Picrosma quassioides</i>	Valleys; frequent.
ORDER 32.—OCHNACEÆ.			
1	<i>Ochna pumila</i>	Rare.
ORDER 33.—BURSERACEÆ.			
Tribe 1. AURANTIÆ.			
1	<i>Boswellia serrata</i> [thurifera, Roxb.]	Sale, gugar...	Bhābar.
	var. <i>glabra</i> ...	Gugai ...	"
2	<i>Garuga pinnata</i> ...	Kharpat ...	"
ORDER 34.—MELIACEÆ.			
1	<i>Melia Azadirachta</i> [Indica] ...	Nim ...	Introduced. Bhābar.
2	" <i>Azadirach</i> [sempervirens Roxb.]	Betain ...	Common.
3	<i>Haynes trijuga</i> , [Zanthoxylon conaroides, W. & A.]	...	Bhābar.
4	<i>Cedrela Toona</i> ...	Tūn ...	Valleys.
5	" <i>serrata</i> ...	Dāl ...	Very common.

DIVISION I.—POLYPTETEE—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 35.—OLACINEÆ.			
1	<i>Olex scandens</i>	Bhābar.
2	„ <i>nana</i>	Frequent.
ORDER 36.—ILICINEÆ.			
1	<i>Nox dipyræa</i>	Kandāli	Very common.
2	„ <i>odorata</i>	Gar—kaula, garahon,	Frequent.
3	„ <i>excelsa</i>	Kandāli ..	Common
ORDER 37.—CELASTRINEÆ.			
1	<i>Eucynus grandiflorus</i> [lacerus, Don].	..	Bhābar, & to 3,000 feet.
2	„ <i>tingens</i>	Kasuri ...	Frequent, April.
3	„ <i>fimbriatus</i>	Rare, 9,000 feet, April.
4	„ <i>frigidus</i>	„ 10,000 „
5	„ <i>echinatus</i>	Frequent, April
6	„ <i>pendula</i> [japonicus, Wall].	..	„
7	„ <i>Hamiltonianus</i>	Agna ...	Common.
8	<i>Celastrus paniculata</i> [ratius, Roxb]	Malkāgni ...	March. Blum Tāl.
9	<i>Gymnosporia ruft.</i> [Celastrus ruft, Brandis.]	..	Rare.
10	„ <i>Boyleana</i> , [C. spinosus, Lam.]	Gwala-dārim,	Common.
11	„ <i>montana</i> [C. zenzigensis, Lam.]	..	Rare.
12	<i>Elmolenion glaucum</i> [Rothburghii, W. & A., Neri-jadichotoma, Roxb].	Shauria jam- rā s i a n d dehri	Common. A white aphid feeds on this tree.
13	<i>Microtropis discolor</i>	Frequent.
ORDER 38.—RHAMNACEÆ.			
1	<i>Ventilago calyculata</i> [madraspatana, Roxb.]	Rakt p i t a, kāla lag	Common.
2	<i>Zizyphus jujuba</i>	Bair	Cultivated.
3	„ <i>nummularia</i>	„
4	„ <i>vulgaris</i>	„
5	„ <i>oxyphylla</i>	Frequent up to 6,000 feet.
6	„ <i>xylopyrus</i>	Bhābar.
7	„ <i>rugosa</i>	„
8	<i>Burchardia floribunda</i>	Kāla lag	Frequent
9	„ <i>flavescens</i>	„
10	„ <i>Kalgeworthii</i>	Rameshwar.
11	<i>Rhamnus dahuricus</i> [virgatus, Roxb.]	Chātua ...	Common, April
12	„ <i>purpurea</i>	Common.
13	„ <i>triqueter</i>	Ghont ...	„
14	„ <i>procumbens</i>	About 7,000 feet : on limestone.
15	<i>Hovenia dulcis</i>	Rare : cultivated
16	<i>Sageretia oppositifolia</i>	Agalala ...	Common.
17	<i>Gonania leptotachya</i>	Kāla lag ...	„
18	„ <i>nepalensis</i>	Bhābar
19	<i>Hedius lanetolatus</i>	Common : valleys.

DIVISION.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note
ORDER 39.—AMPELIDÆ.			
1	<i>Vitis</i> (Simplicifoliæ) quadrangularis [Cissus edulis].	Harjora	Bhābar.
2	" pubila	...	Rare.
3	" repanda [roosa, Royle Ill.]	...	Frequent; Bhābar.
4	" adnata	...	Rare.
5	" pedicellata	...	"
6	" l-nata	Purcin	Common, May.
7	" var. rugosa	Assānja	" July.
8	" latifolia	Pan-laguli	" Bhābar.
9	" vinifera	...	Wild?
10	" parvifolia	Barain	Common.
11	" (Trifoliata) carnosā	...	Bhābar.
12	" [Ampelopsis, Wight] li-malayana.	Chappar-tang,	Common.
13	" divaricata [tomentosa, W. and A.]	Amila	"
14	" (Digitatæ) obtecta	...	"
15	" (Pedatæ) capriolata [Cissus serrulata, Roxb.]	...	"
16	" lanceolata [Cissus feminea, Roxb.]	Panj-pata	Rānibagh,
17	<i>Leea</i> (Simplicifoliæ) macrophylla	...	Frequent
18	" (Pinnatæ) alata	...	"
19	" aspera	Kurmali	Common.
20	" (Bipinnatæ) sambucina	...	Rare.
ORDER 40.—SAPINDACÆ.			
TRIBE SAPINDÆ.			
1	<i>Cardiospermum Halicacabum</i>	...	Bhābar.
2	<i>Æsculus</i> (Pavia) indica	Pānkar, pan-gri.	Common.
3	<i>Schleichera trijuga</i>	Kusām	Bhābar.
4	<i>Sapindus Mukorossi</i>	Rita.	"
5	" var. detergens	...	Rare.
6	" var. acuminatus	...	Common.
7	<i>Acer</i> (Integrifoliæ) oblongum	Patangtiya...	Common; cups are made of the wood.
8	" levigatum	...	Common.
9	<i>Acer</i> (Trilobata) pentapomicum	...	"
10	" (Quinquelobata) caesium	Kilu	Frequent.
11	" villosum [sterculiaceum Wall?]	...	8,000 feet; cups made of the wood.
12	" candatum [pectinatum Wall.]	...	10,000 feet.
13	" (Septemlobata) pictum [cultatum, Wall.]	...	Very common.
TRIBE DODONÆÆ.			
14	<i>Dodonaea viscosa</i>	...	Rare; Bhābar.
TRIBE MELIANTHÆÆ.			
15	<i>Melanthus major</i>	...	Introduced.

DIVISION I.—POLYPETALÆ—(continued)

No.	Scientific name.	Vernacular name.	Note.
ORDER 40.—SAPINDACEÆ—(concluded).			
Tribe STAPHYLEÆ.			
14	<i>Staphylea emodi</i> ...	Chitra ...	Frequent: Gágar and Sílhl Dabl.
ORDER 41.—SABIACEÆ.			
1	<i>Meliosma dillenifolia</i> ...	Gwep ...	Common about 7,000 feet.
2	" <i>pungens</i> ...	Gardar ...	Frequent; banks of streams.
3	<i>Sabia campanulata</i> ...	Bakal pata ...	Common.
4	" <i>parviflora</i>	" about 5,000 feet.
5	" <i>paniculata</i>	" Bhábar.
ORDER 42.—ANACARDIACEÆ.			
1	<i>Rhus (Simplicifoliæ) cothuis</i> ...	Tung-ani ...	Very common
2	" (<i>Trifoliatæ</i>) <i>parviflora</i> ...	Ranel ...	"
3	" (<i>Impari pinnatæ</i>) <i>semialata</i> ...	Daknila ...	Common.
4	" <i>Wallichii</i> [<i>verniciifera</i> , Royle.]	Bholiao ...	"
5	" <i>succedanea</i> [<i>acuminata</i> , Dec.] ...	Titari ...	"
6	<i>Pistacia integerrima</i> , [<i>Rhus integririma</i> , Wall.]	Kakar slugi,	Valleys; common.
7	<i>Olinia Wedler</i> ...	Jhingho ...	Common; valleys
8	<i>Semecarpus Anacardium</i> ...	Bhiláwa ...	Bhábar.
9	<i>Mangifera indica</i> ...	Am ...	Common; valleys.
10	<i>Buchanania latifolia</i> ...	Pial ...	Bhábar.
11	<i>Spondias mangifera</i> ...	Amra ...	" and valleys.
ORDER 43.—CORIARIACEÆ.			
1	<i>Coriaria nepalensis</i> ...	Mansuri mokola.	Common.
ORDER 44.—MORINGEÆ.			
1	<i>Moringa pterygosperma</i> ...	Salujan ...	Common; Bhábar.
ORDER 45.—LEGUMINOSÆ.			
Tribe PODALYRIÆ.			
1	<i>Piptanthus nepalensis</i> ...	Shulgari ...	Above 8,000 feet.
2	<i>Thermopsis barbata</i>	" 10,000 "
Tribe GENISTEÆ.			
3	<i>Argyrolobium flaccidum</i> [<i>Cytisus flaccidum</i> , Royle]	...	Common, July and August, on limestone.
4	" <i>rosemæ</i>	"
5	<i>Crotalaria (Diffuse) humifusa</i> ...	Gungri ...	Common.
6	" <i>hirsuta</i>	Bhábar.
7	" (<i>Alatæ</i>) <i>slata</i> ...	Gungri ...	Common, autumn.
8	" (<i>Calycineæ</i>) <i>mysorensis</i>	Valley.
9	" <i>albida</i>	Common.
10	" <i>calycina</i>	Frequent.

DIVISION L—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note
ORDER 45.—LEGUMINOSÆ—(continued).			
TRIBE GENISTEÆ—(concluded).			
11	<i>Crotalaria</i> (<i>Calycina sessiliflora</i> [anthylloides, Lam.]	...	Common.
12	" (<i>Erecta</i>) <i>aricea</i> ...	Gungri ...	Valleys, common.
13	" (<i>Kriocarpa</i>) <i>tetragona</i>	"
14	" (<i>Trifoliolata</i>) <i>medicaginea</i> ,	Very common.
15	" " <i>incana</i>	Naturalized.
TRIBE TRIFOLIEÆ.			
SECTION DIGITATÆ.			
16	<i>Trifolium pratense</i>	Cultivated; rare.
17	" <i>repens</i>	" common.
18	<i>Paroetus communis</i> [<i>oxalidifolia</i> , Boyle.]	...	Common, damp ground.
SECTION PINNATÆ.			
19	<i>Trigonella Fœnanum-græcum</i> ...	Methi ...	Cultivated.
20	" <i>polycerata</i> [<i>incisa</i> , Royle]	...	"
21	" <i>gracilis</i>	July, on limestone.
22	" <i>pubescens</i>	Col. Davidson.
23	" <i>corniculata</i>	Frequent.
24	" <i>emodi</i>	Col. Davidson.
25	<i>Medicago parviflora</i>	Cultivated.
26	" <i>alba</i> ...	Garila ...	Common.
27	<i>Medicago lupulina</i>	"
28	" <i>denticulata</i>	"
TRIBE LOTÆ.			
29	<i>Lotus corniculatus</i>	Very common.
TRIBE GALGANEÆ.			
30	<i>Cyamopsis psoraleoides</i>	Bhābar.
31	<i>Indigofera</i> (<i>Sphæroidophora</i>) <i>linifolia</i>	"
32	" (<i>Pancifoliolata</i>) <i>trifoliata</i>	Common.
33	" " <i>tsita</i>	"
34	" (<i>Multifoliolata</i>) <i>hirsuta</i> ...	Nānā-sakina,	
35	" <i>Gerardiana</i> [<i>heterantha</i> , Wall.	Mā-sakina...	Very common, May.
36	" " <i>atropurpurea</i> ...	Kālā-sakina,	Common, August.
37	" " <i>pulchella</i> [<i>arorea</i> , Roxb.]	Sakina ...	Very common, March.
38	" " <i>Dosua</i> [<i>heterantha</i> , Wall.	Sakina ...	Very common, May, June.
39	<i>Colutea arboreocens</i> , var. <i>nepalensis</i> ,	...	Above 10,000 feet, July, August.
40	<i>Mellettia auriculata</i> [<i>Robinia macrophylla</i> , Ro. & B.]	Gojha ...	Very common; valleys, May.
41	<i>Tephrosia candida</i> ...	Lehtia ...	Bhābar. Leaves used to poison fish.
42	" <i>purpurea</i>	Bhābar.

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 45.—LEGUMINOSÆ—(continued).			
TRIBE GALEGÆÆ (concluded).			
43	<i>Sebania egyptiaca</i>	Jait ...	Cultivated.
44	<i>Caragana pygmaea</i> [Genista, versicolor, Royle].	...	Common above 10,000 feet.
45	" <i>Gerardiana</i> [<i>Caragana spinosissima</i> , Royle.]	Dámá ...	8,000 feet, May.
46	" <i>polyacantha</i>	" ...	Rare.
47	" <i>brevispina</i>	" ...	Common above 8,000 feet.
48	" <i>crassicaulis</i>	" " 12,000 feet.
49	<i>Guldenstädtia himalaica</i> ...	Dámá ...	" " 11,000 feet.
50	<i>Astragalus</i> (Pogonophace) <i>trichocarpus</i> (<i>sebanoides</i> , Royle).	Common.
51	" (<i>Pterolotus</i>) <i>hosackioides</i>	Common.
52	" (<i>Hypoglottis</i>) <i>himalayana</i>	Above 10,000 feet.
53	" " <i>Maddenianus</i>	" "
54	" " <i>lencocephalus</i>	Common.
55	" (<i>Phaca</i>) <i>chlorostachys</i>	
56	" " var. <i>trichocarpoides</i> Kurz,	W. Waterfield, Naini Tál.
57	" " <i>emodi</i>	Above 10,000 feet.
58	" <i>gravcolens</i>	Frequent.
59	" (<i>Myobroma</i>) <i>Candolleanus</i>	Above 8,000 feet.
60	" (<i>Ægacantha</i>) <i>polyacanthus</i>	Frequent.
TRIBE HEDYSARÆÆ			
SECTION ESTIPELLATÆ.			
61	<i>Onobrychis sativa</i>	Cultivated.
62	<i>Lespedeza</i> (<i>Eulespedeza</i>) <i>sericea</i>	Common.
63	" " <i>Gerardiana</i> ,	Very common.
64	" (<i>Oxyrambha</i>) <i>macrostyla</i> ,	Common.
65	" " <i>criocarpa</i>	"
66	<i>Hedysarum kumaonense</i>	Rare; found by Strachey and Winterbottom.
67	<i>Zornia diphylla</i> (<i>angustifolia</i> , D.C.)	Common.
68	<i>Smithia senatiiva</i>	Bhábar.
69	" <i>geminiflora</i>	"
70	" <i>ciliata</i>	Common near streams.
71	<i>Neochynomea indica</i>	Up to 4,000 feet.
SECTION STIPELLATÆ.			
72	<i>Urania picta</i>	Common.
73	" <i>crinita</i>	"
74	" <i>lagopus</i>	"
75	" <i>prunellifolia</i>	Valleys.
76	<i>Alysicarpus</i> (<i>Microcalycina</i>) <i>vagnalis</i> ,	Common to 4,000 feet.
77	" (<i>Microcalycina</i>) <i>brunneifolia</i> ,	" "

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 45 —LEGUMINOSÆ—(continued).			
TRIBE HEDYSARÆÆ—(concluded).			
SECTION STIPELLATÆ—(concluded).			
78	<i>Alysicarpus</i> (<i>Microcalycine</i>) <i>rugosus</i> ,	S a i n d a s asinda.	Common to 4,000 feet.
79	<i>Oncoclinia dalbergioides</i> [<i>Dalbergia on-</i> <i>jeinensis</i> , Roxb.]	...	Up to 4,000 feet, March.
80	<i>Desmodium</i> (<i>Phyllodium</i>) <i>pulchellum</i> ,	...	Bhābar.
81	" (<i>Pteropoma</i>) <i>triquetrum</i> ,	...	Common.
82	" (<i>Scorpiurus</i>) <i>laxiflorum</i> ,	...	" Naini Tāl.
83	" (<i>Podocarpus</i>) <i>polycar-</i> <i>pum</i>	Rare.
84	" (<i>Dolineria</i>) <i>floribundum</i>	...	Very common, August.
	[<i>multiflorum</i> D. C.]		
85	" " <i>oxyphyllum</i>	...	Common; Binsar "
	[<i>polycarpum</i> , Wall.]		
86	" " <i>tiliaefolium</i> [<i>nutans</i> ,	...	Very common, July.
	Wall., <i>argenteum</i>		
	Wall.]		
87	" (<i>Heteroloma</i>) <i>Gangeticum</i> ,	...	Valleys.
88	" " <i>latifolium</i>	Frequent.
89	" " <i>aequat</i>	"
90	" " <i>concinnum</i>	...	Common up to 7,000
	[<i>penduliflorum</i> , Wall.]		feet.
91	" (<i>Nicolsonia</i>) <i>polycarpum</i>	...	"
	D.C. [<i>angulatum</i> , Wall.]		
92	" (<i>Sagotis</i>) <i>triflorum</i>	Very common,
			autumn.
93	" " <i>parviflorum</i>	"
94	" (<i>Pleurolobium</i>) <i>gyrans</i>	Common.
95	" " <i>gyroides</i>	Bhābar.
TRIBE VICIÆÆ.			
96	<i>Acris precatorius</i> ...	Gunchi rakti,	Bhābar and valleys.
97	" <i>pulchellum</i> ...	Laguli-imli,	"
98	<i>Cicer arletinum</i> ...	Chana ...	Cultivated; Bhābar.
99	" <i>soongaricum</i> [<i>microphyllum</i> ,	...	About 12,000 feet;
	Royl.]		rare.
100	<i>Vicia</i> (<i>Ervum</i>) <i>tetrasperma</i>	Rare.
101	" " <i>hirsuta</i> ...	Kuri ...	In cultivated fields.
102	" (<i>Cracca</i>) <i>tenera</i>	Frequent; Adwani,
			April.
103	" " <i>molli</i>	High levels; rare.
104	" " <i>pallida</i>	Common.
105	" " <i>rigidula</i>	Rare.
106	" (<i>Euvicia sativa</i> ...	Kur-kosha ...	Cultivated.
107	<i>Lathyrus</i> (<i>Kalathyrus</i>) <i>sphaca</i> ...	Ghora-kolon,	Common.
108	" " <i>sativus</i> ...	Matar ...	Cultivated.
109	" " <i>sphaericus</i> [<i>an-</i> <i>gulatus</i> , Royl.]	Ningla-kosha,	Common.
110	" (<i>Orbus</i>) <i>lutens</i>	Frequent.
111	<i>Pisum sativum</i> ...	Kolai ...	Cultivated.
112	" <i>arvense</i> ...	"	Cultivated and wild.

DIVISION I.—POLYPETALÆ— (continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 45. LEGUMINOSÆ- (continued).			
Tribe PHASEOLÆ.			
SECTION GLYCINÆ.			
113	<i>Shuteria vestita</i> , var. <i>involucrata</i>	Common.
	var. <i>densiflora</i>	Rare.
114	<i>Dumasia villosa</i>	Common.
115	<i>Glycine soja</i> [<i>Soja hispida</i> , Moench.]	Bhat ...	Cultivated; very common.
SECTION ERYTHRINÆ.			
116	<i>Mucuna atropurpurea</i> ...	Baldaka ...	Bageshwar.
117	" <i>pruriens</i> ...	Goncha ...	Valleys; common, May.
118	<i>Erythrina indica</i> ...	Rungra ...	February; Bhábar.
119	" <i>resupinata</i>	Frequent; Háwalbagh, March and October.
120	" <i>suberosa</i> ...	Rungra ...	March, up to 4,000 feet.
121	" <i>arborescens</i> ...	"	August, common.
SECTION GALACTULÆ.			
122	<i>Spatholobus Roxburghii</i> [<i>Butea parviflora</i> , Roxb.]	Maola ...	Bhábar.
123	<i>Butea frondosa</i> ...	Dhák ...	Up to 4,000 feet.
124	<i>Butea superba</i>	Near Talli Mainari hill Loch, September.
125	<i>Pueraria tuberosa</i> ...	Billi-kand ...	Very common, March, April.
126	" <i>Stracheyi</i>	Rare: Kálimandi, 7,900 feet.
SECTION EUPHABOLÆ.			
127	<i>Phaseolus vulgaris</i>	Cultivated; Bhábar <i>parviflora</i> .
128	" <i>trilobus</i> ...	Chhihin ...	" Bhábar.
129	" <i>aconitifolius</i> ...	Moth ...	" Bhábar.
130	" <i>mungo</i> ...	Mung ...	" not common.
	" var. <i>radiatus</i> ...	Urd ...	" common.
131	" <i>calcaratus</i> [<i>torosus</i> , Roxb.]	Gurudh ...	" and wild red and white varieties.
132	<i>Vigna</i> (<i>Euvigna</i>) <i>catiang</i> [<i>Dolichos sinensis</i> , Linn.]	Riansh ...	" common, three varieties.
	" var. <i>sinensis</i> ...	Lohiya ...	"
133	" (<i>Plectropis</i>) <i>vevillata</i> [<i>Phaseolus angustifolius</i> , Wall.]	Ban-mung ...	Very common, autumn.
134	<i>Pachyrrhizus angulatus</i>	Cultivated; Bhábar, root edible.
135	<i>Dolichos Lablab</i> ...	Shimi ...	Gardens, cultivated up to 7,000 feet.
136	" <i>biflorus</i> [<i>uniflorus</i> , D. C.]	Gahat ...	Cultivated up to 3,000 feet.
137	" <i>falcatus</i> ...	Gahatiya? ...	Common.

DIVISION I. -- POLYPETALÆ (continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 45. LEGUMINOSÆ (continued).			
Tribe PHASEOLEÆ (concluded).			
SECTION CAJANINÆ.			
138	<i>Alysicia</i> (<i>Alycia</i>) <i>mollis</i>	Common.
139	" " <i>grandiflora</i>	Rare; Bageshwar.
140	" " [<i>Cantharo</i> sp. <i>minum</i>] <i>scarabaeoides</i>	Common; Bhábar and outer hills.
141	" " <i>platycarpa</i> ,	Rare.
142	<i>Cajanus</i> <i>flavus</i> ...	Arhar ...	Cultivated; Bhábar.
143	<i>Eriosema</i> <i>chinense</i> [<i>Crotalaria</i> <i>tuberosa</i> , Don]	Common; Simota hill.
144	<i>Rhynchosia</i> <i>pseudo-cajan</i> ...	Shuli, phú-sarpata, ...	"
145	" <i>minima</i>	Low elevations; rare.
146	" <i>himalensis</i>	"
147	<i>Flemingia</i> (<i>Ostrya</i> <i>lium</i>) <i>strobilifera</i> ,	Very common.
148	" " <i>var. fruticulosa</i>	"
149	" " <i>Chappier</i>	Rare.
149	" (<i>Chalaria</i>) <i>paniculata</i>	At low elevations.
150	" (<i>Flemingiastrum</i>) <i>congesta</i> [<i>semialata</i>], ...	Bhatua ...	Common.
151	" " <i>var. nana</i>	Common.
151	" (<i>Rhynchosoides</i>) <i>vestita</i> , ...	Muskila ...	Edible tuber; common.
Tribe DALBERGIEÆ.			
SECTION ALTERNIPOLIOLITÆ.			
152	<i>Dalbergia</i> <i>Sisoo</i> ...	Sisoo ...	Bhábar and valleys.
153	" <i>rimosa</i>	"
154	" <i>lanceolaria</i> [<i>frondosa</i> , Roxb.]	Frequent.
155	" <i>volubilis</i> ...	Bhatia ...	Common.
156	<i>Pterocarpus</i> <i>marsupium</i> ...	Bijai-sal ...	Near Barindeo.
SECTION OPOSITOPOLIOLITÆ.			
157	<i>Pongamia</i> <i>glabra</i> ...	Pit pápar ...	Bhábar, April
158	<i>Derris</i> <i>robusta</i> ...	Baro ...	Up to 5,000 feet
Tribe SOPHOREÆ.			
159	<i>Sophora</i> <i>mollis</i> [<i>Edwardsia</i> <i>mollis</i> , Royle].	Pahār-gangri, ...	Very common, March.
Tribe CÆSALPINIÆ.			
160	<i>Cæsalpinia</i> [<i>Gutlandina</i>] <i>Bouducella</i> , ...	Karonj ...	Up to 2,500 feet
161	" [<i>Poinciana</i>] <i>pulcherrima</i> , ...	Krishna churra, ...	Gardens.
162	" " <i>sepiaria</i> ...	Aira ...	Common.
163	<i>Parkinsonia</i> <i>aculeata</i> ...	Wiláyati karkar, ...	Introduced.
164	<i>Poinciana</i> <i>regia</i>	Gardens; rare.

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 45.—LEGUMINOSÆ—(continued).			
TRIBE CASSIÆ.			
165	<i>Cassia (Fistula) Fistula</i> ...	Rajbriksh, kistota.	May, June, up to 4,000 feet.
166	" (<i>Senna</i>) <i>occidentalis</i> ...	Banār ...	Bhābar.
167	" " <i>Sophora</i> [purpurea, Roxb.] ...	" ...	"
168	" " <i>Tora</i> ...	" ...	Common.
169	" " <i>angustifolia</i> [lanceolata, Royle.] ...	Sanna, makki.	Cultivated; Bhābar.
170	" (<i>Pterohegma</i>) <i>glauca</i> [arborescens, Roxb.] ...	" ...	Bhābar.
171	" (<i>Lasiorhagma</i>) <i>Abana</i> [coccinea, Wall.] ...	Banār ...	Common.
172	" " <i>pumila</i> ...	Silpatiya ...	"
173	" " <i>mimoides</i> ...	" ...	Very common.
...	" " <i>var. dimidiata</i> ...	" ...	Common.
...	" " <i>var. Wallichiana</i> ...	" ...	"
TRIBE AMHERSTIÆ.			
174	<i>Saraca indica</i> [<i>Joncsia asoka</i> , Roxb.] ...	Asok ...	Introduced; Sitaban, Bhābar.
TRIBE BAUHINIÆ.			
175	<i>Bauhinia (Dacrydium) racemosa</i> [parviflora Vahl.] ...	" ...	Up to 5,000 feet, common, April.
176	" " <i>malabarica</i> ...	Amli ...	Rare, October.
177	" (<i>Triandra</i>) <i>rotunda</i> [emarginata, Wall.] ...	Kāndla ...	Common, September.
178	" " <i>Vahlil</i> ...	Māln ...	Very common, April, May.
179	" " <i>purpurea</i> ...	Khairwāl ...	Bhābar and valleys, October.
180	" (<i>Pentandra</i>) <i>variegata</i> ...	" ...	Common, April.
...	" " <i>var. candida</i> , ...	" ...	"
TRIBE MIMOSÆ.			
181	<i>Entada scandens</i> ...	" ...	Rare.
182	<i>Parkia biglandulosa</i> ...	" ...	Cultivated; Bhābar.
183	<i>Leucæna glauca</i> [<i>Acacia frondosa</i> , D. C.] ...	" ...	Common, valleys, July.
184	<i>Mimosa pudica</i> ...	Lajwantl ...	Naturalized, Bhābar.
185	" <i>rubraulis</i> [<i>octandra</i> , Roxb.] ...	Agfa ...	August; common up to 4,000 feet.
TRIBE ACACIÆ.			
186	<i>Acacia (Globiferae) Farnesiana</i> ...	Willāyuti babāl.	Bhābar and valleys.
187	" " <i>arabica</i> ...	Babāl ...	"
188	" " <i>churra</i> ...	Bhes ...	Bank of streams.
189	" (<i>Spicatar</i>) <i>Catechu</i> ...	Khair ...	Very common; Bhābar.
190	" " <i>modesta</i> ...	Phillal ...	Rare, Bhābar.
191	" " <i>lenticularis</i> ...	Khain ...	Barnico.
192	" (<i>Vulgaris</i>) <i>concinna</i> [rugata, D.C.] ...	Ban-ritha ...	Bhābar.
193	" " <i>Intala</i> [Cassia, Linu.] ...	Katrar ...	Very common, Bhābar.
194	" " <i>pennata</i> ...	Agfa ...	"

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 45.—LEGUMINOSÆ—(concluded).			
TRIBE INCÆ.			
195	<i>Albisia</i> (<i>Multifoliolata</i>) <i>Lebbek</i> ...	Tantiya, halsis	Common, April, June.
196	" " <i>odoratissima</i> ...	Stris ...	Bhábar, April, June.
197	" " <i>ser. mollis</i> ...	" ...	" May
197	" " <i>procera</i> ...	Baro ...	Up " to 5,000 feet,
198	" " <i>Julibrissin</i> ...	Lál stris ...	May.
199	" " <i>stipulata</i> (<i>Smithiana</i> , Wall.)	Up to 4,000 feet, May.
200	" <i>Páucifoliolata</i> (<i>luclida</i>	Bhábar, April.
201	<i>Pithecoleblum</i> (<i>Unguiculata</i>) <i>dulce</i> ...	Willáyati (mali, Dakhni baddi.	Gardens, Bhábar.
202	" <i>Clypearia</i> (<i>bigeminum</i> ...	Kachlora ...	Up to 3,000 feet, April.
ORDER 46.—ROSACEÆ.			
TRIBE PRUNÆ.			
1	<i>Prunus</i> (<i>Amygdalus</i>) <i>persica</i> ...	Ara ...	Cultivated; very common.
2	" (<i>Armeniaca</i>) <i>armeniaca</i> ...	Chufaru and sardaru.	"
3	" (<i>Cerasus</i>) <i>avium</i>	Cultivated up to 2,000 feet.
4	" " <i>cerasus</i>	"
5	" " <i>Jacquemontii</i> ...	Padam pays	Rare, Malári, Edgeworth.
6	" " <i>Puddum</i>	Very common, November.
7	" (<i>Euprurus</i>) <i>communis</i> var., <i>domestica</i> var.	Bhotiya bádám.	Cultivated; Almora.
8	" (<i>Laurocerasus</i>) <i>Pandus</i> (<i>cornuta</i> , Royle).	Ladaki Jáman ...	Common.
9	" " <i>nepalensis</i> ...	"	About 6,000 feet.
10	" " <i>undulata</i> (<i>capriculata</i> , Wall.) ...	"	"
11	<i>Prinsepia</i> <i>utilis</i> ...	Bekia, jhatila,	Very common.
TRIBE SPIRÆÆ.			
12	<i>Spiræa</i> (<i>Ulmaria</i>) <i>vestita</i> [<i>Kamtschatica</i> , Lindley.	...	Above 8,000 feet.
13	" (<i>Arunus</i>) <i>aruncus</i> [<i>triternata</i> , Wall.]	" 10,000 feet.
14	" (<i>Sorbaria</i>) <i>sorbifolia</i> [<i>Lindley and Wall.</i>]	Dhaniyál ...	" 8,000 "
15	" (<i>Chamaedrya</i>) <i>bella</i> ...	Jhár ...	Common.
16	" " <i>canescens</i> (<i>cuneifolia</i> , Wall.) ...	" ...	"
17	" " <i>vacciniifolia</i>	Rare.

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name	Vernacular name.	Note.
ORDER 46.—ROSACEÆ—(continued).			
TRIBE RUM &.			
18	<i>Rubus</i> (<i>Simplicifolius</i>) <i>calycinus</i> [Dall.] barda calycina, D.C.]	.	Rare.
19	" " <i>acuminatus</i> ...	Lāl hīālū ...	"
20	" " <i>paniculatus</i> [tilliacus, Wall.]	Kālā " ...	Common.
21	" " <i>mollicarpus</i> [rogatus, Smith].	Lāl " ...	" low levels
22	" " <i>reticulatus</i> ...	Kālā " ...	" high levels
23	" " <i>lunatus</i> [tilliacus, St. & W.]	" ...	About 6,000 feet.
24	" (<i>Digitatus</i>) <i>alpestris</i> ...	"	Rare.
25	" (<i>Prifoliatus</i>) <i>saxatilis</i> ...	Lāl hīālū	Above 10,000 feet.
26	" " <i>nudans</i> ...	Pingla " gangur.	Very common at high elevations.
27	" " <i>nivens</i> ...	Lāl " ...	Very common.
...	" " <i>var. concolor</i> ...	Pingla " ...	Common, China hill.
28	" " <i>macilentus</i> [super Royle.]	" " ...	"
29	" " <i>ellipticus</i> [rotundifolius, Wall; Gower-phyll, Roxb. flavus, Don.]	Hīālū	March, very common. Fruit much eaten in April and May.
30	" (<i>Pinnatus</i>) <i>biflorus</i> ...	Pingla hīālū,	Frequent.
31	" " <i>lasiocarpus</i> [roseoflorus, Roxb.]	Kālā " ...	Very common, Bhābar, and up to 7,500 feet.
32	" " <i>foliolosus</i> ...	Safel " ...	Frequent
...	" " <i>roseifolius</i> ...	Lāl " ...	"
TRIBE POTENTILLA &.			
33	<i>Geum</i> (<i>Engelm</i>) <i>urbanum</i>	Rare.
34	" (<i>Sieversia</i>) <i>elatum</i> [Potentilla adnata, Wall.]	.	Common, above 8,000 feet.
35	<i>Fragaria</i> (<i>Duchesnea</i>) <i>indica</i> ...	Kaphaliya	" up to 8,000 feet.
36	" (<i>Eufragaria</i>) <i>vesca</i> [nubicola, Wall.]	"	above 7,000 feet
37	<i>Potentilla</i> (<i>Sibbaldia</i>) <i>Sibbaldi</i>	" Dākuri, at 10,000 feet
38	" " <i>avillifera</i>	Rare.
39	" (<i>Trichothalamus</i>) <i>fruticosus</i>	Above 12,000 feet.
40	" " <i>ambigua</i>	Colonel Davidson.
41	" " <i>ericarpa</i>	Above 12,000 feet.
42	" (<i>Eupotentilla</i>) <i>Mosmanni</i> [polyphylla, Wall.]	.	High elevations.
43	" " <i>fulgens</i> [splendens, Wall.]	.	Common
44	" " <i>Leschenaultiana</i> .	.	"
45	" " <i>fragarioides</i>	Very common
46	" " <i>perlacularis</i>	Rare. Malāri.
47	" " <i>leucantha</i>	Rare, high levels
48	" " <i>microphylla</i>	"

DIVISION I.—POLYPETALÆ--(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 46.—ROSACEÆ—(continued).			
TRIBE POTENTILLÆ—(concluded)			
49	Potentilla (Eupotentilla) multifida	Rare, Niti pass, 16,000 feet.
50	" " sericea	Rare, high elevations.
51	" " nepalensis	Common.
52	" " argyrophylla	Rare.
...	" " var. atrosanguinea.	...	Common.
53	" " nivalis	Rare, high elevations.
54	" " Kleiniana	Common.
55	" " alpina	"
TRIBE POTERIFÆ.			
56	Agrimonia Eupatorium	Common, June, July.
TRIBE ROSÆ.			
57	Rosa (Eleutherostylax) macrophylla.	...	Frequent; China and Binsar.
58	" " Webblana	Above 8,000 feet, common.
59	" " sericea [tetrapetala, Royle.]	...	" "
60	" (Syntylax) monchata [Brunonia, Wall].	...	Very common; May-June.
TRIBE POMÆ.			
61	Cydonia vulgaris ...	Bibi ...	Cultivated up to 5,000 feet.
62	Eriobotrya japonica ...	Loquat ...	" Bhábar.
63	Pyrus (Malus) malus ...	Seb ...	" and perhaps wild
64	" " baccata ...	Ban melol ...	Common; sometimes cultivated.
65	" (Eupyrus) communis ...	Nashpáti ...	Cultivated.
66	" " Pashia [variolosa, Wall].	Melal ...	Very common.
67	" " kumaoni ...	" ...	Common.
68	" (Aria) lanata ...	Galion ...	Above 8,000 feet, common.
69	" " vestita ...	Mauli ...	Common.
70	" (Sorbus) Aucuparia	Above 12,000 feet.
71	" " foliolosa [ursina, Wall].	Salia halia ...	" 8,000 feet, very common
72	Stranvossia glaucescens [Photinia dubia, Mucken].	Garuchal, sund.	Very common about 5,000 feet.
73	Cratogeomys Oxycnemitha	Cultivated; rare.
74	" " ornulata ...	Gingáru ...	Very common.
75	Cotoneaster (Planifolia) tuccillaris ...	Raus, rois ...	Common. Walking sticks made of the wood.

DIVISION I. —POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 46.—ROSACEÆ—(concluded).			
TRIBE POMES (concluded).			
76	<i>Cotoneaster (Planifolia) acuminata</i> ...	Rana, rois ...	Common. Walking sticks made of the wood
77	" " <i>multiflora</i>	Rare; Nitt pass.
78	" " <i>nummularia</i>	Colonel Davidson.
79	" (<i>Recurvifolia</i>) <i>microphylla</i> ...	Gari ...	Very common everywhere.
80	" " <i>thymifolia</i> ...	" ...	Common.
81	" " <i>lucifolia</i> ...	" ...	"
ORDER 47.—SAXIFRAGACEÆ.			
TRIBE SAXIFRAGÆ.			
1	<i>Astilbe rivularia</i> [<i>Spiraea barbata</i> , Wall.]	Common.
2	<i>Saxifraga (Nephrophyllum) odontophylla</i>	Above 10,000 feet.
3	" (<i>Hirculus</i>) <i>palpebrata</i>	Rare; found near Râ-lam at 15,000 feet.
4	" " <i>sagittoides</i>	" 10,000 "
5	" " <i>diversifolia</i>	" 10,000 "
6	" (<i>Thoraphila</i>) <i>micrantha</i>	" 12,000 "
7	" (<i>Microphylla</i>) <i>imbricata</i>	" 10,000 "
8	" (<i>Spinulase</i>) <i>brachypoda</i>	" 11,000 "
9	" " <i>fimbriata</i>	" 10,000 "
10	" " <i>filicollis</i>	" 10,000 " Pin-dari.
11	" (<i>Flagellares</i>) <i>Brunoniana</i> ,	China peak.
12	" (<i>Bergenia</i>) <i>ligulata</i> ...	Silphora ...	March; very common.
...	" <i>var. ciliata</i> , Royle	" ...	Very common.
13	<i>Chrysosplenium (Oppositifolia) nepalense</i>	Common.
...	" <i>var. sulcatum</i> ,	"
14	" " <i>trichospermum</i> ,	"
15	" (<i>Alternifolia</i>) <i>tenellum</i>	Near Kedar Kanta.
16	<i>Parnassia nubicola</i>	Common.
TRIBE HYDRANGÆÆ.			
17	<i>Hydrangea (Truncata) altissima</i>	Frequent.
18	" " <i>aspera</i> ...	Bhâjhatâ ...	"
19	" " (<i>Coronata</i>) <i>vestita</i> , ...	" ...	"
20	<i>Deutzia corymbosa</i>	May, common, high elevations.
21	" <i>staminea</i>	May, very common.
22	" <i>macrantha</i>	May, rare.
23	<i>Philadelphus coronarius</i> , <i>var. toukousus</i> , ...	Bukyal, dhaniyal.	June, frequent at high elevations. Wân.
TRIBE ESCALONIEÆ.			
24	<i>Itea nutans</i> ...	Garkath ...	Common at low elevations.

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 47.—SAXIFRAGACEÆ—(concluded).			
TRIBE RIBESIÆ.			
25	<i>Ribes</i> (<i>Grossularia</i>) <i>grossularia</i>	Common about 10,000 feet.
26	" (<i>Ribesia</i>) <i>glaciale</i> [acuminatum, Wall.] ...	Kokaliya ...	" above 10,000 feet.
27	" " <i>nigrum</i> ...	l'áphar ...	" about 10,000 feet.
28	" " <i>rubrum</i> ...	" ...	" "
ORDER 48.—CRASSULACEÆ.			
TRIBE ISOSTEMONES.			
1	<i>Tillaea pentandra</i>	Common.
2	<i>Crassula indica</i>	Frequent, Gágár.
3	<i>Cotyledon oreades</i>	Above 13,000 feet.
TRIBE DIPLASTEMONUS OPPOSITIFOLIA.			
4	<i>Bryophyllum calycinum</i>	Bhábar; introduced.
5	<i>Kalanchoe spathulata</i> [varians, Wall.] ...	Bakal patta,	Very common. Goats poisoned by eating this plant.
TRIBE DIPLASTEMONUS ALTERNIFOLIA.			
6	<i>Sedum</i> (<i>Rhodiola</i>) <i>Rhodiola</i> [rosea, Linn.]	Common above 12,000 feet.
7	" " <i>crenulatum</i>	Frequent above 12,000 feet.
8	" " <i>quadrifidum</i>	Common above 12,000 feet.
9	" " <i>asiaticum</i>	" above 11,000 feet.
10	" " <i>linearifolium</i>	" on trees.
11	" " <i>trifidum</i> [sinuatum, Royle]	" September.
12	" (<i>Eusedum</i>) <i>renatum</i>	Common, April May.
13	<i>Sedum</i> (<i>Eusedum</i>) <i>adenotrichum</i>	Common, April, and May.
14	" " <i>trullipetalum</i>	Above 11,000 feet.
15	" " <i>Ewersii</i> [rubrum, Royle]	Frequent.
16	" " <i>multicaule</i>	Common, July.
17	<i>Sempervivum macronatum</i>	Above 10,000 feet.
18	" <i>scabroides</i>	" 10,000 feet.
ORDER 49.—DROSERACEÆ.			
1	<i>Drosera Burmanni</i>	Bhábar.
2	" <i>peltata</i>	Very common.
ORDER 50.—HALORAGACEÆ.			
1	<i>Myriophyllum spicatum</i>	Naini Tál lake.
2	<i>Callitriche verna</i>	"

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 51.—COMBRETACEÆ.			
TRIBE COMBRETÆ.			
1	<i>Terminalia</i> (Catappa) <i>belerica</i> ...	Beliara ...	Frequent; Ganai, March.
2	" " <i>Chesula</i> ...	Har, harira, ...	Common up to 5,000 feet, April, May.
3	" (Pentaptera) <i>Arjuna</i> [glabra, Roxb.] ...	Sáj ...	Bhábar, April, May.
4	" " <i>tomentosa</i> ...	" ...	Common up to 4,000 feet, April.
5	<i>Anogeissus latifolia</i> ...	Bákla ...	" outer range.
6	<i>Combretum</i> (Polvra) <i>decandrum</i> ...	" ...	" Feb., March.
7	" (<i>Eucombretum</i>) <i>nanum</i> ...	Pharsiya ...	Bhábar; very common, March, April.
8	<i>Quiequalis indica</i> ...	" ...	" gardens.
ORDER 52.—MYRTACEÆ.			
TRIBE LEPTOSPERMÆ.			
1	<i>Eucalyptus obliqua</i> ...	Stringy bark ...	Introduced.
2	" <i>globulus</i> ...	Blue gum ...	"
TRIBE MYRTÆ.			
3	<i>Paidium</i> <i>Guyava</i> ...	Amrúd ...	Introduced.
4	<i>Myrtus communis</i> ...	Wiláya ti mendhi. ...	Cultivated, up to 8,000 feet.
5	<i>Eugenia</i> (Jambosa) <i>Jambos</i> ...	Guláb jáman, ...	Cultivated; Bhábar.
6	" (<i>Syzygium</i>) <i>operculata</i> ...	Rái " ...	Bhábar.
7	" " <i>Jambolana</i> ...	Jáman. ...	" and valleys.
TRIBE BARRINGTONIÆ.			
8	<i>Barringtonia acutangula</i> ...	" ...	Banks of streams; Bhábar.
9	<i>Careya herbacea</i> ...	" ...	Bhábar, March.
10	" <i>arborea</i> ...	Kumbh ...	" "
ORDER 53.—MELASTOMACEÆ.			
1	<i>Osbeckia chinensis</i> [angustifolia, Don.] ...	Kukar-makri, ...	Frequent, autumn.
2	" <i>stellata</i> ...	" " ...	Common "
ORDER 54.—LYTHRACEÆ.			
TRIBE AMMANNIÆ.			
1	<i>Ammannia</i> (Rotala) <i>peplodes</i> ...	" ...	Rare. In damp places.
2	" " <i>rotundifolia</i> ...	Darmiya ...	Very common
3	" (<i>Euammannia</i>) <i>salicifolia</i> , ...	" ...	Bhábar swamps, and up to 4,000 feet.
4	" " <i>senegalensis</i> , ...	" ...	Frequent up to 5,000 feet.

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 54.—LYTHRACEÆ (concluded).			
TRIBE LYTHRÆ.			
5	Woodfordia floribunda [Grislen to- mictosa, Roxb.]	Dhal ...	Very common, April.
6	Lawsonia alba ...	Mendhi ...	Common gardens.
7	Lagerstromia (Velagu) indica ...	Gulbahār ...	August, common, gardens.
8	" " parviflora ...	Dhaura ...	Very common, Bhābar.
9	" (Adamben) Flos-Reginæ,	Jāral ...	Rare. Gardens.
10	Punica Granatum ...	Dārim ...	Very com. on up to 5,000 feet.
ORDER 55.—ONAGRACEÆ.			
SECTION POLYSPERMÆ.			
1	Epilobium (Chamænerium) angustifolium.	...	Frequent.
2	" " latifolium	High elevations.
3	" (Lysimachion) hirsutum,	...	Common.
4	" " roseum	Bhimi Tāl.
5	" " tetragonum...	...	Naini Tāl.
6	" " laxum, Royle,	...	
7	Acnothera rosea	Naturalized.
8	" nocturna	"
9	" grandiflora	"
SECTION OLIGOSPERMA.			
10	Circæa latetiana [repens, Wall]	...	Common.
	" alpina [intermedia, Wall]	...	"
ORDER 56.—SAMYDACEÆ.			
1	Casaria graveolens ...	Chilla, chilla-ra.	Fruit used to poison fish.
	" tomentosa ...	" "	" "
ORDER 57.—PASSIFLORACEÆ.			
1	Passiflora, species 1 ...	Cultivated.
ORDER 58.—CUCURBITACEÆ.			
SECTION SIGMATEPÆ.			
1	Trichosanthes palmata ...	Indrāyan ...	Common.
2	" cucumerina	Bhābar.
3	" " anguina ...	Chichinda ...	Cultivated.
4	Herpetospermum caudigerum	Rare.
5	Lagenaria vulgaris ...	Lauka, lauki, tauri.	Cultivated.
6	Luffa aegyptiaca [pentandra, Roxb.]	Ghis torai ...	"
7	" acutangula [triandra]	Torai ...	"
8	Benincasa olerifera ...	Bhuja, petha, kumhra.	"
9	Momordica Charantia ...	Karcia ...	"

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 58.—CUCURBITACEÆ—(concluded).			
SECTION SIGMATOPHEÆ—(concluded).			
10	<i>Momordica dioica</i>	Frequent.
11	<i>Cucumis trigonus</i>	Airālu ...	Common.
12	" <i>Melo</i> [utillissimus, Roxb.] ...	Kakri ...	Cultivated.
13	" <i>sativus</i>	Kakri, khira,	
14	<i>Citrullus Colocynthis</i>	Tarbaza ...	Wild and cultivated.
15	" <i>vilgaris</i>	" ...	Cultivated.
16	<i>Cephalandra indica</i> [<i>Coccinia indica</i> , W. and A.]	Kunderu ...	Bhābar.
17	<i>Cucurbita maxima</i>	Kadu ...	Cultivated.
SECTION ORTHOPHEÆ.			
18	<i>Hyosia laciniosa</i>	Common.
19	<i>Mukia scabrella</i>	"
20	<i>Zehneria umbellata</i>	Gwāla-kakri,	"
	" <i>var. nepalensis</i>	"	"
21	<i>Melothria odorata</i>	Rare.
22	<i>Edgaria darjeelingensis</i>	"
23	<i>Gynostemma pedata</i>	"
ORDER 59.—BEGONIACEÆ.			
1	<i>Begonia</i> (<i>Knebeckia</i>) <i>picta</i>	Common, autumn.
2	" (<i>Trilobaria</i>) <i>amara</i> [tinella, Don.]	" "
ORDER 60.—DATICACEÆ.			
1	<i>Datiscia Cannabina</i>	1 Bhang-jala...	Common.
ORDER 61.—CACTACEÆ.			
1	<i>Opuntia Dillenii</i>	1 Nāg-phansa,	Introduced. Common up to 6,000 feet.
ORDER 62.—FICOIDÆ.			
1	<i>Trianthema crystallina</i>	Bhābar.
2	" <i>pentandra</i>	"
3	<i>Mollugo stricta</i>	Common.
ORDER 63.—UMBELLIFERÆ.			
SECTION HETEROSCHIADÆ.			
1	<i>Hydrocotyle javanica</i>	Common.
2	" <i>retundifolia</i>	Rare.
3	" <i>asiatica</i>	Bhābar.
4	<i>Sanicula europæa</i> [elata, Don.]	Common, June, Bānj- lagr.

DIVISION I.—POLYPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 63.—UMBELLIFERÆ—(concluded).			
SECTION HAPLOSTYLÆ.			
TRIBE AMMINEÆ.			
5	<i>Vicatia conifolia</i>	Khāti Pindari, May.
6	" <i>Stewartii</i>	Rare, Garhwāi.
7	<i>Bupleurum lanceolatum</i>	Common, Junc, Bānj lagr.
8	" <i>Candollii</i>	Above 8,000 feet.
9	" <i>diversifolium</i>	" 7,000 "
10	" <i>falcatum</i> , var. <i>marginata</i> ,	Naini Tāl, common.
11	" <i>longicaule</i>	Above 8,000 feet.
12	" <i>tenuis</i>	Common.
13	<i>Carum</i> (Glabræ) <i>Carui</i>	Rare.
14	" (Hispide) <i>Roxburghianum</i>	Cultivated.
15	" <i>anthifolium</i>	Almora, common.
16	<i>Pimpinella</i> (Glabræ) <i>achilleifolia</i>	Common.
17	" [Heutera, Edgw.] <i>acuminata</i>	Naini Tāl, common.
18	" [Aconema, Edgw.] <i>tenera</i>	Naini Tāl, on oaks, common.
19	" (Papillææ <i>diversifolia</i> tri- foliata, Wall.	Dhanjari ...	Very common.
20	" " <i>Stracheyi</i>	Rare; banks of Dhaulī.
21	" " <i>capitata</i>	" "
22	<i>Osmorrhiza Claytoni</i> [Lava, Royle]	Frequent.
23	<i>Cherophyllum villosum</i>	"
	" <i>reflexum</i>	Rare.
TRIBE SESELINÆ.			
24	<i>Seseli trilobum</i>	Rare; Niti and Milam.
25	<i>Feniculum vulgare</i> [panmorium, Roxb.]	Sonf ...	Cultivated.
26	<i>Ananthe stolonifera</i>	Low elevations, com- mon.
27	<i>Ligusticum marginatum</i>	Naini Tāl, common.
28	" <i>elatum</i> [Cortia <i>elata</i> , Edgw.]	" "
29	<i>Selinum tenuifolium</i> [Selinum Can- dollii, Edgw.]	Common.
30	" <i>vaginatum</i> [Cortia <i>vagina- ta</i> , Edgw.]	Frequent.
31	<i>Cortia Lendell</i>	
32	<i>Hypurospermum</i> [Hymenolena, D. C.] <i>Govanianum</i>	High elevations.
33	" "	"
34	" <i>Candollii</i>	
35	" <i>angelicoides</i> "	Rare; Bālam river, 11,500 feet.
36	" <i>dentatum</i> "	Frequent.
37	" <i>stellatum</i> "	Rare; Milam glacier.
37	" <i>densiflorum</i> "	High elevations.

DIVISION I.—POLYPETALÆ—(concluded).

No.	Scientific name.	Vernacular name.	Note.
ORDER 63.—UMBELLIFERÆ—(concluded).			
SECTION HAPLOXYGIÆ—(concluded).			
TRIBE PEUCEDANÆ.			
38	<i>Peucedanum graveolens</i> [Anethum sowa, Roxb.]	See ...	Cultivated.
39	" <i>Dhana</i>	Up to 4,000 feet.
40	<i>Heracleum Brunonis</i>	Rare.
41	" <i>candicans</i>	Common.
42	" <i>nepalense</i>	Frequent.
SECTION DIPL. XYGIÆ.			
43	<i>Coriandrum sativum</i> ...	Dhaniya ...	Cultivated.
44	<i>Daucus Carota</i>	"
45	<i>Caucalis Anthriscus</i>	Very common.
ORDER 64.—ARALIACEÆ.			
SECTION IMBRICATÆ.			
1	<i>Aralia cisalfolia</i>	Rare.
2	" <i>cachemirica</i> [<i>Panax triplinatum</i> , Wall.]	...	Pindari, autumn.
3	<i>Pentapanax parasiticum</i> ...	Kot semal ...	Up to 7,000 feet.
4	" <i>Lenchenaultii</i> , var. <i>umbellatum</i> [<i>Hedera fragrans</i> , Don.] [<i>Panax brijugum</i> , Wall.]	...	"
SECTION FALVATÆ.			
5	<i>Heptapleurum impressum</i> [tomentosum, Don.]	...	Common.
6	" <i>elatum</i>	Rare.
7	" <i>venulosum</i> [<i>Hedera terebinthacea</i> , Wall.]	...	Up to 3,000 feet.
8	<i>Heteropanax fragrans</i> [<i>Panax fragrans</i> , Roxb.]	...	Low elevations, November.
9	<i>Hedera helix</i> , var. <i>maculifolia</i> ...	Bānda ...	Very common.
ORDER 65.—CORNACEÆ.			
1	<i>Alangium Lamarchii</i> [hexapetalum, Roxb., decapetalum, Wight].	...	Bhābar, rare. March.
2	<i>Maria bigoniifolia</i> ...	Tumri ...	Bageswar, May.
3	<i>Cornus macrophylla</i> ...	Kāshi ...	Common, May.
4	" <i>oblonga</i>	Frequent, September.
5	" <i>nervosa</i> , Royle	...	Rare.
6	" <i>capitata</i> [<i>Benthamia fragifera</i>],	Bānaura ...	Common. Fruit edible, summer.

DIVISION II.—MONOPETALÆ.

No.	Scientific name.	Vernacular name.	Note.
ORDER 1.—CAPRIFOLIACEÆ.			
TRIBE SAM. UCEÆ.			
1	<i>Viburnum c. tinifolium</i> , Don. ...	Gwiya ...	Very common, June
2	" <i>Mullaha</i> , Don [<i>stellulatum</i> , Wall.].	Lāl tittmuliya,	Common, July.
3	" <i>cylinrifolium</i> , Don. [<i>coriaceum</i> , Blume.].	Kāla tittmuliya,	" January.
4	" <i>punctatum</i> , Don. [<i>acuminatum</i> , Wall.].	...	Common, Binsar, July.
5	(<i>Trilocularia</i>) <i>nervium</i> , Don. [<i>grandiflorum</i> , Wall.].	...	" Upper Sārju.
6	" <i>erubescens</i> , Wall.]	Common, April-June.
TRIBE LONICERÆ.			
7	<i>Tricostium himalayana</i> ...	Dur-bank ...	Above 8,000 feet.
8	<i>Abelia triflora</i> ...	Mauri ...	Common, June.
9	<i>Lonicera diversifolia</i> , Wall. [<i>L. quinquelocularis</i> , Harl.].	Bhat-kukri,	" June and July.
10	<i>Lonicera orientalis</i> [<i>Govaniana</i>]	Rare; Bhatkot, June.
11	" <i>alpigena</i> [Webbiana, Wall.]	Rare; Pindari, June.
12	<i>Leycesteria formosa</i> [<i>Lonicera quinquelocularis</i> , Harl.]	Nalkaru, karnaliya.	Common. Confused by Hardwicke with No. 10.
ORDER 2.—RUBIACEÆ.			
TRIBE NAUCLEÆ.			
1	<i>Anthocephalus</i> (<i>Naucllea</i>) <i>Cada ba</i> ,	Kadani ...	Cultivated, June.
2	<i>Adina</i> (<i>Naucllea</i>) <i>cordifolia</i> ...	Haidu ...	Common; Haidwāni, July.
3	<i>Stephagyne</i> (<i>Naucllea</i>) <i>parvifolia</i> ...	Phaidu ...	" Bhābar, June.
TRIBE CINCHONACEÆ.			
4	<i>Hymenopogon parasiticus</i> ...	Bhainal ya - padera.	(On oaks at Binsar.
5	"	"	"
6	<i>Hymenodictyon excelsum</i> ...	Bhalena ...	Common, June.
7	" <i>flaccidum</i>	Rare, June.
8	<i>Wendlandia exserta</i> [W. <i>cinerea</i> Rondeletia <i>exserta</i>].	Chilkiya ...	Common, March.
9	" <i>tinctoria</i>	Bhābar, February.
TRIBE HEDYOTIDÆ.			
9	<i>Arrostemma sarmentosum</i>	Frequent.
10	<i>Hedyotis Lindleyana</i>	Common.
11	" <i>calycina</i>	"
12	<i>Oldenlandia Burmanniana</i>	"
13	" (<i>Kohautia</i>) <i>coccinea</i>	"
14	" <i>gracilis</i>	"
15	" <i>stipulata</i>	Rare.
16	<i>Ophiorhiza nana</i>	"
17	" <i>Mangus</i>	"
18	" <i>bracteolata</i>	"

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 2.—RUBIACEÆ—(concluded).			
TRIBE GARDENIÆ.			
10	<i>Gardenia turgida</i> [longispina, Mutten.]	Thaulea ...	Bhābar.
20	" <i>latifolia</i> ...	Pindālu ...	"
21	<i>Randia tetrasperma</i> ...	Bhotiya gindgar u a n d Bara garl.	Very common.
22	" <i>dumetorum</i> ...	Malphul ...	Bhābar.
23	" <i>nigrosa</i> ...	Pindālu ...	"
TRIBE KNOXIÆ.			
24	<i>Knoxia mollis</i>	Frequent.
TRIBE IXORIÆ.			
25	<i>Pavetta tomentosa</i> [<i>Ixora tomentosa</i> , Roxb.]	Pudera ...	Common.
26	<i>Coffea bengalensis</i> , ...	Kath jahi ..	Frequent, March.
TRIBE PADERIÆ.			
27	<i>Hamiltonia scabra</i> , Dec. [suavcolens, var. <i>asurca</i> , Wall.]	Jogiya-padera,	Common, November.
28	<i>Leptodermis lanceolata</i> ...	Pudera ...	Very common, summer.
TRIBE SPERMACEIÆ.			
29	<i>Spermococo articularis</i>	Common.
30	" <i>lasiocarpa</i> [stricta, Roxb.]	...	"
TRIBE GALIACEÆ.			
31	<i>Rubia cordifolia</i> ...	Mājethi	Common.
32	" <i>nervosa</i> , Edgc.	Rare; Gangoli bridge.
33	" <i>Munjiata</i>	Rare.
34	" <i>alata</i>	"
35	<i>Galium asperifolium</i> ...	Kuri ...	Common.
36	" <i>elegans</i> ...	" ...	"
37	" <i>Hamiltonii</i> [<i>G. latifolium</i> , Don.]	" ...	Frequent.
38	" <i>hirtiflorum</i> [<i>G. ciliatum</i> , Don.]	" ...	" Binsar.
39	" <i>sparino.</i> ...	" ..	Introduced.
40	<i>asperuloides</i> , Edgc., [tricornis, Dr. King.]	" ..	Very common; above 10,000 feet.
ORDER 3.—VALERIANIÆ.			
1	<i>Nerlostachys grandiflora</i>	Rare.
2	<i>Valeriana Wallichii</i> ...	Shani ...	Frequent.
3	" <i>Harlowickii</i> [<i>V. elata</i> , Don.]	" ...	Common.
4	" <i>tenera</i> ..	" ...	Frequent.

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 4.—DIPSACFÆ.			
1	<i>Morina Wallichiana</i> [bre v i f l o r a , Edg.?	...	Common.
2	" <i>longifolia</i>	Above 10,000 feet.
3	<i>Dipsacus strictus</i> , Don.	China, 8,700 "
4	" <i>inermis</i>	Very common.
5	" <i>levis</i> ? ...	Narso ...	Common.
6	<i>Scabiosa Candollii</i>	" May.
ORDER 5.—COMPOSITÆ.			
TRIBE VERNONIACEÆ.			
1	<i>Vernonia</i> (<i>Stengelia</i>) <i>anthelmintica</i> ,	Rare; autumn.
2	" (<i>Tephrosia</i>) <i>cinerea</i> ...	Kalajira ...	Very common up to 8,000 feet.
3	" " <i>montana</i> ...	" ...	Common.
3	" (<i>Xipholepis</i>) <i>terres</i>	Col. Davidson.
TRIBE EUPATORIÆ.			
4	<i>Aidenostemma viscosum</i> , var. <i>latifolium</i>	Common up to 2,500 feet.
5	<i>Ageratum conyzoides</i>	Autumn, common up to 6,000 feet.
6	<i>Eupatorium longicanle</i>	October, common.
TRIBE ASTEROIDEÆ.			
7	<i>Solidago virga aurea</i> [<i>Amphiraphis pubescens</i> , D. C.]	Summer, common.
8	<i>Dichrocephala latifolia</i>	"
9	<i>Cyathocline lyrata</i>	Common up to 4,000 feet.
10	<i>Myriactis nepalensis</i> [olcosa, Edgw.]	Autumn, common.
11	" <i>Wallichii</i> [racilis, Edgw.]	"
12	" <i>Gmelini</i>	October, frequent.
13	<i>Rhynchospermum verticillatum</i>	Autumn; rare.
14	<i>Callistephus chinensis</i> ...	Naipala ...	Introduced.
15	<i>Aster</i> (<i>Kuaster</i>) <i>amelius</i>	Aug., above 8,000 feet.
16	" (<i>Alpigenia</i>) <i>alpinus</i>	July, above 11,000 feet.
17	" (<i>Dr. honeria</i>) <i>uoliusculus</i>	May and June; frequent.
18	" " <i>altaicus</i>	Aug., above 9,000 feet.
19	" " <i>spartioides</i>	"
20	" (<i>Calimeria</i>) <i>peduncularis</i>	September, October; common.
21	<i>Erigeron canadense</i> ...	Salto ...	Introduced, but now very common.
22	" <i>acre</i> [Royel, D. C.] ...	Mirch mál ...	May, August; very common.
23	" <i>multiradiatum</i>	July, August; rare.
24	<i>Microglossa cabulica</i> [<i>Aster ferrugineus</i> , Edgw.].	...	Rare; Badrinath.
25	<i>Conyza veronicifolia</i>	September, common.
26	" <i>viscidula</i> [<i>polyccephala</i> , Edgw.].	...	Feb., up to 4,000 feet.
27	" <i>absinthifolia</i>	Very common, autumn.

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 5.—COMPOSITÆ—(continued).			
Tribe INULOIDÆ.			
28	<i>Blumea Wightiana</i>	Spring; Bhābar.
29	" <i>laccra</i>	" very common.
30	" <i>hieracifolia</i>	May, common.
31	" <i>runcinata</i> [lacinata, Roxb.]	frequent.
32	" <i>oxylonta</i>	"
33	<i>Blumea densiflora</i>	November; Col. Davidson.
34	<i>Laggera alata</i>	Aundok ...	October; common.
35	<i>Sphæranthus hirtus</i> [mollis, Roxb.]	Spring; "
36	<i>Pillago arvensis</i> , var. <i>lutescens</i>	Common.
37	<i>Leontopodium alpinum</i>	Above 12,000 feet.
38	<i>Anaphalis cinnamomea</i>	Jhula, bokla,	Autumn; common.
39	" <i>tripplinervis</i>	"
40	" <i>nubigena</i>	" above 9,000 feet.
41	" <i>mucronata</i>	" above 12,000 feet.
42	" <i>tenella</i> [contorta, Don.]	" very common.
43	" <i>margaritacea</i>	" Col Davidson.
44	" <i>adnata</i>	" common.
45	" <i>araneosa</i>	"
	var. <i>sepidocurrens</i>	" very common.
46	<i>Phagnalon niveum</i>	June; banks of Dhaul near Niti pass.
47	<i>Gnaphalium hypoleucum</i>	Autumn; very common.
48	" <i>luteo album</i>	" common.
49	<i>Cæulia axillaris</i>	October; rice fields; Ganai.
50	<i>Inula</i> [Corvisartia] <i>Royleana</i>	Above 10,000 feet.
51	" (<i>Bubonium</i>) <i>vestita</i>	Spring; up to 3,000 feet.
52	" <i>nervosa</i> [nitida, Edgw.]	Autumn; frequent.
53	" <i>barbata</i>	"
54	" (<i>Cappa</i>) <i>Cappa</i>	Tamagari ...	November; very common.
55	" <i>cuspidata</i>	October; common.
56	" <i>rubicaulis</i>	March; frequent.
57	<i>Vicos indica</i>	Common.
58	<i>Carpesium cernuum</i> var. <i>nepalense</i>	Autumn; common.
	" var. <i>pedunculatum</i> ,	
	" var. <i>trachelifolium</i> ,	
59	" <i>abrotanoides</i> [racemosum, Wall.]	Autumn; rare.
Tribe HELIANTHOIDÆ.			
60	<i>Adenocaulon himalaicum</i>	September; rare.
61	<i>Xanthium strumarium</i>	Up to 5,000 feet.
62	<i>Zinnia elegans</i>	Naturalized.
63	" <i>multiflora</i>	"
64	<i>Boisbeckia orientalis</i>	(lobariya ...	October; very com-

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 5.—COMPOSITÆ—(continued).			
TRIBE HELIANTHOIDEÆ—(concluded).			
65	<i>Eclipta alba</i> , var. <i>erecta</i>	Common.
	<i>prostrata</i>	
66	<i>Blainvillia latifolia</i> ...	Karshaniya...	Very common.
67	<i>Wedelia biflora</i> ?	Rare.
68	<i>Coreopsis</i> (<i>Calliopsis</i>) <i>tinctoria</i>	Naturalized.
69	<i>Bidens</i> (<i>Platycarpus</i>) <i>tripartita</i> [repens, Don.]	June, October ; common.
70	<i>(Pallocarpus pilosa</i>	Very common.
71	<i>decomposita</i> ,	
72	<i>Glossogyne pinnatifida</i>	Low elevations.
73	<i>Chrysanthellum indicum</i>	September; introduced.
74	<i>Galinsoga parviflora</i>	"
TRIBE HELENIOIDEÆ.			
75	<i>Tagetes patula</i>	Gardens and naturalized.
76	<i>erecta</i>	"
TRIBE ANTHEMIDEÆ.			
77	<i>Achillea millefolium</i> , var. <i>cuspidata</i> , Wall.	Rare.
78	<i>Allardia tridactylites</i> ?	Above 15,000 feet.
79	<i>tomentosa</i>	Above 12,000 feet.
80	<i>Contula anthemoides</i> ?	February ; rare.
81	<i>Tanacetum longifolium</i>	September, about 11,000 feet.
82	<i>Artemisia</i> (<i>Dracunculus</i>) <i>scoparia</i> ...	Jhao ...	October, common.
83	<i>parviflora</i>	
84	(<i>Abrotanum</i>) <i>vestita</i> ...	Deo pati ...	September, October ; rare ; Jagtswar.
85	<i>vulgaris</i> [in- dica, Roxb.] ...	Pati ...	September, October ; very common.
	<i>var. hypoleuca</i> , Edgw. "	...	Common.
86	(<i>Absinthium</i>) <i>sierversiana</i> ,	About 12,000 feet ; Jhelam.
TRIBE SENECIENIDEÆ.			
87	<i>Tussilago farfara</i>	Common; banks of streams above 10,000 feet.
88	<i>Doronicum Roylei</i> [Fullertonia Ru- maonensis, D.C.]	High elevations.
89	<i>Gynura nepalensis</i> , var. <i>Thomsoni</i>	March, common, Rānibagh.
90	<i>Emilia sonchifolia</i>	Very common.
91	<i>Senecio</i> (<i>Synotis</i>) <i>alatus</i>	October, November ; Pindari, China hill.
92	<i>densiflorus</i>	October to April; very common.

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 5.—COMPOSITÆ—(continued).			
TRIBE SENECEOIDÆ—(concluded).			
93	<i>Senecio</i> (Synotis) <i>rufinervis</i>	November, frequent.
94	" " <i>Kunthianus</i>	September, above 10,000 feet
95	" " <i>Candolleanus</i>	" "
96	" (<i>Eusenecio</i>) <i>gracilliflorus</i>	October, high levels.
97	" " <i>chrysanthenoides</i> ,	" Pindari.
98	" " <i>pallens</i> [<i>violaceus</i> , Wall].	Rat patiya ...	March, June, very common.
99	" (<i>Ligularia</i>) <i>sibiricus</i>	September, Pindari.
100	" " <i>arnicoides</i>	" "
101	" <i>Cacalia</i> <i>quinquefolia</i>	Above 10,000 feet.
TRIBE CYNAROIDEÆ.			
102	<i>Echinops</i> <i>niveus</i> ...	Kanilla, jao-kanda.	Very common.
103	<i>Carduus</i> <i>nutans</i>	Above 8,000 feet, Jhelam near Niti.
104	<i>Culeus</i> <i>arvensis</i>	Rare.
105	" <i>eriphorus</i> var. <i>involucratus</i> ,	Autumn, common.
106	" <i>ferox</i> , Edgw.	May common.
107	" <i>argyranthus</i>	June common.
108	" var. <i>nepalensis</i>	October, common.
109	" <i>Wallichii</i> (<i>arachnoides</i> , Wall),	...	" "
110	<i>Saussurea</i> (<i>Bracteata</i>) <i>obvallata</i>	September, above 10,000 feet.
111	" (<i>Acaules</i>) <i>Kunthiana</i>	" "
112	" (<i>Congesta</i>) <i>gonypina</i> (<i>sacra</i> , Edgw.)	Kanwal ...	September, common above 10,000 feet.
113	" " <i>grammifolia</i>	" about 10,000 feet.
114	" (<i>Coarctata</i>) <i>pterocaulon</i>	" about 10,000 feet.
115	" (<i>Corymbosæ</i>) <i>candicans</i>	Very common.
116	" " var. <i>scaposa</i>	Common, April, Panwa Naula.
117	" " <i>albescens</i>	September, frequent.
118	" " <i>denticulata</i> ,	September, common.
119	" " <i>hypoleuca</i>	" "
120	<i>Jurinea macrocephala</i> (<i>Dolomieu macrocephala</i> , D. C.).	Kálá tagar ...	" Pindari.
121	<i>Serratula pallida</i>	May, June, very common.
122	<i>Tricholepis furcata</i>	November, Pindari.
123	" <i>elongata</i>	October, common.
124	<i>Voltairella divaricata</i>	March, Haldwani.
125	<i>Carthamus tinctorius</i> ...	Kusam ...	" Cultivated, Bhábar.
TRIBE MUTISACEÆ			
SECTION SUBÆQUALIFLORÆ.			
126	<i>Leucomeria spectabilis</i>	May, common.
127	<i>Ainslia pteropoda</i>	Spring, very common.
	" <i>aptera</i>	" "

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 5.—COMPOSITÆ—(concluded).			
SECTION LABIATÆFLORÆ.			
128	<i>Gerbera lanuginosa</i> (Chaptalia goasypina, Royle)	...	June, high elevations.
129	" <i>nivea</i>	September "
130	" <i>macrophylla</i> (Chaptalia maxima, Don).	...	" "
THIRD CICHORACEÆ.			
131	<i>Cichorium intybus</i>	Cultivated.
132	<i>Pteris hieracioides</i>	Mulpat near Lohaghāt, October.
132	<i>Crepis</i> (<i>Barkhausia</i>) <i>fetida</i>	May, June, very common.
133	" (<i>Youngia</i>) <i>lyrata</i> [runcinata Wall].	...	Very common. An Lactuca?
134	<i>Taraxacum Dens leonis</i>	Very common.
...	" <i>var. eriopodium</i>	
...	" <i>var. parvulum</i>	High elevations.
135	<i>Lactuca</i> (<i>Brachy ramphus</i>) <i>obtus</i> ...	Jangli gobhi,	April, May, very common; yellow.
...	" [<i>Microhynchus nudicaulis</i> and <i>patens</i> , D.C.]	...	
136	" (<i>Scariola</i>) <i>scariola</i>	Low elevations; yellow.
...	" <i>var. sativa</i>	Cultivated "
137	" (<i>Cicerbita</i>) <i>auriculata</i>	May, July, very common; blue.
138	" " <i>longifolia</i>	October, fields, common; blue.
139	" (<i>Lactucopsis sagittarioides</i> ,	...	February, common; blue. An <i>Crepis</i> ?
140	" " <i>Brunoniana</i> [Prenanthes Brunoniana, D.C.]	...	August, October, common; blue.
141	" (<i>Melanoseris</i>) <i>hastata</i>	Autumn, common; Lohaba; blue or white.
142	" " <i>lavigata</i> [macrorhizon, Royle.]	...	September, very common; blue.
143	" " <i>violæfolia</i>	September, high elevations; blue.
144	" " <i>Lessertiana</i>	Common, October; blue.
145	" " <i>dubyra</i> [<i>Dubyra hispid</i> , D.C.]	...	September, very common; yellow.
146	" (<i>Ixeria</i>) <i>glabra</i>	Winter, low elevations; yellow.
147	" " <i>polycephala</i>	" "
148	<i>Prenanthes hispidula</i>	Rare.
149	<i>Sonchus asper</i> ...	Ufat kandia nalaba.	Spring, very common.
150	" <i>oleraceus</i> ...	"	Spring, common, edible.
151	" <i>arvensis</i> ...	"	Spring, common.
152	<i>Microhynchus secundus</i>	Rare; Royle.
153	<i>Tragopogon junceum</i>	June, July, very common.

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 6.—LOBELIACÆ.			
1	<i>Lobelia trigona</i> , Roxb. [trialata, Don.]	...	Common.
2	" <i>pyramidalis</i> , Wall	...	Frequent.
3	" <i>rosea</i> , Wall	...	Rare; Malwa Tāl.
ORDER 7.—CAMPANULACÆ.			
1	<i>Cephalostigma hirsuta</i> ...	Gol ghanna,	Frequent.
2	<i>Wahlenbergia viridis</i> , Edgw.	...	Wācham, near Pindari.
3	" <i>agrestis</i>	Frequent, Banks of Koel.
4	<i>Campanula latifolia</i>	Rare.
5	" <i>sylvatica</i> [integer r i t n a Don.]	...	Common under trees.
6	" <i>canescens</i>	Very common.
7	" <i>argyrotricha</i> [pa l l i d a, Madden].	...	"
8	" <i>colorata ramulosa</i> , Wall]	...	Rare.
ORDER 8.—ERICACÆ.			
1	<i>Gaultheria nummularia</i> ...	Bhalu bor	Common, D ā k u r l, Bhatkot.
2	" <i>trichophylla</i>	Frequent, above 10,000 feet.
3	<i>Andromeda</i> (Pieris) <i>ovalifolia</i> , [A. capricida, Don.]	Alyār	Very common.
4	" (<i>Caasiopie fastigiata</i> [A. cupressiformis, Wal.]	Kamba	Milam and Plinda r i, 12,000 feet.
5	<i>Rhodolend r n arboreum</i> ...	Burāns	Very common.
6	" <i>rar. roseum</i>	Abov 8,000 feet.
7	" <i>campanulatum</i> ...	Chimul	" 10,000 "
8	" <i>anthopogon</i> ...	Tallisfār	" 12,000 "
9	" <i>lepidotum</i>	" 12,000 "
ORDER 9.—MONOTROPEÆ.			
1	<i>Monotropa nepalensis</i> ...	Chhāo	Hinvar, Septem b e r; found by Mrs. Govan.
ORDER 10.—LENTIBULARIÆ.			
1	<i>Utricularia flexuosa</i> [stell a r i s, ... Wight].	...	Common.
2	<i>Pinguicula alpina</i>	Rare.
ORDER 11.—PRIMULACÆ.			
1	<i>Primula</i> (<i>Sphondylia</i>) <i>floribunda</i>	Down to 2,400 feet.
2	" (<i>Primulastrum</i>) <i>petiolaris</i> [nana, Wall].	...	Above to 10,000 feet.
3	" (<i>Athritica</i>) <i>purpurea</i>	"

DIVISION II.—MONOPETALÆ—(continued)

No.	Scientific name.	Vernacular name.	Note.
ORDER 11.—PRIMULACEÆ—(concluded).			
4	<i>Primula Stuartii</i> ...	Bish kopra...	Common.
5	" (<i>Aleuritia</i>) <i>speciosa</i>	Above 10,000 feet
6	" <i>involuta</i>	Common, high elevations.
7	" <i>denticulata</i> , var. <i>erosa</i>	"
8	<i>Androsace sarnientosa</i> , Wall	Above 12,000 feet.
9	" <i>lanuginosa</i> , Wal.	Common.
10	" <i>rotundifolia</i> , Harlw.	Valleys and Bhábhar winter and spring.
11	" <i>incisa</i> Wall, [<i>rotundifolia</i> , Don].	...	Very common above 10,000 feet.
12	<i>Lysimachia</i> (<i>Ephemerum</i>) <i>lobeloides</i> ,	Common.
13	" <i>pyramidalis</i>	"
14	" (<i>Lysimachium</i>) <i>alternifolia</i>	"
15	" <i>debilis</i>	"
16	<i>Anagallis arvensis</i> , var. <i>corulca</i> ...	Jank-mara ...	"
17	<i>Samolus Valerandii</i>	Rare, Sutl river.
ORDER 12.—MYRSINACEÆ.			
1	<i>Marsa</i> (<i>Baccharis</i>) <i>indica</i> ...	Kalsis ...	Common. Low elevations
2	" <i>argentea</i> ...	Phnsara ...	Frequent, Binsar.
3	<i>Myrsine africana</i> ...	Pahári chái, baiberang.	Common, March.
4	" <i>semiserrata</i> ...	Chupra ...	"
5	" <i>acuminata</i>	"
6	<i>Embellia robusta</i> ...	Baiberang ...	Bhábar.
7	<i>Ardisia floribunda</i>	Rare.
ORDER 13.—SAPOTACEÆ			
1	<i>Bouala butyracea</i> ...	Chíurá and phalcl.	Common at 4,000 feet, near Pctoragari.
2	" <i>latifolia</i> ...	Manús ...	Bhábar.
ORDER 14.—EBENACEÆ.			
1	<i>Diospyros Melanoxylon</i> ...	Tendu ...	Bhábar.
2	" <i>montana</i> ...	Tendu ...	"
3	" <i>lanceifolia</i> ...	Ardinia ...	Kota Dán
4	" <i>Embryopteris</i> ...	Gab ...	Bhábar.
ORDER 15.—SYMPLOCACEÆ.			
1	<i>Symplocos</i> (<i>Hopcia</i>) <i>salenta</i> ...	L. dli, bhú'ya.	Rare, November
2	" <i>acuminata</i>	Rare, Sial devi.
3	" <i>racemosa</i>	Bhábar, November.
4	" <i>ramosissima</i> (<i>corvosa</i> , Madden)	...	Rare, Gágar, Binsar.
5	" (<i>Pá'u'a</i>) <i>cratægoides</i> (S. pinnulata, Mullen).	Ladh ...	Very common, June and July.

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 16. OLEACEÆ.			
1	<i>Fraxinus excelsior</i> ...	Angu ...	Common.
2	" (<i>Ornus</i>) <i>floribunda</i> ...	Angu ...	Common.
3	<i>Syringa emodi</i> ...	Chin ...	Common, above 10,000 feet.
4	<i>Olea cuspidata</i> [ferruginea, Royle III.]	Kalap briksh,	Patāi dehi near Almorā.
5	" <i>acu inata</i> var. <i>longifolia</i> .	Silang ...	Very common, wild and planted.
6	" <i>glandulifera</i> ...	Gair ...	Common.
7	<i>Ligustrum nepalense</i>	Banks of streams, May.
8	" <i>obovatum</i>	Frequent, Binsar, June.
9	" <i>robustum</i>	Common, May.
10	" <i>compactum</i>	"
ORDER 17.—JASMINEÆ.			
1	<i>Jasminum</i> (<i>Simplexifolium</i>) <i>Sambac</i>	Cultivated.
2	" <i>pubescens</i>	Frequent, Canal.
3	" <i>arborescens</i>	" Bhābar.
4	" <i>nepalense</i> [<i>dichotomum</i> Don].	Jāhi ...	Common.
5	" (<i>trifolium</i>) <i>diapernum</i> ...	Sarmāl. ...	Common. Leaves, generally trifol, but sometimes simple and sometimes pinnate.
6	<i>Jasminum</i> (<i>Alternifolium</i>) <i>revolutum</i> [<i>chrysanthum</i> , Roxb.]	Sonajāhi ...	Common.
7	" " <i>pubigerum</i>	Rare.
8	" (<i>Pinnatifolium</i>) <i>orbiculare</i> , ...	Chambeli ...	Introduced.
9	" " <i>granuliflorum</i> , ...	Chambeli and jāhi.	Common.
10	<i>Nyctanthes Arbor-tristis</i> ...	Parijāt, harsinghār of plains.	
11	<i>Schrebera swietenoides</i> ...	Moka ...	Bhābar tree
ORDER 18.—APOCYNACEÆ.			
Tribe CARISSYÆ.			
1	<i>Carissa diffusa</i> ...	Timuktiya ...	Bhābar.
Tribe PLUMERIÆ.			
2	<i>Tabernaemontana coronaria</i> ...	Chandi ...	"
3	<i>Vinea pumila</i> [V. <i>parvidora</i> , Roxb.]	"
Tribe PASSIFLOREÆ.			
4	<i>Vallisneria spiralis</i> ...	Dūdhī ...	Common.
Tribe WRIGHTIÆ.			
5	<i>Wrightia multiflora</i> ...	Darhela ...	"

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 18.—APOCYNAC Æ (concluded).			
TRIBE ALSTONIEÆ.			
6	<i>Alstonia scholaris</i>	Bhābar.
7	<i>Blumea populifolia</i>	Dūdhi ...	Common.
TRIBE ECHITEÆ.			
8	<i>Holarrhena antidysenterica</i>	Kuer ...	"
9	" <i>pubescens</i>	"
10	<i>Nerium olerum</i>	Kanyūr ...	
11	<i>Conemorpha macrophylla</i> , [Echites macrophylla, Roxb.]	Garbadra ...	Bhābar.
12	<i>Ichnocarpus frutescens</i>	Dūdhi ...	Common.
ORDER 19.—ASCLEPIADEÆ.			
TRIBE PERIPLICEÆ.			
1	<i>Cryptolepis reticulata</i>	Rui kosa ...	Up to 4,000 feet.
2	<i>Periploca calophylla</i> [Streptocaulon calophyllum, Wright.]	Dāl bhengla,	Common.
TRIBE ASCLEPIADEÆ.			
3	<i>Vincetoxicum canescens</i> [Cynanchum glaucum, Wall.]	...	Common.
4	" <i>montanum</i> [V. kunu- warensis, Madden.]	...	Frequent.
5	<i>Caotroplepis gigantea</i>	Ak, madār ...	Common, Bhābar.
6	" <i>procera</i> [Hamiltonii]	" ...	Rare, Bhābar.
7	<i>Cynanchum lamhousei</i>	Frequent.
TRIBE STAFELIÆ.			
SUB-TRIBE PERGULARIÆ.			
8	<i>Tylophora tenerrius</i>	Frequent.
9	" <i>Govanii</i> [Iphisia Govani- ana, Wright.]
10	<i>Marasmiella Roylei</i>	Murkila ...	Common.
11	" <i>lucida</i> , Edgw.	Dūdhi ...	Frequent.
12	" <i>tenacissima</i>	Up to 4,000 feet.
13	<i>Pergularia pallida</i>	Sūrkila ...	Common, Bhābar.
14	<i>Gongronoma nepalense</i>	Frequent on rocks.
SUB-TRIBE CEROPEGIÆ.			
15	<i>Orthanthura viminea</i>	Chapkiya ...	Bhābar.
16	<i>Moya lanceolata</i>	Common, Bhābar, and up to 7,000 feet.
17	<i>Ceropegia longifolia</i>	Rare.
18	" <i>Wallitchii</i>	Frequent.
19	" <i>gracilis</i> , Edgw.	"

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 20.—LOGANIACEÆ.			
1	<i>Gardneria ovata</i>	Banjāhl ...	Frequent.
2	<i>angustifolia</i>	"
3	<i>Buddleia crispa</i> [paniculata, Roxb.]...	Dhirlīya, bhātī, dhaula.	Common.
4	<i>asiatica</i> [neoma, Hamilton],	...	"
ORDER 21.—GENTIANACEÆ.			
TRIBE CHIRONIÆ.			
1	<i>Exacum tetragonum</i>	1 Uda, tītkāna,	Frequent, Ragswar.
TRIBE CHLOREÆ.			
2	<i>Pladera pusilla</i>	Common.
3	<i>Canescora diffusa</i> [virgata, Roxb.]	Bhābar.
4	<i>decussata</i>	"
TRIBE SWERTICEÆ.			
5	<i>Gentiana</i> (<i>Chondrophyllum</i>) <i>pedicilata</i>	Common.
6	<i>capitata</i>	"
7	<i>aprica</i>	Bhābar.
8	<i>decussata</i>	Common.
9	<i>marginata</i>	"
10	<i>Ophelia paniculata</i>	Chiretta Tītkāna.	Autumn annual.
11	<i>purpureascens</i>	" ..	" "
12	<i>cordata</i>	" ..	" "
13	<i>var. floribus dilute roseis</i>	W. Waterfield, Naini Tāl.
14	(<i>Agathotes</i>) <i>angustifolia</i> ...	" ..	Autumn annual.
15	" <i>clirata</i>	" ..	" perennial.
16	" <i>alata</i>	" ..	" annual.
17	<i>Halenia elliptica</i>	Above 10,000 feet.
18	<i>Swertia cuneata</i>	Rare.
19	<i>caerulea</i>	"
20	<i>speciosa</i> [perfoliata, Royl.],	Simariya ...	Common, above 10,000 feet.
ORDER 22.—BIGNONIACEÆ.			
1	<i>Calceanthus indica</i>	Pharkath ...	Bhābar.
2	<i>Millingtonia hortensis</i> [<i>Bignonia suberosa</i> , Roxb.],	Nim chaubell	Introduced.
3	<i>Stereospermum</i> [<i>Bignonia</i> , Roxb.] <i>suaveolens</i> .	Pātal ...	Bhābar.
4	<i>Amphicomme arguta</i>	Banks of Patāiganga.
5	<i>Tecoma undulata</i>	Ganai valley, May.

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Notes.
ORDER 23.—PEDALIACEÆ.			
1	<i>Martynia proboscidea</i> [<i>M. diandra</i> , Madden].	...	Naturalised.
2	<i>Sesamum indicum</i> ...	Til ...	"
3	<i>Pedaliium murex</i>	Gokhrn ...	Bhābar.
ORDER 24.—CRYPTANDRACEÆ.			
1	<i>Lysionotus ternifolia</i> [<i>L. serrata</i> , Don].	...	Common up to 4,000 feet.
2	<i>Didymocarpus subalternans</i>	Rare.
3	" <i>aromatica</i>	Frequent.
4	" (<i>Acaules</i>) <i>macrophylla</i> ...	Pathar laung,	Common.
5	" <i>pedicellata</i>	Bhābar.
6	" <i>lanuginosa</i>	Common.
7	<i>Chirita Edgeworthii</i> ...	Sankh pushpa,	"
8	" <i>bifolia</i> ...	" ...	"
9	<i>Rhynchoglossum obliquum</i> [<i>Wu-fenia</i> <i>obliqua</i> , Wall]	...	" up to 6,000 feet.
10.	<i>Platystemma violoides</i>	Common.
11	<i>Æchynanthus perrottetiana</i>	Bhābar.
12	" <i>grandiflora</i>	Har silang. October.
13	<i>Polycnuron dimidiata</i>	Col. Davidson.
ORDER 25.—POLEMONIACEÆ.			
1	<i>Cyananthus lobata</i>	Pindari, above 10,000 feet.
2	" <i>integra</i>	Kailā 12,000 feet.
ORDER 26.—CONVOLVULACEÆ.			
TRIBE ARGYREÆ.			
1	<i>Rives ornata</i>	Bhābar.
2	<i>Argyrea speciosa</i>	"
3	" <i>setosa</i>	Sarju, bank near Kap- kut, September.
4	" <i>capitata</i>
TRIBE CONVULVEÆ.			
5	<i>Quamoclit vulgaris</i>	Bhābar.
6	<i>Batatas pentaphylla</i>	"
7	<i>Pharbitis Nil</i> ...	Baurā ...	Very common.
	var. <i>intergrifolia</i> [<i>cœrulea</i> , Roxb.]	" ...	Common.
8	<i>Calonyction speciosum</i>	Bhābar
9	<i>Ipomœa vitifolia</i>	"
10	" <i>Pœa-tigridis</i>	"
11	" <i>Pilosa</i>	Kota Dîn.
12	" <i>sp.</i> [<i>hirsuta</i> , Madden]	"

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 26.—CONVOLULACEÆ—(concluded).			
TRIBE CONVULVLEÆ—(concluded).			
13	<i>Ipomœa acutiflora</i> [pharoccephala, Roxb.]	...	Very common.
14	" <i>chrysoides</i> [muricata, Mad-den].	...	Bhābar.
15	<i>Arisœa barlerioides</i>	Kosi bank, Khaṛna.
16	<i>Convolvulus pluricaulis</i>	Bhābar
17	" <i>arvensis</i> [Malcolmi Roxb.]	...	Naini Tāl, Mr. Water-field.
18	<i>Porana racemosa</i> ...	Maeti ...	Common.
19	" <i>paniculata</i> ...	" ...	"
20	<i>Evolvulus alsinoides</i> [sericea, Madlen].	...	Very common.
TRIBE CUSANTEÆ.			
21	<i>Cuscuta reflexa</i> [verrucosa] ...	Akash leguli,	Common
22	" <i>macrantha</i> [grandiflora, Wall].	" ...	Frequent.
ORDER 27.—BORAGINACEÆ.			
TRIBE CORDIFÆ.			
1	<i>Gynœon vestitum</i> ...	Pyā ...	Common up to 3,000 ft.
2	<i>Cordia latifolia</i> , Roxb ...	Borā ...	Bhābar variety.
3	" <i>Myxa</i> , Linn. ...	Kodah ...	Common.
TRIBE EHRETICÆ.			
4	<i>Ehretia serrata</i> ...	Pānya ...	Common, April
5	" <i>lævis</i> ...	Kodah ..	" Bhābar, March.
6	" <i>viminea</i> [Rhabdia sericea Madden].	...	" in river beds.
TRIBE HELIOTROPÆ.			
7	<i>Heliotropium strigosum</i> [brevifolium Wall].	..	Common.
TRIBE BORAGÆ.			
SUB-TRIBE ECHINÆ.			
8	<i>Macrotomia Benthani</i> ...	1 ...	Badrināth.
SUB-TRIBE LITHOSPERMÆ.			
9	<i>Onosma echioides</i>	Above 10,000 feet.
10	<i>Myosotis pallens</i>	" " Chau-patta.
11	" <i>longiflora</i>	" " "
12	<i>Bothriospermum tenellum</i> [Cynoglossum prostratum, Don].	...	Rare, Bīnār.

DIVISION II.—MONOPETALÆ--(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 27.—BORAGINACEÆ—(concluded).			
SUB-TRIBE CYNOGLOSSÆ.			
13	<i>Eritrichium rotundifolium</i>	Above 10,000 feet; rare.
14	<i>Echinopspermum glochidiatum</i> (Cynoglossum uncinatum, Royle).	Kura ...	" " "
15	<i>Cynoglossum furcatum</i> ...	" ...	Very common.
16	" <i>micranthum</i> [canescens, Wall].	" ...	" "
17	" <i>glochidiatum</i> ...	" ...	Frequent.
18	<i>Trichodroma indicum</i>	Common.
ORDER 28.—SOLANACEÆ.			
TRIBE SOLANÆ.			
1	<i>Lycopersicon esculentum</i> ...	Wiláyati baigan.	Cultivated.
2	<i>Solanum</i> (<i>Pachystemon</i>) <i>tuberosum</i> ...	Ahi ...	"
3	" <i>Roxburghii</i> [rubrum] ...	Chhota ghivain.	Common.
4	" <i>verbascifolium</i> ...	Ushita ...	"
5	" (<i>Leptostemon</i>) <i>indicum</i> ...	Kant kárl ...	"
6	" <i>xanthocarpum</i>	"
	var. <i>Jacquini</i>	"
	var. <i>Schraderei</i> [diffusum]	"
7	" <i>esculentum</i> [melongena], ...	Baigan ...	Cultivated.
8	<i>Capsicum frutescens</i> ...	Lál uirch ...	"
9	<i>Nicantra physaloides</i>	Introduced.
10	<i>Physalis peruviana</i>	"
TRIBE DATURÆ.			
11	<i>Datura Wallichii</i> [ferox, Makden] ...	Kála dhatúra, ...	Valleys.
12	" var. <i>double</i>	Cultivated.
	" <i>alba</i> ...	Dhatúra ...	Very common.
	" var. <i>double</i>	Cultivated.
TRIBE NICOTIANÆ.			
13	<i>Nicotiana Tabacum</i> ...	Tamáku ...	Cultivated.
14	<i>Petunia Phœnicea</i>	Introduced.
ORDER 29.—SCROPHULARIACEÆ.			
TRIBE ANTIRRHINÆ.			
1	<i>Verbascum Thapsus</i> ...	Ekalbír ...	Common.
2	<i>Celsia coromandeliana</i>	Rare, valleys.
3	<i>Linaria ramosissima</i>	Common.
4	" <i>incana</i>	"
5	<i>Antirrhinum orontium</i>	" May fields.
6	<i>Scrophularia Edgeworthii</i>	Above 12,000 feet.

DIVISION II.—NONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 29.—SCROPHULARIACEÆ—(continued)			
Tribe ANTIRRHINEÆ—(concluded).			
7	<i>Scrophularia polyantha</i> ...	Gajalla ...	Common.
8	" <i>himalensis</i> ...	" ...	"
9	" <i>obtusa</i> (Madden) ...	" ...	Frequent.
10	<i>Mimulus nepalensis</i> ...	" ...	Common.
11	" <i>gracilis</i> ...	" ...	Frequent.
12	<i>Mazus surculosus</i> ...	" ...	Common.
13	" <i>rugosus</i> ...	" ...	"
14	<i>Lindenbergia grandiflora</i> ...	" ...	"
15	" <i>urticifolia</i> [Stemodia ruderalis, Wall].	" ...	Very common.
16	" <i>macrontachya</i> ...	" ...	Rare.
17	<i>Limonophila menthustrum</i> ...	Jangli sonf .	Common.
18	" <i>hypericifolia</i> ...	" ...	"
19	<i>Herpertia Monnieria</i> ...	Jāl nīm ...	"
20	<i>Torenia cordifolia</i> ...	" ...	"
21	<i>Vandellia crustacea</i> [pedunculata, Madden].	" ...	"
22	" <i>nummularifolia</i> ...	" ...	Very common. August.
23	<i>Bonnaya brachiata</i> ...	" ...	Frequent. September.
24	" <i>grandiflora</i> ...	" ...	"
Tribe RHINANTHEÆ.			
25	<i>Hemiphragma heterophyllum</i> ...	" ...	Common at 10,000 feet, Dākuri and Dýáritál.
26	<i>Digitalis purpurea</i> ...	" ...	Introduced.
27	<i>Wulfenia Antheratiana</i> [Pæderota Antheratiana, Wall].	" ...	Common near water above 8,000 feet.
28	<i>Veronica</i> (Beccabunga) <i>himalensis</i> [grandiflora, Wall].	" ...	Above 10,000 feet.
29	" <i>anagallis</i> ...	" ...	Up to 5,000 feet.
30	" <i>punctata</i> ...	" ...	"
31	" (<i>Chamaedrya</i>) <i>deltoidea</i> ...	" ...	Common.
32	" <i>laxa</i> ...	" ...	Above 11,000 feet.
	" <i>var. of V. chamaedrya?</i> ...	" ...	Kedarnáth.
33	" (<i>Onphalospora</i>) <i>biloba</i> ...	" ...	Common at 5,000 feet. March.
34	" " <i>Maddenii</i> Edge. ...	" ...	"
35	" " <i>agrestis</i> ...	" ...	"
36	<i>Buchnera hispida</i> ...	" ...	"
37	<i>Striga hirsuta</i> ...	" ...	"
...	" <i>var. lutea</i> ...	" ...	"
38	" <i>euphrasoides</i> ...	" ...	"
39	<i>Leptorhabdos parviflora</i> [Gerardia and <i>Sopubia parviflora</i>].	" ...	"
40	<i>Sopubia trifida</i> ...	" ...	Grassy slopes at 8,000 feet.
41	<i>Centranthera hispida</i> ...	" ...	Bagewar. Frequent.
...	" <i>var. nepalensis</i> ...	" ...	"
42	<i>Euphrasia officinalis</i> ...	" ...	Above 7,000 feet.

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 29.—SCHOPHULARIACEÆ—(concluded).			
TRIBE RHINANTHEÆ—(concluded).			
43	<i>Pedicularis (Cyclophyllum) pectinata.</i>	...	Frequent.
44	" " <i>porrecta,</i>	..	"
...	" [<i>elegans</i> , Madden]	..	"
45	" (<i>Siphonanthus</i>) <i>megalantha.</i>	..	Above 10,000 feet.]
46	" (<i>Edentula</i>) <i>carnea</i>	...	Common.
ORDER 30.—OROBANCHACEÆ.			
1	<i>Phellipsea aegyptiaca</i>	...	Rare.
2	<i>Æginetia indica</i>	...	Frequent, Bagewar.
ORDER 31.—ACANTHACEÆ			
TRIBE THUNBERGIEÆ.			
1	<i>Hexacentris coccinea</i>	Kuljug	Bhābar and valleys. Winter.
TRIBE HYGROPHILLEÆ.			
2	<i>Hemiladelphis polyasperma</i>	.. 1	Bhābar.
TRIBE RUELLEÆ.			
3	<i>Petalidium barlerioides</i>	Bakraola	...
4	<i>Hemigraphis latebrosa,</i> [<i>Ruellia latebrosa</i>].	...	Common.
5	<i>Æchmanthera Wallichii</i> [<i>gossypina</i> , Madden].	Jaundela	Very common. From this bees make the best honey.
6	<i>Goldfussia pentstemonoides</i>	...	Frequent.
7	<i>Strobilanthes auriculatus</i>	Tilkapuri	Bhābar.
8	" <i>Wallichii</i>	...	Above 10,000 feet.
9	" <i>attenuatus</i>	Kapur nali	Common.
10	" <i>glutinosus</i>	...	"
TRIBE BARLERIEÆ.			
11	<i>Barleria dichotoma</i>	...	Frequent, September.
12	" <i>nepalensis</i>	...	"
13	" <i>ciliata</i> (rosea)	Sarp jiba	Common.
14	" <i>cristata</i>	...	"
15	<i>Lepidagathis hyalina</i>	...	"
16	" <i>undulata</i>	...	About 4,000 feet.
17	" <i>cuspidata</i>	...	Common.

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 31.—ACANTHACEÆ—(concluded).			
Tribe JUSTICEÆ.			
18	<i>Phlogacanthus thyrsoiflorus</i>	Bhābar and valleys. common. March
19	<i>Kranthemum nervosum</i>	Bhābar and up to 6,000 feet. May.
20	<i>Rostkilmaria procumbens</i>	Very common.
21	<i>Adhatoda vasica</i> [<i>Justicia Adhatoda</i>],	Bashang. Arūn of plains.	Common.
22	<i>Rangia parviflora</i>	"
23	<i>Dicliptera Roxburghii</i>	" up to 5,000 feet.
24	<i>Periatrophe speciosa</i>	Bhābar.
25	" <i>bicalyculata</i>	Rare, Vishnuganga.
26	<i>Rhaphidospora ciliata</i>	
ORDER 32.—PHYRMACEÆ.			
1	<i>Phryma leptostachya</i> ... 1	...	Common in woods.
ORDER 33.—VERBENACEÆ.			
Tribe VERBENEÆ.			
1	<i>Verbena Bonariensis</i>	Naturalized.
2	" <i>officinalis</i>	Common.
3	<i>Lippia (Zapanaia) nobiliflora</i>	Up to 4,500 feet.
4	<i>Lantana alba</i> [L. dubia, Royle III.]	...	" 3,000 "
5	<i>Gmelina arborea</i> ..	Kumār	" 2,000 "
Tribe VITEÆ.			
6	<i>Caryopteris Wallichiana</i> [<i>Clerodendron odoratum</i> , Muell. Ar.]	Mohini	Very common up to 6,000 feet. April, Bhābar.
7	<i>Premna (Gumina) serratifolia</i> [spinosa, Roxb.]	...	
8	" (<i>Premna</i>) <i>mucronata</i> ..	Agnūn	"
9	" " <i>burhata</i>	Common.
10	" " <i>herbacea</i>	Rare.
11	<i>Callicarpa arborea</i> , Roxb. ..	Ghiwāl	Common.
12	" <i>macrophylla</i> , Vahl	Daiya	Frequent, Sarju bank.
13	<i>Clerodendron serratum</i>	Common.
	" <i>var. ternatifolium</i>	" Bageswar.
14	" <i>infundatum</i>	Bhābar.
15	" <i>siphonanthus</i>	Bhābar and outer hills, April.
16	<i>Vitex negundo</i> ..	Shiwāl	
17	<i>Holuskoldia sanguinea</i> ..	Kubtolliya	Common, October.
ORDER 34.—LABIATÆ.			
Tribe OCIMEÆ.			
1	<i>Geniospermum strobiliferum</i>	Banks of Sarju.
2	<i>Ortholipon rubicundus</i>	Common.
3	" <i>virgatus</i>	Banks of Kosi.
4	<i>Plectranthus (Isodon) scrophularioides</i> ,	...	Frequent.
5	" " <i>Geraniifolius</i>	"
6	" " <i>striatus</i>	Very common October.

DIVISION II.—MONOPETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 24.—LABIATÆ—(continued).			
TRIBE QUINÆÆ—(concluded).			
7	<i>Plectranthus (Isodon) Coctus</i>	Very common. Octr.
8	" <i>rugosus</i>	Common up to 6,500 ft.
9	" (<i>Pyramidium</i>) <i>ternatifolius</i> ,	" low elevations.
10	" (<i>Colcoides</i>) <i>cordifolius</i>	"
11	<i>Coleus barbatus</i> ...	Piwal ...	Very common.
TRIBE SATYRÆ.			
SUB-TRIBE ELIALTHIÆ.			
12	<i>Pogostemon plectranthoides</i> ...	Rudra ...	Common.
13	<i>Dysophylla cruciata</i>	" in swamps.
14	<i>Colobrochia oppositifolia</i> ...	Dulshat ...	"
15	<i>Elialthia flava</i> ...	Bhangria ...	" down to 5,000 feet.
16	" <i>polystachya</i> ...	Bhangria ...	Very common.
17	" <i>incisa</i> ...	Set bhangria.	Rare.
18	<i>Elialthia pilosa</i>	Frequent.
19	" <i>crisostachya</i>	Rare.
20	" (<i>Cyclostegia</i>) <i>strobilifera</i>	Common.
SUB-TRIBE MENTHÆ.			
21	<i>Perilla ocimoides</i> ...	Bhangria ...	Cultivated, September.
22	<i>Mentha viridis</i> ...	Pulina ...	Naturalized.
23	" <i>Royleana</i>	Rare. Near Bhim Tal.
SUB-TRIBE THYMÆ.			
24	<i>Origanum normale?</i> [vulgare?] ...	Ban tulsi ...	Very common, autumn.
25	<i>Thymus scripyllum</i> ...	" ajwain ...	Common, except outer range.
SUB-TRIBE MELLISÆ.			
26	<i>Micromeria biflora</i>	Very common.
27	<i>Calamintha (Clinopodium) repens</i>	"
28	" <i>umbrosa</i> [Melissa,	June, under trees.
29	<i>Hedeoma nepalensis</i>	Frequent.
TRIBE MOKARDÆ.			
30	<i>Salvia (Drymosphace) glutinosa</i> [hians, Royle.]	Frequent, outer range.
31	<i>Salvia (Aethiops) lanata</i> ...	Gariya ...	Very common, March.
32	" <i>Mourcroftiana</i>	Above 10,000 feet.
33	" (<i>Notiosphace</i>) <i>plebeia</i>	Bhabar and valleys.
TRIBE NEPITÆ.			
34	<i>Nepeta (Pycnonopeta) spicata</i> , var.	Common, Binaar.
	" <i>elata</i> , Royle	Yellow, morning, blue afternoon.
35	" <i>elliptica</i>	Common, Binaar.
36	" (<i>Cataria</i>) <i>rupestris</i>	Frequent.
37	" <i>leucophylla</i>	Common.
38	" <i>graciliflora</i>	Bhabar.
39	" (<i>Macronepeta</i>) <i>goraniana</i>	Frequent.
40	<i>Dracopis alpinum acanthoides</i>	Niti.
41	<i>Loliumantia Royleana</i> ...	Balangra ...	Rare.

DIVISION II.—MONOPETALÆ—(concluded).

No.	Scientific name.	Vernacular name.	Note.
ORDER 34.—LABIATÆ—(concluded).			
TRIBE STACHTYDÆ.			
SUB-TRIBE SCUTELLARIÆ.			
42	<i>Prunella vulgaris</i>	Common, June, Binsar.
43	<i>Scutellaria linearis</i>	" spring.
44	" <i>repens</i>	" spring and autumn.
45	" <i>angulosa</i> [scandens, Don].	...	" autumn.
SUB-TRIBE MARRUBIÆ.			
46	<i>Craniotome versicolor</i>	Very common.
SUB-TRIBE LAMIÆ.			
47	<i>Anisomeles ovata</i>	Common.
48	<i>Colquhounia coccinea</i> [vestita, Wall]	Bhilmora and angiriya.	" at Naini Tāl and Badrināth.
49	<i>Stachys sericea</i>	Common.
50	" <i>splendens</i>	Bhainsia aj-wāin.	Rare.
51	<i>Leonurus pubescens</i>	Common.
52	<i>Lamium amplexicaule</i>	"
53	" <i>petiolatum</i>	Tit pāti ...	"
54	<i>Royia elegans</i>	Guma ...	Frequent.
55	<i>Leucas</i> [Hemistoma] <i>urticaefolia</i>	Common.
56	" [Ortholeucas] <i>lanata</i>	Rare.
57	" " <i>mollissima</i>	Common.
58	" [Plagiostoma] <i>hysopifolia</i> ,	"
59	" " <i>aspera</i> [cephalotes, Blume.]	...	Bhābar & Haldwāni.
60	<i>Leonotis nepetæfolia</i>	Rare.
61	<i>Phlomis lamellifolia</i>	
TRIBE AJUGÆ.			
62	<i>Teucrium Royleanum</i> (var.) <i>albidum</i>	Common, valleys.
63	" <i>quadrifarium</i>	Bhābar.
64	<i>Ajuga lobata</i>	Very common.
65	" <i>remota</i>	High elevations.
66	" <i>parviflora</i>	Very common.
67	" <i>bracteosa</i>	Rare.
68	" <i>decumbens</i>	
ORDER 35.—PLUMBAGINACÆ.			
69	<i>Plumbago Zeylanica</i>	Common.
ORDER 36.—PLANTAGINACÆ.			
1	<i>Plantago</i> (<i>Polyneuron</i>) <i>major</i> , var. <i>asiatica</i> .	Bār tang ...	Rare.
2	" " <i>erosa</i> , Don, [longica, Jacq.]	Lohuriya ...	Very common.
3	" " <i>lanceolata</i>	Rare.

DIVISION III.—APETALÆ.

No.	Scientific name.	Vernacular name.	Note.
ORDER 1.—PHYTOLACCACIÆ.			
1	<i>Phytolacca, decandra, var. acinosa, ...</i>	Jirrag ...	Cultivated up to 10,000 feet.
ORDER 2.—CHENOPODIACÆ.			
1	<i>Beta vulgaris, var. Bengalensis, Roxb.,</i>	Palang ...	Cultivated.
2	<i>Chenopodium album ...</i>	Bhatuwa ...	Common.
3	" (<i>Ambrina</i>) <i>Botrys ...</i>	...	"
ORDER 3.—AMARANTACÆ.			
TRIBE CELOSIFÆ.			
1	<i>Deeringia celosioides ...</i>	Kāla lohāri...	Common.
2	<i>Colocasia argentea ...</i>	Sirāl a n d ghogiya.	"
TRIBE ACHYRANTHÆ.			
3	<i>Amarantus caudatus ...</i>	...	Gardens.
4	" <i>anardana [farinaceus] ...</i>	Ramdāna and chua.	Cultivation.
5	" <i>spinosus ...</i>	Sāji ...	Common.
6	<i>Acris scandens ...</i>	Sāji ...	"
7	<i>Achyranthes aspera ...</i>	...	Not known at Kew
8	" species [<i>bidentata</i> ?],	Very common.
9	<i>Cyathula tomentosa [Fupalis sequax],</i>	Jhat kuri ...	
	Madden.	...	Rare.
10	" <i>capitata.</i>	...	
TRIBE GOMPHRENEÆ.			
11	<i>Alternanthera nodiflora ...</i>	Bhimrāj ...	Common. Bhim Tāl.
ORDER 4.—NYCTAGINACÆ.			
1	<i>Borhaavia diffusa ...</i>	...	Common.
2	<i>Mirabilis Jalapa ...</i>	...	Introduced.
ORDER 5.—POLYGONACÆ.			
TRIBE TRIPOCARPÆ.			
1	<i>Rheum emodi ...</i>	Dolu ...	Above 10,000 feet.
2	<i>Oxyria elatior ...</i>	...	" 9,000 "
TRIBE APTEROCARPÆ.			
3	<i>Rumex (Lapallum) Wallibohli ...</i>	...	Bhābar.
4	" <i>nepalensis ...</i>	Haluniya ...	Common.
5	" (<i>Acetosia</i>) <i>hastatus ...</i>	Almora ...	Very common.
6	<i>Kenigia moandra or islandica ...</i>	...	Above 12,000 feet.
7	<i>Polygonum (Avicularia) effusum</i>	...	Common.
	[<i>herniarioides</i> .]	...	
1	" <i>Roxburghii ...</i>	...	Rare.
2	" <i>ilicobroides ...</i>	...	"

DIVISION III.—APETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 5.—POLYGONACEÆ—(concluded).			
Tribe ATEROCARPEÆ (concluded).			
10	<i>Polygonum ellipticum</i>	Rare.
11	" <i>cornutum</i>	14,500 feet.
12	" <i>perumbena</i>	Common. September.
13	" (<i>Periscaria</i>) <i>Hamiltonii</i> [<i>hiapidum</i> , Don.],	...	Frequent. Bhábar, November.
14	" <i>hirsutum</i>	Bhábar. Common.
15	" <i>donii</i>	Frequent.
16	" <i>posumbu</i>	Bhábar.
17	" <i>flaccidum</i> [<i>glabrum</i> , Roxb.],	...	
28	" <i>amphibium</i>	Naini Tál lake.
19	" <i>lanigerum</i>	Frequent.
20	" <i>scabrinervium</i>	Naini Tál lake.
21	" (<i>Bistorta</i>) <i>viviparum</i> [<i>bulbiferum</i> , Royle].	...	15,000 feet.
22	" <i>sphaerostachyum</i>	Very common.
23	" <i>amplexicaule</i>	Common.
24	" <i>affine</i> [<i>Brunonia</i> , Wall.],	...	At high elevations.
25	" <i>enodii</i>	Rare.
26	(<i>Didymocarpum</i>) <i>delicatulum</i>	12,000 feet.
27	" <i>nummifolium</i> ,	14,500 feet.
28	" <i>stipulae</i> [<i>ciliatum</i> , Don.].	...	Rare.
29	" <i>humile</i>	"
30	" <i>perforatum</i> [<i>punctatum</i> , Don.].	...	"
31	" <i>nepalense</i>	Common.
32	" <i>sphaerocarpum</i> ,	Frequent.
33	" <i>capitatum</i> [<i>repens</i> , Wall.],	...	Very common.
34	" <i>sinuatum</i>	8,500 feet.
35	" (<i>Corymbocarpum</i>) <i>chinense</i> , (<i>Rhinocarpum</i>) <i>horridum</i>	Very common.
36	" (<i>Tiniaria</i>) <i>convolutum</i>	Bhím Tál. Common.
37	" <i>pterocephalum</i>	"
38	" (<i>Aconogonon</i>) <i>frondosum</i>	7,500 feet.
39	" <i>polystachyum</i>	12,000 feet.
40	" <i>Fagopyrum esculentum</i> [<i>vulgare</i> , Madden].	Ogal ...	Cultivated, lower hills.
41	" <i>emarginatum</i>	Pápag ...	Higher hills.
42	" <i>cymosum</i> [<i>corymbosum</i> , Madden].	Ban-ogal ...	Common. The native names given by Royle are incorrect.
ORDER 6.—LAURINEÆ.			
SECTION EXINVOLUCRATÆ.			
1	<i>Cinnamomum Tamala</i> , var. <i>albiflorum</i> ,	Tejpát, dáchhi- ni.	Common. February— May.
2	<i>Phoebe lanceolata</i>	Sau kanwal...	Bhábar and valleys, February—May.
3	" <i>pallida</i>	Kanwal ...	Common.
4	" <i>paniculata</i>	Kapua kanwal,	Bhábar and valleys.
5	<i>Machilus odoratissima</i>	Common. March— August.

DIVISION III.—APETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 6.—LAURINÆ—(concluded).			
SECTION INVOCRATEÆ.			
7	<i>Tetranthera laurifolia</i> [apetalæ, Koch., var. <i>Roxburghii</i>].	Gar-bijur ...	Valleys. June.
"	<i>monopetala</i> ...	Katunara, singrao.	Common. April.
8	<i>Litsea innuclonæ</i>	Rare.
9	" <i>conculilla</i> [T. pallens, Don.].	Chirra and chirchira.	Common, interior of hills.
10	<i>Daphnidium pulcherrimum</i>	Frequent.
11	" <i>bifarium</i>	"
ORDER 7.—THYMELACÆÆ.			
1	<i>Daphne papyracea</i> [D. cannabina, Wall.],	...	Common.
"	var. <i>purpurea</i> ...	Baruwa ...	
"	" <i>alba</i> ...	Set baruwa...	Used, for paper-making.
2	<i>Wikstrœmia salicifolia</i> [Daphne sericea, Madden]	Chunliya ...	Common, used for making inferior paper.
ORDER 8.—ELEAGNACÆÆ.			
1	<i>Hippophae salicifolia</i> ...	Dur chuk ...	Common above 10,000 feet.
2	<i>Elœagnus parvifolia</i> ...	Chiwain ...	Common.
3	" <i>latifolia</i> [ferruginea, D.C.]	...	Bhâbar.
ORDER 9.—MANTALACÆÆ.			
1	<i>Oxyris arborea</i> , [nepalensis, Madden.],	Bâkar dhârâ,	Common.
2	<i>Thesium himalense</i>	Rare, Bhâbar.
ORDER 10.—ARISTOLOCHIACÆÆ.			
1	<i>Aristolochia succata</i>	Frequent; about 7,000 feet.
ORDER 11.—EUPHORBIACÆÆ.			
TRIBE PHYLANTHINÆ.			
1	<i>Amirachne corallifolia</i> [Leptopus corallifolius, Decr.]	...	Very common.
2	<i>Anticarsia diandra</i> ...	Sarshoti âmlî	Bhâbar.
3	<i>Phyllanthus</i> [Hemiglochidion] nepalensis [Bradleya ovata, Wall. Glochidion bifaria, Royle].	Bair mao
"	" <i>velutinus</i> ...	" " ...	Very common.
4	" (Emblia) Emblica ...	Amia ...	Frequent.
5	(Paraphyllanthus) urinaria ...	Slâhl ...	Common.
6	(Ruphyllanthus) parvifolius	Frequent.
7	<i>Putorijiva</i> Roxburghii ...	Putrâ jivâ ...	Bhâbar.
8	<i>Securidaca</i> (Fluggen) obovata [Phyllanthus retusus, Wall.]	Dhâni ...	Common.
9	" <i>leucopyrus</i> [Phyllanthus albicans, Wall.]	Ainta
10	<i>Blachodis Zeylanica</i> [Andrachne trifoliata, Roxb.].	Koraa

DIVISION III.—APETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 11.—EUPHORBIACEÆ—(concluded).			
TRIBE BRIDELIÆ.			
13	<i>Bridelia stipularis</i>	Rare.
14	" (<i>Cluytia</i>) <i>montana</i>	Karnaliya ...	Common.
14	" <i>retusa</i>	Khaja ...	Up to 3,000 feet.
TRIBE ACALYPHÆ.			
16	<i>Frewia nudiflora</i>	Támri ...	Bhábar.
16	<i>Mallotus philippinensis</i> [<i>Rettlera tinctoria</i>].	Ryáni ...	Very common.
17	<i>Ricinus communis</i>	Ind, rendi ...	Common.
18	<i>Homonoya riparia</i> [<i>Adelia</i> , Madden].	Kandágar ...	Frequent.
TRIBE HIPPOCAMPÆ.			
19	<i>Jatropha Curcas</i>	Safed ind ...	Naturalized, Bhábar.
20	<i>Baliospermum montanum</i> [<i>Croton polyanthus</i> , Roxb.]	...	Common, Sarju bank.
21	<i>Euxcarica</i> (<i>Triadica</i>) <i>s. bifera</i>	Introduced.
22	" (<i>Euxcarica</i>) <i>insignis</i>	Common.
23	" (<i>Euxcarica</i>) <i>acrifolia</i> [<i>sapium</i> , Madden, <i>Stillingia himalayensis</i>]	Phutkiya ...	Very common, Sarju bank.
TRIBE EUPHORBIEÆ.			
24	<i>Euphorbia pentagona</i> [<i>Royleana</i>] ...	Sohund ...	Up to 6,000 feet.
2	" <i>angustifolia</i> [<i>dracunculoides</i> , Roxb.]	Dúdhila and Mahábir.	Common up to 6,000 feet.
26	" <i>hirsuta</i> [<i>longifolia</i> , Don.],	"	Above 6,000 feet (like <i>Helioscopia</i>).
27	" <i>peplodes</i>	Frequent.
28	" <i>pilulifera</i>	Very common.
29	" <i>notoptera</i>	Ganai valley.
30	" <i>hypericifolia</i>	Col. Davidson.
31	" <i>thymifolia</i>	"
32	" <i>pilosa</i>	"
ORDER 12.—DAPHNIPHYLLACEÆ.			
1	<i>Daphniphyllum himalense</i> [<i>Gyrantra laurina</i> , Wall. <i>Illicium</i> <i>sp.</i> , Royle.]	Rakt bhandan rakt angliya.	Frequent. Used in marking the tikú mark on forehead.
ORDER 13.—BUXACEÆ.			
1	<i>Buxus sempervirens</i>	Papar ...	Milam and Wán.
2	<i>Sarcococca saligna</i> Dec. [<i>S. nepalensis</i> , Royle].	Sukat sing...	Common, high levels.
ORDER 14.—CANNABINACEÆ.			
1	<i>Humulus lupulus</i>	Introduced.
2	<i>Cannabis sativa</i>	Bhang ...	Very common.

DIVISION III.—APETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 15.—ULMACEÆ.			
1	<i>Ulmus Wallichiana</i> [cross, Roth] ...	Chambar mā-ya.	Frequent. Flowers precocious.
2	" <i>parvifolia</i> [virgata, Roxb.] ...	"	Frequent. Flowers coetaneous.
3	<i>Holoptelea integrifolia</i> [<i>Ulmus integrifolia</i> , Roxb.]	Kanjo	Common.
4	<i>Celtis australis</i> [tetrandra., Roxb] ...	Kharak	Very common and planted.
5	<i>Spondia politoria</i> [<i>Celtis tetrandra</i> , Wall ?]	Khaoi, khāk-si.	Bhābar and valleys.
ORDER 16.—MORACEÆ.			
1	<i>Strobilina aspera</i> [<i>Trophis aspera</i> ...]	Rosa	Bhābar.
2	<i>Plecospirium spinosum</i> [<i>Batis spinosa</i>]	"	Rare.
3	<i>Morus indica</i> ...	Shalitūt	Frequent.
4	" <i>laevigata</i> ...	"	"
5	" <i>serrata</i> ...	Kimn	Common. In mountains.
6	<i>Ficus indica</i> ...	Bar	Valleys.
7	" <i>infectoria</i> ...	Pākar	Bhābar
8	" <i>Talicta</i> ...	Kathbar	Planted.
9	" <i>religiosa</i> ...	Pipal	Valleys.
10	" <i>cordifolia</i> ...	Kabul pipal,	Common.
11	" <i>retusa</i> [nitida] ...	"	Bhābar.
12	" <i>Carica</i> ...	Anjir	Cultivated.
13	" <i>virgata</i> ...	Beru	Very common.
14	" <i>parasticta</i> ...	"	Bhābar.
15	" <i>scandens</i> ...	"	Valleys.
16	" <i>trachycarpa</i> ...	"	"
17	" <i>Cnna</i> ...	Kunla	Common.
18	" <i>glomerata</i> ...	Gulār	Planted and banks of Kali.
19	" <i>macrophylla</i> ...	Timla	Fruit eaten.
20	" <i>oppositifolia</i> ...	Totmilla	Common.
21	" <i>Laducea</i> ...	Kabhila	"
22	" <i>laurifolia</i> ...	Dadhila	"
23	" <i>ovata</i> ...	Betull	"
24	" <i>Chincha</i> ...	Kiamira	"
25	" <i>saxatilis</i> ...	"	"
26	" <i>laminosa</i> ...	"	"
27	" <i>acuminata</i> ...	"	Bhābar.
ORDER 17.—ARTOCARPEÆ.			
1	<i>Artocarpus Lakoocha</i> ...	Dhāo	Introduced.
2	<i>Cudrania javanensis</i> ...	Mādu	Bhābar.
ORDER 18.—URTICACEÆ.			
Tribe URENEÆ.			
1	<i>Urtica parviflora</i> ...	Shishama	Eaten; very common.
2	<i>Girardinia heterophylla</i> ...	Awā bichhu.	Very common.

DIVISION III.—APETALÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 18.—URTICACEÆ—(concluded).			
TRIEF PRACERIDÆ.			
3	<i>Pilea scripta</i> [sagado, Wall.]	Common; Naini Tāl.
4	<i>Lecanthus peduncularis</i> ...	Saochāla ...	Very common; July, August.
5	<i>Elatostemma acule</i> [<i>Procris punctata</i> , Don].	" ...	"
6	" <i>platyphyllum</i> ...	Gargela ...	Common.
7	" <i>Stracheyanum</i>	Rare.
8	" <i>diversifolium</i>	Common; August.
TUM. BOEHMERIÆ.			
9	<i>Boehmeria rugulosa</i> [nervosa, Madden]	Geti ...	Common; dishes made of the wood.
10	" <i>macrophylla</i> ...	Saochāla golka	Common.
11	" <i>platyphylla</i> , var. <i>rotunda</i> folia.	Gargela ...	"
12	<i>Chamaetina squamigera</i>	Bhābar.
13	<i>Pouzolzia viminea</i> [sanguinea]	Frequent.
14	<i>Memorialis pentandra</i> [<i>Urtica pentandra</i> , Koch.	Jaiphal jari,	Common.
15	" <i>hirta</i>	Frequent.
16	" <i>quinque nervia</i>	"
17	<i>Villebrunea frutescens</i> , [<i>Urtica frutescens</i> , Madden]	Poi dhaula and kāzāhl.	Common.
18	<i>Debregeusia longifolia</i>	Rare.
19	" <i>bicolor</i> [<i>Boehmeria salicifolia</i> , Madden]	Tushyārī ...	Very common; April.
20	<i>Maoutia Puya</i> , [<i>Boehmeria frutescens</i> , Don.]	P'husar patta and dhaula-kāzāhl.	Common up to 4,000 feet; Bhim Tāl and Bageswar.
ORDER 19.—PIPERACEÆ.			
1	<i>Piper sylvaticum</i> ...	Pippula mūl,	Common; Bhābar and valleys.
2	<i>Piperomia reflexa</i> . [saxatilis, Mad- den].	" ...	Very common on trees.
ORDER 20. CORYLACEÆ.			
1	<i>Carpinus faginea</i> ...	Gish cham- kharak.	Common.
2	" <i>vilina</i>	"
3	<i>Corylus lacra</i> ...	Kapāl, bho- tiya ba lān.	Above 10,000 feet. Common, Jōshi-math.
4	<i>Castanopsis tribuloides</i> ...	Kātonj ...	Frequent.
5	<i>Quercus incana</i> ...	Bānj ...	Very common.

DIVISION III.—APETALÆ—(concluded).

No.	Scientific name.	Vernacular name.	Note.
ORDER 20.—CORYLACEÆ—(concluded).			
6	<i>Quercus lanata</i>	Blanj ...	Frequent; Lariya kán-ta.
7	" <i>semicarpifolia</i>	Karahn ...	Common, high elevations.
8	" <i>dilatata</i>	Tilonj ...	Very common.
9	" <i>annulata</i>	Phalyant ...	Common, valleys.
ORDER 21.—JUGLANDACEÆ.			
1	<i>Juglans regia</i>	Akhrot, ákor	Common.
2	<i>Engelhardtia Colebrookiana</i> ...	Mao ...	"
ORDER 22.—MYRICACEÆ.			
1	<i>Myrica aspl'a</i>	Kaiphai ...	Very common.
ORDER 23.—BETULACEÆ.			
1	<i>Betula Bhojpatra</i>	Bhojpatra ...	Above 10,000 feet.
2	" (<i>Betula</i>) <i>acuminata</i>	Itaoul ...	" 8,000 "
	[<i>cylindrostachys</i>]	Shinoul ...	Common.
3	" <i>nitida</i>	Pnya-ulash,	"
4	<i>Alnus nepalensis</i>	Udash ...	Banks of all rivers.
ORDER 24.—SALICACEÆ.			
1	<i>Salix</i> (<i>Picean</i>) <i>tetrasperma</i> ...	Gar-bains ...	Common near streams.
2	" " <i>acutophylla</i> [<i>dealbata</i> , Jacq.]	Frequent.
3	" (<i>Diandra</i>) <i>babylonica</i> ...	Majnuv ...	Introduced.
4	" " <i>Wallichiana</i> ...	Bains ...	Common.
5	" " <i>elegans</i> [kumaonensis, Lindley]	Top of China.
6	" " <i>daphnoides</i>	Above 10,000 feet.
7	" " <i>Lindleyana</i>	"
8	<i>Populus nigra</i> [<i>fastigiata</i>]	Introduced.
9	" <i>ciliata</i>	Gar pipal ...	Common.
ORDER 25.—LORANTHACEÆ.			
1	<i>Viscum stellatum</i> , Don. ...	Bánda ...	Common on <i>Pyrus</i> .
2	" (<i>Aphylla</i>) <i>articulatum</i> ...	Ditto ...	Common on <i>Quercus</i> , <i>incana</i> and <i>dilatata</i> .
3	" <i>attenuatum</i>	Ditto ...	Ditto.
4	<i>Loranthus</i> (<i>Euloranthus</i>) <i>lignastrinus</i> ,	Frequent; June.
5	" (<i>Scurrula</i>) <i>pulverulentus</i> ,	Common; winter.
6	" " <i>cordifolius</i>	Rare; December.
7	" " <i>vestitus</i>	Frequent; winter.
8	" (<i>Pentamera</i>) <i>longiflorus</i> [bicolor, Roxb.]	Common; spring.

DIVISION IV.—GYMNOSPERMÆ.

No.	Scientific name.	Vernacular name.	Note.
ORDER 1.—GNETACEÆ.			
1	<i>Ephedra vulgaris</i> [Gerardiana]	Above 10,000 feet ; fruit edible.
ORDER 2.—CONIFERÆ.			
1	<i>Pinus longifolia</i> ...	Chir, salla ...	Very common.
2	" <i>Gerardiana</i>	Above 12,000 feet.
3	" <i>excoles</i> ...	Rai salla ...	" 10,000 feet, Kanol, Painkhanda.
4	<i>Abies Smithiana</i> ...	Rágha ...	Above 10,000 feet.
5	" <i>Brunoniana</i> [dumosa] ...	"	Rare ; found by Captain Webb.
6	<i>Picea Pindrow</i> ...	"	Common above 10,000 feet.
7	" <i>Webbiana</i> ...	"	" " 12,000 feet.
8	<i>Cedrus Deodara</i> ...	Diyár ...	Introduced.
ORDER 3.—CUPRESSINEÆ.			
1	<i>Cupressus torulosa</i> ...	Sarai ...	Naini Tál and Joshimath.
2	" <i>sempervirens</i>	Introduced.
3	<i>Juniperus</i> (<i>Oxycedrus</i>) <i>communis</i> , ...	Chichiya ..	Milam and Nítl, 12,000 feet.
4	" " <i>recurva</i> ...	Bhedara ..	Rare.
	" <i>var. squamata</i> ...	Bel bhedara,	Iláni Páni, 8,000 feet.
5	" (<i>Sabina</i>) <i>excoles</i> ...	Pádám ...	Milam, 12,000 feet.
6	" " <i>Wallichiana</i> ...	Bhedara ...	Milam and Nítl.
	<i>Var. squamosa</i> , Madden ...	Bel bhedara,	Pindari, Painkhanda, Rikholi gudri. A kind of yeast is made from this plant.
ORDER 4.—TAXACEÆ.			
1	<i>Taxus baccata</i> ...	Thuner ...	Common.
2	" <i>nucifera</i>	Rare ; only found by Boyle.

DIVISION V.—ENDOGENÆ.

No.	Scientific name.	Vernacular name.	Note.
ORDER 1.—ZINGIBERACEÆ.			
1	<i>Curcuma angustifolia</i> ...	Ban haldi ...	Up to 6,000 feet; May.
2	" <i>longa</i> [Kuchoor, Boyle] ...	Haldi ...	Cultivated up to 4,000 feet.
3	<i>Hedychium spicatum</i> ...	Kachûr-kachri,	Common.
4	" <i>coccineum</i> ...	Rakt haldi ...	"
5	" <i>villosum</i> [Maiden]	Rare, Piyûra.
6	" <i>tenuiflorum</i> [Murdan Ali]	"
7	" <i>coronarum</i> [flavum, Madden].	...	Gardens.
8	<i>Roseoea purpurea</i>	Common; autumn.
9	" <i>alpina</i>	"
10	" <i>lutea</i> [gracilis]	"
11	" <i>elatior</i>	"
12	<i>Zingiber capitatum</i>	Dakurj b'ndyak.
13	" <i>ligulatum</i>	Bhâbar.
14	" <i>var. chrysanthemum</i>	"
15	" <i>casumunar</i> [elatum, Mad-den?].	...	Bhim Tâl.
16	" <i>officinale</i> ...	Adrak ...	Bhâbar; eaten by pigs.
17	<i>Amomum subulatum</i> ...	Hatchi ...	Cultivated.
18	<i>Costus speciosus</i>	Gardens.
19	<i>Gloiba secunda</i>	Rare; valleys. Autumn.
20	" <i>Orizensis</i> ?	...	Bhâbar.
21	"	...	"
ORDER 2.—MARANTACEÆ.			
1	<i>Canna speciosa</i> ...	Kiwâra ...	Gardens.
ORDER 3.—MUSACEÆ.			
1	<i>Musa sapientum</i>	Cultivated.
2	" <i>nepalensis</i>	River banks, Askot to Dhârchûla.
ORDER 4.—AMARYLLIDACEÆ.			
1	<i>Crinum toxicarium</i> ...	Chandar kan-wal.	Common; July.
2	" <i>speciosum</i>	Introduced.
ORDER 5.—HYPOXIDACEÆ.			
1	<i>Caralligo recurvata</i> ...	Pitâri ...	Bhim Tâl.
2	" <i>orchoides</i> ...	"	Common up to 6,400 feet.
3	<i>Hypoxis minor</i>	Very common.
ORDER 6.—IRIDACEÆ.			
1	<i>Pardanthus sinensis</i> ...	Katâr patâ ...	Very common.
2	<i>Iris nepalensis</i> ...	Nîl-kan wal	Common.
3	" <i>kumaonensis</i>	"
4	" <i>decora</i>	Frequent.
5	<i>Tigridia pavonia</i>	Introduced.

DIVISION V.—ENDOGENÆ—(continued).

No	Scientific name.	Vernacular name.	Note.
ORDER 7.—HYDROCHARIDACE.			
1	<i>Vallisneria spiralis</i>	Bhim Tal.
2	<i>Hydrilla verticillata</i>	"
ORDER 8.—ORCHIDACEÆ.			
TRIBE MALAXIDACE (Kad.)			
SUB-TRIBE PLURITALLACE.			
1	<i>Oncoclis tridifolia</i> ...	Bānda ...	Low elevations.
2	<i>Microstylis Wallichii</i>	Colonel Davidson.
3	<i>Dicella cylindrostachya</i>	Common, Binsar, Chi-na, August.
4	<i>Liparis nepalensis</i>	Lariya Kanta, Colonel Davidson. August.
5	<i>Pholidota articulata</i>	Common, April.
6	<i>Celogyne nuda</i>	Valleys.
7	" <i>præcox</i>	Kotahgar. October.
8	" <i>ovalis</i>	Binsar; common, May.
SUB-TRIBE DENDROBIIUM			
9	<i>Eria convallarioides</i> [<i>Ocrotomeria spicata</i>].	Gara panja,	Common on oaks, June
10	<i>Dendrobium alpestre</i>	Frequent.
11	" <i>Picardii</i> [roseum?]	Common, June.
TRIBE EPIDENDRUM (K.D.)			
12	<i>Apaturia Smithiana</i>	Bhābar.
13	<i>Phaius albus</i>	Very common near Bhim Tal, June.
14	" <i>bicolor</i>	Colonel Davidson.
TRIBE VANDÆ.			
15	<i>Cymbidium triste</i> [<i>Lalsia trichorhiza</i>].	...	Frequent, low levels.
16	<i>Eulophia campestris</i>	Rare Bhābar. Salep made from root.
17	" <i>herbacea</i>	Frequent, sweet scented.
18	<i>Cyrtopora fusca</i> (flava, Royle III.)	Common up to 4,500 feet. May.
19	<i>Vanda cristata</i> ...	Bānda ...	Common, on trees. June.
20	<i>Saccolabium guttatum</i>	Very common, low elevations. May, June.
21	" <i>bapilioeum</i> [<i>Cymbidium premorsum</i> , Roxb.].	...	Haldwani, on <i>Ulmus integrifolia</i> .
22	<i>Aerides affine</i>	Common. June.
23	" <i>odoratum</i> [cornutum, Roxb.].	...	Colonel Davidson.
24	<i>Calanthe plantaginea</i> ...	Hyacinth orchis.	Very common. April.

DIVISION V.—ENDOGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 8.—(ORCHIDACEÆ—(concluded).			
TRIBE ORCHIDÆ.			
26	<i>Orchis lotifolia</i>	Phulasin near Jo-himath, June.
26	<i>Aceras angustifolia</i>	Very common.
27	<i>Platanthera Suannum</i>	Ranigár, September.
28	" <i>obconata</i>	Near Nulni Tal Brow- ery. September.
29	" <i>acuminata</i>	Naini Tal. Coone i Davidson.
30	<i>Hemipilia cordifolia</i>	Near Khairna, Sep- tember.
31	<i>Peristylus goodyeroides</i>	Frequent.
32	<i>Hermidium Monorchis</i>	Top of Lariya Kánta, August.
33	" <i>gramineum</i>	Rare.
34	<i>Habenaria pectinata</i>	Common, September.
35	" <i>internellia</i>	
36	" <i>marginata</i>	Very common, August, September.
37	" <i>ensifolia</i>	Colonel Davidson.
38	" <i>plantaginea</i>	
39	" <i>rostrata</i> , Madden	...	Perhaps same as "emar- ginata. No. 36.
40	<i>Gymnadenia commelynefolia</i> , Mad- den.	...	On Kálmath, 6,000 feet.
41	<i>Satyrium nepalense</i> ...	Pakwa-dinini,	Very common, August, September.
TRIBE GASTRODIEÆ.			
42	<i>Epipogon Gmelini</i>	Colonel Davidson.
TRIBE NLOTILÆ.			
43	<i>Spiranthes australis</i>	Very common, August, September.
44	<i>Epipactis latifolia</i>	Common.
45	" <i>acrostachya</i>	"
46	<i>Goodyera repens</i>	"
47	" <i>procera</i>	Binsar, August.
48	<i>Zeuxine sulcata</i>	Winter, Bhábar.
TRIBE ARTHRODIEÆ.			
49	<i>Cephalanthera acuminata</i>	Common, under trees, May.
TRIBE CYPRIPEDIEÆ.			
50	<i>Cypripedium cordigerum</i>	Lariya Kánta, May. Colonel Davidson.

DIVISION V.—ENDOGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 9.—PALMACEÆ.			
1	<i>Chamerops Martiana</i> ...	Thākīl ...	Frequent.
2	<i>Wallichia densiflora</i> [Harrison oblongifolia.]	Kāla-aunsa...	Common, valleys.
3	<i>Phoenix sylvestris</i> , var. <i>humilis</i> Mad-den.	Khajur ...	Up to 5,000 feet. Bihar.
4	" <i>acaulis</i>	Bhābar.
5	<i>Calamus Royleanus</i> , [C. Rotang?] ...	Ret ...	"
ORDER 10. PONTEDERIACEÆ.			
1	<i>Pontederia (Monochoria) hastata</i> , [P. vaginalis, Madsen.]	...	Common. Rice fields.
ORDER 11.—MELANTHACEÆ.			
1	<i>Diaporium Pitautum</i>	Common.
2	" <i>Leschenaultianum</i> [Lyncharia Leschenaultii, Madsen.]	...	"
3	" <i>Wallichii</i>	Frequent.
4	" <i>Hamiltonianum</i>	"
5	<i>Ipligenia indica</i> [Anguillaria indica, Brown.]	...	Common.
ORDER 12. LILIACEÆ.			
Tribe Tulipeæ			
1	<i>Tulipa stellata</i> ...	Majaula, nalku,	Common up to 7,000.
2	<i>Gagea elegans</i>	" above 10,000 feet
3	<i>Lloydia serotina?</i>	Common.
4	" <i>himalense</i>	Above 10,000 feet.
5	<i>Fritillaria verticillata</i>	Phulain, 8,000 feet.
6	" <i>Thomsonianum</i>	Common, April.
7	<i>Lilium Wallichianum</i> ...	Findora	Common, Rāmbagh.
8	" <i>nepalense</i>	"
9	" <i>giganteum</i> ...	Bhaniya nar,	Above 10,000 feet.
10	<i>Methonica (gloriosa) superba</i>	Bish nāngal,	Valleys.
11	<i>Hemerocallis fulva</i> [diasticha]	...	Above 10,000 feet.
Tribe Alopecæ			
12	<i>Aloe perfoliata</i> ...	Ghikwār ...	Valleys, April.
13	<i>Yucca gloriosa</i> ...	Rāmbāsa ...	Naturalized.
Tribe Hyacintheæ.			
14	<i>Uropetalum</i>	Rare, Kālmāth.
15	<i>Urginea indica</i> [Sella indica, Madsen.]	Gesua ...	Common.
16	<i>Allium leptophyllum</i> ...	Piriyā lāhsan	Outer range.
17	" <i>Wallichii</i> ...	Paḍariyā lāhsan.	"
18	" <i>ellipticum</i> ...	Shcōllā lāhsan	"
19	" <i>Jibba</i> ...	Jibba ...	Above 10,000 feet.
20	" <i>Argand</i> ...	Argand ...	Rānīpāni, 8,000 feet.

DIVISION V.—ENDOGENÆ—(continued).

No.	Scientific name	Vernacular name.	Note.
ORDER 12.—LILIACEÆ (concluded).			
TRIBE ASPARAGACEÆ.			
21	<i>Asparagus racemosus</i>	Ailora ...	Bhábhar.
22	" <i>filicinna</i>	Khalruwa ...	Common, edible.
23	" <i>adscendens</i>	Common.
24	<i>Polygonatum verticillatum</i>	"
25	" <i>multiflorum</i>	"
26	" <i>cirrhiflorum</i> ? (Sibiricum ?)	"
27	<i>Smilacina pallida</i>	"
28	" <i>alpina</i>	Kedár.
TRIBE OPHIOPOGONACEÆ.			
29	<i>Ophiopogon intermedius</i>	Frequent.
30	" <i>Wallichianus</i> [apicatum, Madden].	...	Common
ORDER 13.—COMMELYNACEÆ.			
SECTION OLIGANDRIÆ.			
1	<i>Commelyna</i> (<i>Encommelyna</i>) <i>salicifolia</i>	Frequent.
2	" <i>bengalensis</i>	"
3	" (<i>Trichyrocarpus</i>) <i>obliqua</i>	Kána ...	Very common.
4	<i>Anellema scapiflorum</i> , [<i>Murdannia scapiflora</i> , Royle].	Common up to 4,000 ft.
5	" <i>lineolatum</i>	Above 5,000 feet.
6	" <i>pratense</i>	Rare.
SECTION HEXANDREÆ.			
7	<i>Dithyrocarpus paniculatus</i> [rufus, Kunth].	Kandera ...	Common.
8	<i>Cyanotis axillaris</i>	Bhábhar.
9	" <i>cristata</i>	"
10	" <i>barbata</i>	Common.
11	" <i>racemosa</i>	Bhábhar.
12	<i>Streptolirion volubile</i>	Rare.
ORDER 14.—ALISMACEÆ.			
1	<i>Sagittaria angustifolia</i>
ORDER 15.—JUNCCEÆ.			
1	<i>Juncus bufonius</i>	Shin ...	Common.
2	" <i>glauus</i>	Frequent.
3	" <i>condensus</i>	"
4	" <i>Wallichiana</i>
ORDER 16.—SMILACEÆ.			
1	<i>Smilax villandii</i> , Royle	Kakar dhára.	Common.
2	" <i>macrophylla</i>	"	Frequent.
3	" <i>acutata</i>	"
4	" <i>vaginata</i>	Rare, China.

DIVISION V.—ENDOGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 17.—DIOSCOREACEÆ.			
1	<i>Dioscorea verticillata</i>	Genti ...	Frequent.
2	" <i>agglutinata</i>	Ta r tagur...	Tubers edible.
3	" <i>pentaphylla</i>	Mugina mu- niya.	Tubers edible.
4	" <i>deltoidea</i>	Gūn ...	Above 6,000 feet.
ORDER 18.—AROLEÆ.			
TRIBE DRACUSCULÆ.			
1	<i>Arisæma speciosum</i>	Sarp ka goga,	Common.
2	" <i>Jacquinii</i>	" ...	Very common.
3	" <i>costatum</i>	" ...	Common.
4	" <i>curvatum</i>	" ...	"
5	<i>Arisæma hastatum</i>	" ...	Common, July.
6	<i>Saurauatum guttatum</i>	" ...	"
7	" <i>punctatum</i> ?	Kāla bank ...	Kāladhūngi.
8	<i>Amorphophallus campanulatus</i> [<i>Arisæma campanulatum</i> , Roxb.]	" ...	Blāhar.
9	<i>Typhonium</i> ?	" ...	Common, July, very fetid.
TRIBE CALADIÆ.			
10	<i>Remusatia vivipara</i>	Bagh pin lālu,	Up to 4,500 feet.
11	" <i>capillifera</i> [<i>Caladium pu- miliun</i> , Don ?]	Bānj pindilo,	Above 5,000 feet, flow- ers, June.
12	<i>Colocasia Roxburghii</i>	Jangli gwa Gar pāpar.	Very common.
13	" <i>himalensis</i>	Gwin and pā- par.	Cultivated.
ORDER 19.—ORONTIACEÆ.			
1	<i>Acorus calamus</i>	Bach bij ...	Rare.
2	<i>Scindapsus officinalis</i> , <i>Pothos offici- nalis</i> , Roxb.]	Hathungliya,	Near Bhim Tāl.
3	" <i>decurvus</i> [<i>Pothos decur- sus</i>].	Kelauniya ...	Binsar.
4	<i>Pothos scanlonii</i>	" ...	"
ORDER 20.—TYPHACEÆ.			
1	<i>Typha elephantina</i>	Pādara, pe- tāra.	Blāhar.
2	" <i>angustifolia</i>	Bora ...	Marahes.
ORDER 21.—JUSGAGINACEÆ.			
1	<i>Potamogeton natans</i>	" ...	Bhim Tāl and Naini Tāl.
2	" <i>macrocarpum</i>	" ...	" "
3	" <i>crispum</i>	" ...	" "
4	" <i>pectinatum</i>	" ...	" "
5	" <i>perfoliatum</i>	" ...	" "

DIVISION V.—ENDOGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 22.—JENNAKÆ.			
1	<i>Jenna minor</i>	Bhitar.
2	" <i>polyrhiza</i> [orbiculata, Roxb.]	Sireams, up to 4,000 feet.
ORDER 23.—GRAMINEÆ.			
TRIBE ORYZÆ.			
1	<i>Oryza sativa</i> ...	Dhan	Cultivated.
TRIBE PHALARIDÆ.			
2	<i>Zea Mays</i> ...	Bhuta	Cultivated.
3	<i>Colo. Lacryma</i> ...	Lakochiya	Common.
4	" <i>gigantea</i> ...	"	Frequent.
TRIBE PANICÆ.			
5	<i>Paspalum longiflorum</i> ...	Kana	Rice fields.
6	<i>Panicum (Hyptaria) cruciatum</i> [Hoyel, Madden]	Frequent.
	" <i>var. distachya</i>	"
	" <i>octostachya</i>	"
7	<i>Panicum (Orhopogon) acuminatissimum</i>	Colonel Davidson.
8	<i>Panicum (Kehnochoa) co'orum</i> ...	Jang'i mandira	Common.
9	" <i>frumentaceum</i> ...	Jhangora and	Cultivated.
10	" <i>homonymum</i> [chamæraphis, Madden] ...	mandira.	Very common.
11	" (<i>Setaria</i>) <i>glauca</i>	"
12	" <i>Ita'icum</i> ...	Kangni koni,	Cultivated.
13	" (<i>Virgaria</i>) <i>millaceum</i> ...	China	"
14	<i>Pennisetum triflorum</i> ...	Bimaliya	Common.
15	<i>Arundine la setosa</i> [Hirata, Madden]	"
16	" <i>nepalensis</i> [millaris, Madden]	"
17	<i>Thysanotus agrostis</i> ...	Aunsa	Very common.
TRIBE STIPACÆ.			
18	<i>Arachne (Piptathrum) ho'eiformis</i>
19	<i>Aristida cyanantha</i> ...	Bimbiya & karkal sun.	Colonel Davidson. Very common.
TRIBE AGROSTIDÆ.			
20	<i>Vilfa elongata</i> [Sporobolus elongatus, Madden]	Common.
21	<i>Agrostis alba</i>	Colonel Davidson.
22	<i>Muhlenbergia syriatica</i>	Rare.
23	" <i>geniculata</i>	"
24	<i>Polypogon fugax</i> ...	Gingia gliolliya,	Common.
25	<i>Perotis latifolia</i>	"
TRIBE ALUNDINACÆ.			
26	<i>Phragmites Roxburghii</i> ...	Bichra, nal...	Common
27	" <i>Karka</i> ...	Karka	"
28	" <i>nepalensis</i> ...	Nal tura	Bhim Tal.

DIVISION V.—ENDOGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 25.—GRAMINÆ—(continued).			
TRIBE CHLORIDÆ.			
29	<i>Chloris decora</i>	Common.
30	<i>Cynodon dactylon</i>	Dūb ...	Very common.
31	<i>Echinochloa crusgata</i>	Manduwa ...	Cultivated.
32	" <i>indica</i>	Common.
TRIBE AVEENÆ.			
33	<i>Avena sativa</i>	Jai ...	Cultivated.
34	" <i>fatua</i>	Jai-ata ...	Common.
TRIBE FIETUÆ.			
35	<i>Poa annua</i>	Introduced.
36	" <i>bulbon</i>	"
37	" <i>trivialis</i>	"
38	<i>Eragrostis cynosuroides</i>	Common.
39	" <i>caudata</i>	"
40	" <i>nutans</i>	Col. Davidson.
41	" <i>ambigua</i>	"
42	" <i>ciliata</i>	Common.
43	" <i>plumosa</i>	Very common.
44	" <i>diarrhena</i>	Col. Davidson.
45	" <i>nardoides</i>	Common.
46	" <i>orientalis</i>	October.
47	<i>Coeleria cristata</i>	Very common, April.
48	<i>Dactylis glomerata</i>	Frequent.
49	<i>Festuca (Brachypodium) nepalensis</i> ,	Common.
50	<i>Tripsacum (Pagiolytrum) filiforme</i> ,	"
TRIBE BAMBUSÆ.			
51	<i>Arundinaria falcata</i>	Ringāl and Ningāl.	Very common.
	(<i>Two varieties</i>)	"
52	<i>Thamnochlamys spathuliformis</i>	Ringāl ...	Dudatol.
53	" <i>Falconeri</i>	Kālī ringāl ...	Common, May.
54	<i>Dendrocalamus strictus</i>	Bāns ...	Bhābar.
55	" <i>sp.</i>	" ...	Dhamas; cultivated.
TRIBE HORDEACEÆ.			
56	<i>Lolium temulentum</i>	Bhābar.
57	<i>Triticum vulgare</i>	Lāl gehūn ...	Cultivated.
58	" <i>striatum</i>	Rare.
59	<i>Hordeum hexastichum</i>	Jao ...	Cultivated.
60	" <i>himalayense</i> [cokate]	Us jao ...	" above 10,000 ft.
TRIBE ROTTHOLLIACEÆ.			
61	<i>Ophurus perforatus</i>	Very common.
62	<i>Rottholliola exaltata</i>	Frequent.
63	" <i>hirsuta</i>	Rare.
64	" <i>lavis</i> [<i>Mucrothia lavis</i>],	"
65	<i>Manisuris granularis</i>	Frequent.

DIVISION V.—ENDOGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 23.—GRAMINÆ—(concluded).			
Tribe ANDROPOGONÆ.			
ANDROPOGON, WITH NUMBER IN STEUDL.			
66	Heteropogon contortus, 37	Kumariya	Common.
67	" polystachyus, 43	"	"
68	Ischnum corollatus, 65	Noniya	Frequent.
69	Ischnum speciosus, 132	"	"
70	Spodiopogon involutus, 117	"	"
71	Apocopia himalayensis, 168	"	Rare.
72	Bathratherum lancifolius, 237	"	Common.
73	" micans, 239	"	"
Tribe ANDROPOGONÆ.			
74	Bathratherum subpatens, 240	"	Common.
75	Cymbopogon pachnodes, 297	"	"
76	" distans, 298	"	"
77	" versicolor, 301	Piriya	Very common. Naini Tal
78	" pendulus, 303	"	Common.
79	" Iwarankusa, 306	Miriya	"
80	" Martini, 309	"	Bhābar.
81	" flexuosa, 310	Bujara	Very common A mora.
82	Andropogon erythrocoleus, 341	"	Common.
83	" punctatus, 343	Pagrimalai	Very common.
84	" controversus, 349	"	Very common. October.
85	" montanus, 352	"	Colonel Davidson.
86	Sorghum halepensis, 384	"	Common.
87	Anatherum muricatus, 393	Gandar, kaskas,	"
88	Chrysopogon caryoleus, 404	Gweriya, par-mal.	"
89	" Trinii, 408	"	Rare.
90	" Royleanus, 425	Salini	Common.
91	" echinulatus, 426	"	"
92	" villisulcus, 427	"	"
93	" glaucopis, 429	"	Rare.
94	" subrepens, 430	"	"
95	" parvispica [Rhapis microstachya], 431	Palkiya, chirsula	Very common.
96	Spodiopogon petiolaris, 443	"	Common, Naini Tal, July.
97	Anthistria arundinacea	Ullu, kanyur,	Bhābar and valleys.
98	" onathera	Jyotish-mati,	Roots luminous.
99	Imperata arundinacea [cylindrica Lam],	Shiro	"
100	Saccharum spontaneum	Jhansh	"
101	" Sara	Sarur	Bhābar.
102	" semidecumbens	Tat, mora	"
103	" exaltatum	"	"
104	" Munja	Munj	"
105	Erianthus olivaceus, Edw.	Kans	Common.
106	" vulpinus, Edw.	"	"
107	" filifolius, Nees.	"	"
108	" rufum, Nees.	"	"
109	" rufipilus, Steud.	"	"
110	Leptatherum Royleanum	"	Rare.
111	Pollinis, sp.	"	Frequent.
112	Kalalia japonica	"	"

DIVISION V.—ENDOGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 21.—CYPERACEÆ.			
TRIBE CYPEREÆ.			
1	<i>Cyperus (Pycnos) capillaris</i> ?
2	" <i>diaphanus</i> ?
3	" <i>angulatus</i>
4	" <i>alopecuroides</i> ?
5	" (<i>Eucyperus</i>) <i>compressus</i>
6	" <i>difformis</i>
7	" <i>Iris</i>	...	Common.
8	<i>Cyperus niveus</i>	...	Very common.
9	" <i>tegetiformis</i>	Motha ...	Cultivated.
10	" <i>tenuliformis</i>	...	Common, August
11	" <i>rotundus</i> [<i>hexastachyos</i> , Kortb.]	...	"
12	" <i>obliquus</i>	...	"
13	" <i>pusillus</i> [<i>pygmaeus</i> , Retz.]
14	" <i>Wallichianus</i>
15	" <i>flabriatus</i>
16	" <i>dilatatus</i>	...	Very common.
17	<i>Mariscus cyperinus</i>	Pan motha and nagar-mutha.	Common.
18	<i>Kyllingia monocephala</i>	Nirbisi	"
19	" <i>triceps</i>	"	"
TRIBE SCIRPEÆ.			
20	<i>Abelgardia monostachya</i>
21	<i>Eleocharis palustris</i>
22	" <i>uniglumis</i>
23	<i>Eleocharis ovata</i>
24	" <i>capitata</i>
25	<i>Scirpus juncoides</i> [<i>muticus</i>]
26	" <i>lacustris</i>	...	Bhim Tal.
27	" <i>affinis</i> [<i>maritimus</i>]
28	<i>Isalepis setacea</i>
29	" <i>aripina</i>
30	" <i>Roylei</i>
31	" <i>trifida</i>	...	Common.
32	<i>Fimbristylis pollicacea</i>
33	" <i>junciformis</i>	...	Very common, August.
34	" <i>suberistachya</i> [<i>crassius</i>]	...	Common.
35	" <i>quinquangularis</i>	...	"
36	<i>Eriophorum comosum</i>	Babar	Very common, au- tumn.
TRIBE HYPOCYPEREÆ.			
37	<i>Lipocarpus argentea</i>	...	Very common, July and August.

DIVISION V.—ENDOGENÆ—(concluded).

No.	Scientific name.	Vernacular name.	Note.
ORDER 24.—CYPERACEÆ—(concluded).			
Tribe RHYNCHOSPHEREÆ.			
38	Rhynchospora chinensis
39	" Wallichiana
Tribe CARICEÆ.			
40	Carex (Androgynæ) longipes
41	" foliosa
42	" nubilosa	Common.
43	" notoleia
44	" Vahlil [alpina]
45	" hæmatostoma
46	" setigera
47	" myosurus	Above 5,000 feet, May.
48	" condensata	Common.
49	" filicina	"
50	" (Biddæ) saxatilis [rigida, Good].
51	Carex notha
52	" acuta [gracilis, Curtis]
53	" (Trifidæ), ligulata
54	" cardiopsis
55	" Royleana
56	" Moorcroftii
57	" emodorum [alopecuroides Don],
58	" obscura
59	" fusiformis
60	" Wallichiana
61	" cruenta
62	" (Incerta) Thomsonii, Boott.,
63	" mitis, Boott.
64	" uncinoides, Boott.
65	" vesiculosa, Boott.
66	Uncinia nepalensis
ORDER 25.—ERIOCAULONÆÆ.			
1	Eriocaulon sexangulare	Very common, swamps.
ORDER 26.—TRILLIACEÆ.			
1	Paris polyphylla	Common.
2	Trillium Govanlanum, Boyle	Kulāra Pass, 11,000 feet.

DIVISION VI.—ACROGENÆ.

No.	Scientific name.	Vernacular name.	Note.
ORDER I.—FILICES.			
TRIBE GLEICHENIACEÆ.			
1	<i>Gleichenia dichotoma</i>	Common.
TRIBE DICKSONIACEÆ.			
2	<i>Woodsia lanosa</i>	Rare, Dudatoli.
3	" <i>elongata</i>	Common.
4	<i>Dicksonia appendiculata</i> [<i>Sitobium</i> , J. Smith].	...	"
TRIBE HYMENOPHYLLACEÆ.			
5	<i>Hymenophyllum poyanthus</i> [exacrum?]	...	On trees, Binsar, October. J. H. Batten.
TRIBE DAVALLIEÆ.			
6	<i>Davallia</i> (<i>Leucostegia</i>) <i>membranulosa</i> ,	...	Frequent.
7	" " <i>immixta</i>	"
8	" " <i>multidentata?</i> <i>pulchra?</i>	...	Rare.
9	" " <i>chaerophylla</i> [<i>cystopteris</i> , D. C.]	...	Common, Binsar.
10	" " <i>nodosa</i> [<i>Acrophorus</i> Presl.]	...	Rare.
11	" (<i>Endavallia</i>) <i>divaricata</i> (<i>cligana</i> , Madden).	...	Very common.
12	" (<i>Microlepia</i>) <i>hookeriana</i>	Bhábar.
13	" " <i>spelunce</i>	"
14	" (<i>Stenoloma</i>) <i>tenuifolia</i>	Common.
TRIBE PTERIDACEÆ.			
15	<i>Adiantum lunulatum</i>	Common.
16	" <i>caudatum</i> [<i>rhizophorum</i> Swartz].	...	Very common.
17	" <i>Capillus-veneris</i> ...	Mubiraka ...	Common.
18	" <i>venustum</i> ...	Hansráj ...	"
19	" <i>pedatum</i>	"
20	<i>Cheilanthes Dalhousii</i>	"
21	" <i>farinosa</i> [<i>dealbata</i>	Very common.
22	<i>Onychium auratum</i>	Common.
23	" <i>japonicum</i>	"
24	<i>Pellaea</i> [<i>Chelolepion</i>] <i>gracilis</i>	Rare; resembles <i>Cryptogramma crispa</i> .
25	" [<i>Allosorus nitidula</i>	Rare.
26	" [<i>Platyloma</i>] <i>calomelanos</i> ...	Bish kutra...	On rocks at low levels.
27	<i>Pteris</i> (<i>Eupteria</i>) <i>longifolia</i>	Very common.
28	" " <i>erecta</i>	"
29	" " <i>var. stenophylla</i>	Common.
30	" " <i>quadri aurita</i>	Very common.
31	" (<i>Pandia</i>) <i>aquilina</i>	Very common.
32	" (<i>Campteria</i>) <i>blanckia</i>	Rare.
33	" " <i>Wallichiana</i>	Common.
34	" (<i>Litobrochia</i>) <i>incisa</i>	Near Raughár, J. H. Batten.

DIVISION VI.—ACROGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 1.—FILICES—(continued).			
TRIBE BLECHNEÆ.			
34	<i>Woodwardia radicans</i>	Low levels, very common.
TRIBE ASPLENEÆ.			
35	<i>Asplenium (Eurasplenium) ensiforme</i>	Common. Binsar, July.
36	" " <i>alternans</i>	...	Very common on walls.
37	" " <i>viride</i>	Above 12,000 feet.
38	" " <i>trichomanes</i>	...	Common.
39	" " <i>resectum</i>	Frequent.
40	" " <i>planicaule</i>	...	Common.
41	" " <i>adiantum nitidum</i>	Rare.
42	" " <i>fontanum</i> ,	...	Frequent.
43	" " <i>varians</i>	"
44	" " <i>bulbiferum</i> ,	...	Rare.
45	" " <i>tenuifolium</i> [<i>concinnum</i> Wall].	...	"
46	" (<i>Athyrium</i>) <i>thelypteroides</i>	Colonel Davidson.
47	" " <i>macrocarpum</i> [<i>foliosum</i> Wall].	...	Common.
48	" " <i>nigripes</i>	Colonel Davidson.
49	" " <i>filix femina</i> [<i>pectinata</i> Wall].	...	Common.
50	" " <i>oxyphyllum</i>	Colonel Davidson.
51	" " <i>imbriatum</i> ,	...	"
52	" " <i>umbrosum</i> [<i>Allantollia australe</i>]	...	Rare.
53	" (<i>Diplazium</i>) <i>polypodoides</i> .	Lingra ...	Common.
54	" " <i>maximiliana</i>	" ...	" edible.
55	" (<i>Anisogonium</i>) <i>esculentum</i> .	" ...	not edible.
56	" (<i>Hemidictyum</i>) <i>Ceterach</i> [<i>Ceterach</i> Wall].	...	Rare; Very like <i>Asplenium alternans</i> .
57	<i>Actinopteris radiata</i> ...	Morpuchh ...	Common, walls, Srinagar.
TRIBE ASPIDIEÆ.			
58	<i>Aspidium (Polystichum) auriculatum</i> ,	...	Common.
59	" " <i>thelypteris</i>	"
60	" " <i>Thomsoni</i>	Above 8,000 feet.
61	" " <i>aculeatum</i>	Common.
...	" " <i>var. rubro-batium</i> , Wall.	...	Very common.
62	" " <i>Prescottianum</i> ,	...	Colonel Davidson.

DIVISION VI.—ACROGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 1.—FILICES—(continued).			
Tribe ASPIDIÆ—(concluded).			
63	<i>Aspidium (Polystichum) aristatum</i> (confolium Wall).	...	Bhābar.
64	" (<i>Cyrtomium</i>) <i>falcatum</i>	Frequent.
...	" var. <i>caryoti-</i> <i>dium</i>	Very common.
65	<i>Nephrodium (Lastrea) patens</i>	Colonel Davidson.
66	" " <i>canum</i>	"
67	" " <i>prolixum</i>	Common.
68	" " <i>filiæ nia</i>	Very common.
...	" " var. <i>cochle-</i> <i>tum</i>	Common.
69	" " <i>flaccidum</i>	"
70	" " <i>spinulosum</i>	Very common.
71	" " <i>sparsum</i>	Common.
72	" <i>odoratum</i> (<i>erlocarpum</i> , H. K.)	Katra ...	Very common; edible.
73	" " <i>angustifrons</i> ,	...	Frequent.
74	" (<i>Eunephrodium</i>) <i>mitum</i>	"
75	" <i>cucullatum</i> (<i>mitum</i> , Sieb.).	...	Bhābar.
76	" " <i>aridum</i>	Colonel Davidson.
77	" " <i>molle</i>	Frequent.
78	" (<i>Sagenia</i>) <i>polymorphum</i> ...	Lingra ...	Up to 4,000 feet.
79	" " <i>variolosum</i>	Rare.
80	" " <i>cicuta tum</i>	Common.
81	<i>Nephrolepis cordifolia</i> [tuberosa. II. K.]	...	Rare; Pīpa kothi.
82	<i>Olcandra nerifolmis</i>	Common.
83	" <i>Wallichii</i>	"
Tribe POLYPODIEÆ.			
84	<i>Polypodium (Phegopteris) cruce-</i> <i>scaus</i> .	..	Low elevations.
85	" " <i>distant</i>	Common.
86	" (<i>Goniopteris</i>) <i>urophyllum</i> ,	...	"
87	" " <i>proliferum</i> ,	...	Bhābar.
88	" " <i>lineatum</i>	Vallaya.
89	" " <i>multilinea-</i> <i>tum</i>	"
90	" (<i>Goniophlebium</i>) <i>amœnum</i> ,	...	Common.
91	" " <i>lachnopus</i>	"
92	" <i>subauricula-</i> <i>tum (argutum, Wall)</i>	Frequent.
93	<i>Polypodium (Niphobolus) adnascens</i> ,	...	Common.
94	" " <i>lingua</i>	"
95	" " <i>stigmatosum</i> ,	...	Vallaya.
96	" " <i>fissum</i>	Common, June.
97	" " (<i>foeculo-</i> <i>sum</i>).	...	Rare.
98	" (<i>Phymatodes</i>) <i>lineare</i> [<i>Pleopellis nuda</i>].	...	Scented fern; very common on trees.
99	" " <i>normale</i>	Common.

DIVISION VI.—ACROGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 1.—FILICES—(concluded).			
TRIBE POLYPODIUM—(concluded).			
100	<i>Polypodium (Phymatodes) membranaceum.</i>	...	Common.
101	<i>trifidum</i> [<i>oxylobum</i> , Wall].	...	Frequent.
102	" " <i>ma'acodon</i> ,	...	Rare.
103	" " <i>ebcuiques</i>	"
104	" (<i>Drynaria</i>) <i>propinqua</i> ,	...	Common.
105	" " <i>rivale</i>	"
106	" " <i>quercifolium</i>	Bhābar.
107	" " <i>juglandifolium</i> ,	...	Common.
108	<i>himalayense</i> [<i>venustum</i> , Wall].	...	Frequent.
TRIBE GRAMNITIDÆ.			
109	<i>Gymnogramma (Leptogramma) Totta</i> ,	...	Colonel Davidson.
110	" (<i>Egymnogramma</i>) <i>vestita</i> ,	Mouse ear fern.	Pyra and China.
111	" " <i>Andersoni</i> ,	...	About 13,000 feet.
112	" " <i>javanica</i>	Binsar. August.
113	" (<i>Bellignea</i>) <i>lanceolata</i> ,	...	Rare.
114	" " <i>involuta</i>	Common.
115	<i>Vittaria (Euvittaria) elonata</i>	Binsar.
116	" (<i>Tenloopsis</i>) <i>scolopendrina</i> ?	...	Binsar, June.
TRIBE ACROSTICHIDÆ.			
117	<i>Acrostichum virens</i>	Bhābar.
	<i>ser. deltigerum</i> [<i>Menis-</i> <i>cium deltigerum</i> , Wall].	...	"
TRIBE OSMUNDACEÆ.			
118	<i>Osmunda Claytoniana</i> [<i>interrupta</i>]	Very common 10,000 feet Chaupatta.
119	" <i>regalis ser. japonica</i>	Rare Khāti 10,000 feet. May.
TRIBE SCHISTACEÆ.			
120	<i>Schizaea digitata</i>	Common.
121	<i>Lygodium scandens</i> ...	Marora, bandhalya.	Bhābar.
122	" <i>japonicum</i> ...	" ...	Common.
TRIBE MARATTIACEÆ.			
123	<i>Angiopteris erecta</i>	Rare.
TRIBE OPHIOGLOSSACEÆ.			
124	<i>Ophioglossum vulgatum</i>	"
125	<i>Botrychium lunaria</i>	Binsar.
126	" <i>denticolium</i>	Common.
127	" <i>virginianum</i>	Very common.
	<i>ser. lanuginosum</i>	Common.

DIVISION VI.—ACROGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 2.—EQUISETACEÆ.			
1	<i>Equisetum debile</i>	Rhābar.
2	" <i>diffusum</i> , Don [<i>s c o p a-</i> <i>rium</i> , Wall].	...	Illī strōama.
ORDER 3.—LYCOPODIACEÆ.			
1	<i>Lycopodium tenellum</i>	Very common.
2	" <i>setaceum</i>	Rare.
3	" <i>circinatum</i>	Frequent, 4,000 feet.
4	" <i>subulifolium</i> ...	Tula muka...	Common.
ORDER 4.—MARSILIACEÆ.			
1	<i>Marsilia quadrifolia</i>	Rhābar.
ORDER 5.—SALVINIACEÆ.			
1	<i>Asolla pinnata</i> ...	Tarāī ...	Common in pools.
ORDER 6.—CHARACEÆ.			
1	<i>Chara verticillata</i>	Nainī Tāl and Bhīm Tāl.
ORDER 7.—BRYACEÆ.			
TRIBE DICRANACEÆ.			
1	<i>Leptotrichum inclinatum</i> , Mitten.	...	Hoti.
2	" <i>molliculum</i> , Mitten	Common.
3	" <i>himalayanum</i> Mitten.	...	"
4	<i>Dicranum himalayanum</i> , Mitten
5	" <i>reflexifolium</i> , Muller.
6	<i>Didymodon laxifolius</i>
TRIBE LEUCOBRYACEÆ.			
7	<i>Oetoblepharum albidum</i>	Rhābar.
TRIBE TRICHOSTOMACEÆ.			
8	<i>Tortula squarrosa</i>	Common.
9	" <i>anomala</i>
10	<i>Anastangium Roylei</i> , Mitten
11	" <i>Thomsoni</i> , Mitten
12	" <i>Stracheyanum</i> , Mitten

DIVISION VI.—ACROGENÆ—(continued).

No.	Scientific name.	Vernacular name.	Note.
ORDER 7.—BRYACEÆ—(continued).			
TRIBE TRICHOPTOMACEÆ—(concluded).			
13	<i>Hymenostylium aurantiacum</i> , Mitten.
14	" <i>curirostrum</i>
15	<i>Barbula vinella</i>
16	" <i>subpellucida</i> , Mitten
17	" <i>gracilentia</i> , Mitten
18	" <i>nigrescens</i> , Mitten
19	<i>Desmatodon Laureri</i>
20	" <i>Wallichii</i>
21	" <i>involutus</i>
22	<i>Eucalypta ciliata</i>
TRIBE GRIMMIACEÆ.			
23	<i>Grimmia spocarpa</i>	Rocks, common.
24	" (<i>Rhacomitrium</i>) <i>subsecunda</i> ,	" "
25	<i>Glyphomitrium tortula</i> , Mitten
TRIBE ORTHOTRICHACEÆ			
26	<i>Orthotrichum Hookeri</i>
27	<i>Macromitrium Moorcroftii</i>
TRIBE FUNARIACEÆ.			
28	<i>Funaria hygrometrica</i>	Very common, summer.
29	" <i>leptopoda</i>
TRIBE SPLACHNACEÆ.			
30	<i>Tayloria indica</i>
31	" (<i>Disodon</i>) <i>tenella</i> , Mitten,	October.
TRIBE BARTRAMIACEÆ.			
32	<i>Bartramia Halleriana</i>
33	" <i>leptoloma</i>	Near <i>Ithyphylia</i> .
34	" <i>subpellucida</i> , Mitten	Resembles "
35	<i>Philonotis Turneriana</i>	Common.
36	" <i>talenta</i>	"
TRIBE BRYACEÆ.			
37	<i>Mitohoforia himalayana</i>	Pindari, 11,000 feet.
38	<i>Webera elongata</i>
39	<i>Bryum filiforme</i> , Mitten
40	" <i>argenteum</i>	Very common.

